



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 06:17 PM GMT

PDB ID : 1A07
Title : C-SRC (SH2 DOMAIN) COMPLEXED WITH ACE-MALONYL TYR-GLU
-(N,N-DIPENTYL AMINE)
Authors : Shewchuk, L.; Jordan, S.
Deposited on : 1997-12-09
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	107	<div> <div></div> <div>73% 23% ..</div> </div>
1	B	107	<div> <div>3%</div> <div>81% 13% ..</div> </div>
2	C	4	<div> <div>75% 25%</div> </div>
2	D	4	<div> <div>75% 25%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2282 atoms, of which 507 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called C-SRC TYROSINE KINASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	105	Total	C	H	N	O	S	0	0	0
			1040	528	205	150	154	3			
1	B	104	Total	C	H	N	O	S	0	0	0
			1020	519	198	147	153	3			

- Molecule 2 is a protein called ACE-MALONYL TYR-GLU-(N,N-DIPENTYL AMINE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	4	Total	C	H	N	O	0	0	0
			36	26	2	3	5			
2	D	4	Total	C	H	N	O	0	0	0
			36	26	2	3	5			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	25	Total	H	O	0	0
			75	50	25		
3	B	20	Total	H	O	0	0
			60	40	20		
3	C	4	Total	H	O	0	0
			12	8	4		
3	D	1	Total	H	O	0	0
			3	2	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: C-SRC TYROSINE KINASE




- Molecule 1: C-SRC TYROSINE KINASE



- Molecule 2: ACE-MALONYL TYR-GLU-(N,N-DIPENTYL AMINE)



- Molecule 2: ACE-MALONYL TYR-GLU-(N,N-DIPENTYL AMINE)



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.50Å 67.20Å 74.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 2.20 41.37 – 2.20	Depositor EDS
% Data completeness (in resolution range)	85.0 (6.00-2.20) 84.7 (41.37-2.20)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.04 (at 2.20Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.192 , (Not available) 0.184 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	31.5	Xtriage
Anisotropy	0.212	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 77.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 11800 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2282	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DIP, PTR, ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.96	0/852	1.68	16/1148 (1.4%)
1	B	1.02	2/839 (0.2%)	1.61	13/1133 (1.1%)
2	C	0.96	0/8	0.87	0/9
2	D	1.16	0/8	0.67	0/9
All	All	0.99	2/1707 (0.1%)	1.64	29/2299 (1.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	163	ARG	CZ-NH1	10.15	1.46	1.33
1	B	163	ARG	CZ-NH2	6.77	1.41	1.33

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	158	ARG	NE-CZ-NH2	-11.01	114.80	120.30
1	B	163	ARG	CD-NE-CZ	-9.81	109.86	123.60
1	A	151	TRP	CD1-CG-CD2	9.27	113.72	106.30
1	A	151	TRP	CE2-CD2-CG	-8.63	100.40	107.30
1	B	159	ARG	NE-CZ-NH2	8.51	124.55	120.30
1	B	208	ARG	NE-CZ-NH2	-8.19	116.21	120.30
1	A	188	CYS	N-CA-CB	-8.07	96.07	110.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	205	TYR	CB-CG-CD2	-7.20	116.68	121.00
1	A	220	ARG	NE-CZ-NH2	-7.17	116.71	120.30
1	B	151	TRP	CD1-CG-CD2	7.16	112.03	106.30
1	B	151	TRP	CE2-CD2-CG	-7.02	101.68	107.30
1	B	205	TYR	CB-CG-CD1	-6.99	116.81	121.00
1	B	151	TRP	CG-CD2-CE3	6.34	139.60	133.90
1	A	151	TRP	CB-CG-CD1	-6.27	118.85	127.00
1	A	178	ARG	NE-CZ-NH2	-6.25	117.17	120.30
1	B	151	TRP	CB-CG-CD1	-6.20	118.94	127.00
1	A	151	TRP	CG-CD1-NE1	-5.99	104.11	110.10
1	B	158	ARG	NE-CZ-NH2	5.83	123.22	120.30
1	A	151	TRP	CG-CD2-CE3	5.81	139.13	133.90
1	A	158	ARG	NE-CZ-NH1	5.63	123.12	120.30
1	B	163	ARG	CB-CG-CD	-5.50	97.29	111.60
1	A	188	CYS	CB-CA-C	5.45	121.29	110.40
1	B	181	GLU	CA-CB-CG	5.39	125.26	113.40
1	A	152	TYR	CB-CG-CD1	-5.31	117.81	121.00
1	A	203	LYS	CG-CD-CE	-5.28	96.05	111.90
1	A	176	LEU	CA-CB-CG	5.10	127.02	115.30
1	A	163	ARG	NE-CZ-NH2	-5.08	117.76	120.30
1	B	151	TRP	CG-CD1-NE1	-5.08	105.02	110.10
1	B	200	LEU	CB-CG-CD1	-5.01	102.48	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	160	GLU	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	835	205	813	12	0
1	B	822	198	789	8	0
2	C	34	2	38	1	0
2	D	34	2	38	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	25	50	0	0	0
3	B	20	40	0	0	0
3	C	4	8	0	0	0
3	D	1	2	0	0	0
All	All	1775	507	1678	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:146:ILE:HD13	1:B:226:LEU:HD12	1.72	0.70
1:A:242:HIS:HD2	1:A:243:ARG:O	1.77	0.68
1:B:159:ARG:HD2	1:B:163:ARG:HH21	1.66	0.60
1:B:159:ARG:HD2	1:B:163:ARG:NH2	2.18	0.59
1:A:209:LYS:HE3	1:A:213:GLY:O	2.06	0.55
1:B:242:HIS:HD2	1:B:243:ARG:O	1.92	0.53
1:A:188:CYS:SG	2:C:101:PTR:HE2	2.50	0.51
1:A:216:TYR:HB3	1:A:222:GLN:HG3	1.92	0.51
1:A:146:ILE:HD11	1:A:151:TRP:CH2	2.46	0.50
1:B:174:THR:HA	1:B:246:THR:O	2.13	0.48
1:A:233:TYR:HA	1:A:236:HIS:O	2.15	0.46
1:B:158:ARG:HH22	2:D:100:ACE:C	2.19	0.46
1:A:209:LYS:HD2	1:A:214:GLY:O	2.15	0.46
1:A:153:PHE:O	1:A:156:ILE:HG23	2.16	0.45
1:A:159:ARG:HH21	1:B:150:GLU:CD	2.22	0.43
1:A:182:THR:HG21	1:B:163:ARG:HB3	2.01	0.42
1:A:146:ILE:HA	1:A:149:GLU:HG2	2.03	0.41
1:A:206:LYS:HB2	1:A:206:LYS:HE3	1.92	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	103/107 (96%)	99 (96%)	4 (4%)	0	100	100
1	B	102/107 (95%)	97 (95%)	5 (5%)	0	100	100
2	C	1/4 (25%)	1 (100%)	0	0	100	100
2	D	1/4 (25%)	1 (100%)	0	0	100	100
All	All	207/222 (93%)	198 (96%)	9 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	87/94 (93%)	86 (99%)	1 (1%)	80	89
1	B	85/94 (90%)	84 (99%)	1 (1%)	78	88
2	C	1/1 (100%)	1 (100%)	0	100	100
2	D	1/1 (100%)	1 (100%)	0	100	100
All	All	174/190 (92%)	172 (99%)	2 (1%)	80	89

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	201	ASN
1	B	227	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	222	GLN
1	A	242	HIS
1	B	227	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	242	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PTR	C	101	2	10,11,17	1.01	1 (10%)	10,13,24	1.07	1 (10%)
2	PTR	D	101	2	10,11,17	0.75	0	10,13,24	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PTR	C	101	2	-	0/4/6/13	0/1/1/1
2	PTR	D	101	2	-	0/4/6/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	101	PTR	CB-CA	-2.28	1.48	1.53

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	101	PTR	CD2-CG-CD1	2.06	121.43	118.13

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	101	PTR	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	105/107 (98%)	-0.16	1 (0%) 84 83	13, 26, 57, 78	0
1	B	104/107 (97%)	-0.25	3 (2%) 55 54	15, 26, 65, 77	0
2	C	1/4 (25%)	-0.52	0 100 100	32, 32, 32, 32	0
2	D	1/4 (25%)	-0.90	0 100 100	28, 28, 28, 28	0
All	All	211/222 (95%)	-0.21	4 (1%) 70 68	13, 26, 65, 78	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	167	ASN	2.6
1	B	183	THR	2.6
1	B	168	ALA	2.5
1	A	198	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	PTR	C	101	11/17	0.93	0.12	-	0,28,34,35	0
2	PTR	D	101	11/17	0.92	0.12	-	0,29,36,38	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.