



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:38 PM GMT

PDB ID : 4A2E  
Title : CRYSTAL STRUCTURE OF A CORIOLOPSIS GALLICA LACCASE AT  
1.7 Å RESOLUTION PH 5.5  
Authors : De La Mora, E.; Valderrama, B.; Horjales, E.; Rudino-Pinera, E.  
Deposited on : 2011-09-26  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	496	<div> <div></div> <div>87%</div> <div>12%</div> <div>.</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4466 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LACCASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	496	Total	C	N	O	S	0	10	0
			3800	2411	640	740	9			

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	39	ASP	TYR	CONFLICT	UNP Q1W6B1
A	151	ASN	GLN	CONFLICT	UNP Q1W6B1
A	178	LYS	ARG	CONFLICT	UNP Q1W6B1
A	182	PRO	ALA	CONFLICT	UNP Q1W6B1
A	183	VAL	ILE	CONFLICT	UNP Q1W6B1
A	229	TYR	HIS	CONFLICT	UNP Q1W6B1
A	256	LEU	ILE	CONFLICT	UNP Q1W6B1
A	287	THR	ASN	CONFLICT	UNP Q1W6B1
A	288	GLN	THR	CONFLICT	UNP Q1W6B1
A	294	THR	VAL	CONFLICT	UNP Q1W6B1
A	314	THR	ALA	CONFLICT	UNP Q1W6B1
A	329	LYS	GLU	CONFLICT	UNP Q1W6B1
A	356	ASN	ARG	CONFLICT	UNP Q1W6B1
A	358	THR	SER	CONFLICT	UNP Q1W6B1
A	423	VAL	ALA	CONFLICT	UNP Q1W6B1

- Molecule 2 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Cu	0	0
			4	4		

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	2	Total	C	N	O	0	0
			28	16	2	10		
3	A	2	Total	C	N	O	0	0
			28	16	2	10		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	606	Total	O	0	0
			606	606		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: LACCASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.38 Å 86.13 Å 152.64 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.63 – 1.80 25.63 – 1.80	Depositor EDS
% Data completeness (in resolution range)	96.6 (25.63-1.80) 96.6 (25.63-1.80)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.88 (at 1.80 Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, $R_{free}$	0.154 , 0.187 0.154 , 0.188	Depositor DCC
$R_{free}$ test set	3389 reflections (5.30%)	DCC
Wilson B-factor (Å <sup>2</sup> )	16.4	Xtriage
Anisotropy	0.110	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 50.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 67354 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4466	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, CU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.48	18/3925 (0.5%)	1.13	16/5398 (0.3%)

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	221	VAL	CB-CG1	-8.43	1.35	1.52
1	A	517	GLN	C-OXT	7.86	1.38	1.23
1	A	464	PRO	CA-CB	7.59	1.68	1.53
1	A	301	TYR	CD1-CE1	7.24	1.50	1.39
1	A	423	VAL	CB-CG2	-7.09	1.38	1.52
1	A	475	PHE	CE2-CZ	6.33	1.49	1.37
1	A	245	SER	CB-OG	6.08	1.50	1.42
1	A	160	VAL	CB-CG2	6.06	1.65	1.52
1	A	329	LYS	CD-CE	5.58	1.65	1.51
1	A	467	TRP	CE3-CZ3	5.56	1.48	1.38
1	A	148	TYR	CE1-CZ	5.55	1.45	1.38
1	A	120	VAL	CB-CG1	5.52	1.64	1.52
1	A	437	TYR	CE1-CZ	5.21	1.45	1.38
1	A	114	PHE	CE2-CZ	5.20	1.47	1.37
1	A	118	PHE	CD2-CE2	5.16	1.49	1.39
1	A	229	TYR	CG-CD2	5.16	1.45	1.39
1	A	242	GLU	CD-OE1	5.11	1.31	1.25
1	A	329	LYS	CE-NZ	5.10	1.61	1.49

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	282	LEU	CB-CG-CD1	7.20	123.23	111.00
1	A	485	MET	CG-SD-CE	-6.92	89.14	100.20
1	A	271	ASP	CB-CG-OD1	6.69	124.32	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	142	ARG	NE-CZ-NH2	-6.40	117.10	120.30
1	A	371	VAL	CG1-CB-CG2	-6.33	100.78	110.90
1	A	343[A]	ASP	CB-CG-OD2	6.05	123.75	118.30
1	A	343[B]	ASP	CB-CG-OD2	6.05	123.75	118.30
1	A	275	ASP	CB-CG-OD1	5.95	123.66	118.30
1	A	248	LEU	CB-CG-CD1	-5.91	100.95	111.00
1	A	142	ARG	NE-CZ-NH1	5.86	123.23	120.30
1	A	460	ARG	NE-CZ-NH2	-5.71	117.44	120.30
1	A	216	TYR	CZ-CE2-CD2	-5.55	114.81	119.80
1	A	80	LYS	CD-CE-NZ	-5.42	99.24	111.70
1	A	64	ARG	NE-CZ-NH1	5.40	123.00	120.30
1	A	329	LYS	CD-CE-NZ	5.10	123.43	111.70
1	A	212	LYS	CD-CE-NZ	-5.05	100.08	111.70

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3800	0	3601	33	0
2	A	4	0	0	0	0
3	A	56	0	50	0	0
4	A	606	0	0	6	0
All	All	4466	0	3651	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:418[B]:HIS:HE1	1:A:470:HIS:CE1	1.36	1.41
1:A:418[B]:HIS:CE1	1:A:470:HIS:CE1	2.09	1.40
1:A:329:LYS:HE2	1:A:329:LYS:H	1.13	1.08

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:418[B]:HIS:CE1	1:A:470:HIS:NE2	2.32	0.97
1:A:418[B]:HIS:HE1	1:A:470:HIS:HE1	1.09	0.89
1:A:199:ALA:HB2	4:A:2317:HOH:O	1.79	0.82
1:A:329:LYS:HE2	1:A:329:LYS:N	1.97	0.78
1:A:422:HIS:HE1	1:A:487:GLU:OE1	1.71	0.73
1:A:22:ALA:HB2	1:A:163:ASP:OD2	1.89	0.72
1:A:276:ASN:HD21	1:A:300:ARG:HE	1.38	0.71
1:A:418[B]:HIS:CE1	1:A:472:HIS:CD2	2.83	0.67
1:A:284:ASN:HD22	1:A:284:ASN:H	1.45	0.65
1:A:329:LYS:CE	1:A:329:LYS:H	2.03	0.62
1:A:276:ASN:ND2	1:A:300:ARG:HE	1.97	0.61
1:A:418[B]:HIS:NE2	1:A:470:HIS:NE2	2.48	0.60
1:A:422:HIS:CE1	1:A:487:GLU:OE1	2.54	0.60
1:A:251:HIS:HE1	4:A:2426:HOH:O	1.92	0.52
1:A:214:LYS:HE2	4:A:2257:HOH:O	2.09	0.52
1:A:33:ASN:ND2	1:A:74:THR:H	2.09	0.51
1:A:135:THR:HA	1:A:474:ASP:OD2	2.09	0.50
1:A:242:GLU:HB3	1:A:265:SER:HB2	1.94	0.49
1:A:199:ALA:CB	4:A:2317:HOH:O	2.47	0.48
1:A:33:ASN:HD22	1:A:74:THR:H	1.62	0.47
1:A:413:PHE:HB2	1:A:414:PRO:HA	1.94	0.47
1:A:418[B]:HIS:ND1	1:A:472:HIS:CD2	2.82	0.47
1:A:436:ASN:HD21	1:A:439:ASN:ND2	2.13	0.47
1:A:92:HIS:HD2	4:A:2147:HOH:O	1.99	0.46
1:A:227:PRO:HA	1:A:284:ASN:HD21	1.82	0.45
1:A:294:THR:CG2	4:A:2301:HOH:O	2.66	0.43
1:A:418[B]:HIS:ND1	1:A:472:HIS:HD2	2.16	0.43
1:A:418[B]:HIS:HB3	1:A:445:VAL:HG22	2.01	0.43
1:A:91:GLN:HE22	1:A:118:PHE:CB	2.33	0.41
1:A:208:ILE:HD13	1:A:279:ILE:CD1	2.51	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	504/496 (102%)	487 (97%)	15 (3%)	2 (0%)	39 23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	226	ASP
1	A	79	LEU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	411/401 (102%)	402 (98%)	9 (2%)	60 45

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	263	ARG
1	A	269	ASN
1	A	282	LEU
1	A	284	ASN
1	A	329	LYS
1	A	407	THR
1	A	439	ASN
1	A	468	PHE
1	A	481	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	ASN
1	A	91	GLN
1	A	92	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	104	ASN
1	A	105	GLN
1	A	154	HIS
1	A	251	HIS
1	A	269	ASN
1	A	272	GLN
1	A	276	ASN
1	A	284	ASN
1	A	356	ASN
1	A	380	GLN
1	A	422	HIS
1	A	439	ASN
1	A	517	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

4 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	A	1522	1,3	14,14,15	1.66	2 (14%)	15,19,21	1.41	3 (20%)
3	NAG	A	1523	3	14,14,15	1.48	2 (14%)	15,19,21	1.56	3 (20%)
3	NAG	A	1524	1,3	14,14,15	1.63	2 (14%)	15,19,21	1.97	4 (26%)
3	NAG	A	1525	3	14,14,15	1.14	0	15,19,21	1.38	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1522	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1523	3	-	0/6/23/26	0/1/1/1
3	NAG	A	1524	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1525	3	-	0/6/23/26	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1524	NAG	C2-N2	2.21	1.50	1.46
3	A	1523	NAG	O4-C4	2.56	1.49	1.43
3	A	1523	NAG	C3-C2	2.67	1.58	1.52
3	A	1522	NAG	C4-C5	2.97	1.59	1.53
3	A	1524	NAG	C1-C2	4.86	1.59	1.52
3	A	1522	NAG	C1-C2	4.93	1.59	1.52

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1524	NAG	C2-N2-C7	-4.42	117.36	123.04
3	A	1525	NAG	O6-C6-C5	-3.22	100.70	111.33
3	A	1524	NAG	O7-C7-C8	-2.76	116.99	122.06
3	A	1523	NAG	O6-C6-C5	-2.31	103.72	111.33
3	A	1522	NAG	O4-C4-C5	-2.22	103.37	109.24
3	A	1525	NAG	O3-C3-C2	-2.19	104.77	109.11
3	A	1522	NAG	O7-C7-C8	-2.12	118.17	122.06
3	A	1524	NAG	O4-C4-C3	-2.09	105.64	110.34
3	A	1523	NAG	C3-C4-C5	-2.07	106.58	110.20
3	A	1525	NAG	C1-O5-C5	2.20	115.04	112.25
3	A	1523	NAG	O7-C7-N2	2.46	126.88	121.86
3	A	1522	NAG	O5-C5-C6	2.77	113.33	107.35
3	A	1524	NAG	O7-C7-N2	3.40	128.80	121.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.6 Ligand geometry

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	496/496 (100%)	-0.30	5 (1%) 84 82	8, 13, 22, 34	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	354	GLY	2.9
1	A	353	ALA	2.3
1	A	317	THR	2.1
1	A	314	THR	2.1
1	A	22	ALA	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	A	1523	14/15	0.90	0.14	1.96	10,17,26,27	0
3	NAG	A	1524	14/15	0.94	0.10	0.29	12,17,26,28	0
3	NAG	A	1522	14/15	0.97	0.07	-0.64	12,15,19,20	0
3	NAG	A	1525	14/15	0.84	0.23	-	16,27,31,37	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CU	A	1521	1/1	0.99	0.03	-2.33	12,12,12,12	1
2	CU	A	1518	1/1	1.00	0.03	-4.26	14,14,14,14	1
2	CU	A	1520	1/1	1.00	0.02	-4.34	17,17,17,17	1
2	CU	A	1519	1/1	1.00	0.03	-5.90	12,12,12,12	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.