



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:40 PM GMT

PDB ID : 4A5O  
Title : Crystal structure of Pseudomonas aeruginosa N5, N10- methylenetetrahydrofolate dehydrogenase-cyclohydrolase (FolD)  
Authors : Eadsforth, T.C.; Gardiner, M.; Maluf, F.V.; Mcelroy, S.; James, D.; Frearson, J.; Gray, D.; Hunter, W.N.  
Deposited on : 2011-10-26  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	<div> <div>2%</div> <div>85%</div> <div>13%</div> <div>..</div> </div>
1	B	286	<div> <div>3%</div> <div>87%</div> <div>12%</div> <div>..</div> </div>
1	C	286	<div> <div>13%</div> <div>84%</div> <div>14%</div> <div>..</div> </div>
1	D	286	<div> <div>12%</div> <div>75%</div> <div>19%</div> <div>• 5%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	1285	-	-	-	X
3	PEG	A	1286	-	-	X	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8632 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BIFUNCTIONAL PROTEIN FOLD.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	283	Total	C	N	O	S	0	4	0
			2139	1349	383	398	9			
1	B	284	Total	C	N	O	S	30	1	0
			2151	1350	389	402	10			
1	C	283	Total	C	N	O	S	115	0	0
			2137	1341	387	400	9			
1	D	273	Total	C	N	O	S	109	0	0
			2057	1296	369	383	9			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	EXPRESSION TAG	UNP Q9I2U6
A	0	HIS	-	EXPRESSION TAG	UNP Q9I2U6
B	-1	GLY	-	EXPRESSION TAG	UNP Q9I2U6
B	0	HIS	-	EXPRESSION TAG	UNP Q9I2U6
C	-1	GLY	-	EXPRESSION TAG	UNP Q9I2U6
C	0	HIS	-	EXPRESSION TAG	UNP Q9I2U6
D	-1	GLY	-	EXPRESSION TAG	UNP Q9I2U6
D	0	HIS	-	EXPRESSION TAG	UNP Q9I2U6

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula:  $C_4H_{10}O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	53	Total 53	O 53	0	0
4	B	54	Total 54	O 54	0	0
4	C	18	Total 18	O 18	0	0
4	D	10	Total 10	O 10	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: BIFUNCTIONAL PROTEIN FOLD




#### • Molecule 1: BIFUNCTIONAL PROTEIN FOLD




#### • Molecule 1: BIFUNCTIONAL PROTEIN FOLD



#### • Molecule 1: BIFUNCTIONAL PROTEIN FOLD







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	61.57Å 82.43Å 109.07Å 90.00° 94.68° 90.00°	Depositor
Resolution (Å)	39.13 – 2.20 39.13 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.9 (39.13-2.20) 99.9 (39.13-2.20)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.41 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.234 , 0.279 0.228 , 0.270	Depositor DCC
$R_{free}$ test set	2804 reflections (5.35%)	DCC
Wilson B-factor (Å <sup>2</sup> )	40.1	Xtriage
Anisotropy	0.336	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 40.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 55245 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8632	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 30.23 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.3545e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.39	0/2188	0.58	0/2982
1	B	0.99	4/2190 (0.2%)	0.81	6/2980 (0.2%)
1	C	0.85	6/2173 (0.3%)	0.74	9/2958 (0.3%)
1	D	0.90	10/2092 (0.5%)	0.76	11/2849 (0.4%)
All	All	0.81	20/8643 (0.2%)	0.73	26/11769 (0.2%)

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	194	ARG	CD-NE	-34.95	0.87	1.46
1	C	68	TYR	CB-CG	-21.90	1.18	1.51
1	D	282	LEU	CA-CB	-21.27	1.04	1.53
1	C	271	LEU	CB-CG	-20.47	0.93	1.52
1	B	56	LYS	CD-CE	-19.87	1.01	1.51
1	D	63	PHE	CB-CG	-18.54	1.19	1.51
1	D	22	ARG	CG-CD	-14.75	1.15	1.51
1	C	70	LEU	CA-CB	-11.60	1.27	1.53
1	D	61	VAL	CA-CB	-10.25	1.33	1.54
1	B	223	GLU	CB-CG	-9.29	1.34	1.52
1	B	27	ARG	CD-NE	-9.26	1.30	1.46
1	D	33	VAL	CA-CB	-8.11	1.37	1.54
1	D	29	GLN	CA-CB	-7.85	1.36	1.53
1	C	14	ASN	CB-CG	-7.42	1.33	1.51
1	C	78	ASP	CA-CB	-7.21	1.38	1.53
1	D	275	THR	CA-CB	-6.64	1.36	1.53
1	D	118	LYS	CA-CB	-6.29	1.40	1.53
1	D	252	ARG	CG-CD	-6.11	1.36	1.51
1	D	194	ARG	CB-CG	-6.06	1.36	1.52
1	C	33	VAL	CA-CB	-5.48	1.43	1.54

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	194	ARG	CG-CD-NE	20.96	155.82	111.80
1	B	56	LYS	CG-CD-CE	13.55	152.57	111.90
1	C	271	LEU	CA-CB-CG	13.46	146.26	115.30
1	B	194	ARG	CD-NE-CZ	12.51	141.12	123.60
1	D	64	LEU	N-CA-CB	12.04	134.48	110.40
1	C	2	THR	N-CA-CB	11.68	132.49	110.30
1	D	282	LEU	N-CA-CB	9.15	128.70	110.40
1	D	63	PHE	CB-CG-CD1	-8.04	115.17	120.80
1	D	118	LYS	CB-CA-C	7.86	126.12	110.40
1	D	22	ARG	CB-CG-CD	7.74	131.73	111.60
1	D	64	LEU	CA-CB-CG	7.62	132.82	115.30
1	B	56	LYS	CD-CE-NZ	7.26	128.41	111.70
1	D	63	PHE	CB-CG-CD2	6.95	125.66	120.80
1	C	43	THR	CB-CA-C	-6.62	93.73	111.60
1	D	63	PHE	CA-CB-CG	6.62	129.79	113.90
1	D	61	VAL	N-CA-CB	6.46	125.71	111.50
1	C	70	LEU	CB-CA-C	6.24	122.06	110.20
1	C	68	TYR	CA-CB-CG	5.98	124.76	113.40
1	C	29	GLN	N-CA-CB	5.95	121.31	110.60
1	D	282	LEU	CA-CB-CG	-5.94	101.64	115.30
1	C	79	LEU	CA-CB-CG	-5.76	102.05	115.30
1	C	63	PHE	CB-CA-C	-5.61	99.18	110.40
1	B	18	GLN	CB-CG-CD	5.51	125.93	111.60
1	B	27	ARG	CG-CD-NE	5.37	123.08	111.80
1	C	9	LYS	CB-CA-C	5.30	121.00	110.40
1	D	61	VAL	CA-CB-CG1	5.12	118.58	110.90

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2139	0	2174	34	0
1	B	2151	0	2194	18	0
1	C	2137	0	2174	20	0
1	D	2057	0	2097	31	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	6	0	8	0	0
3	A	7	0	10	5	0
4	A	53	0	0	0	0
4	B	54	0	0	0	0
4	C	18	0	0	0	0
4	D	10	0	0	0	0
All	All	8632	0	8657	99	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (99) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:167:ALA:HB2	1:C:188:VAL:HG12	1.66	0.76
1:A:157:LEU:HB2	3:A:1286:PEG:H22	1.73	0.70
1:B:167:ALA:HB3	1:B:190:HIS:HB3	1.73	0.70
1:A:120:VAL:HG23	1:A:271:LEU:HD21	1.72	0.69
1:C:150:LEU:HD21	1:C:205:LEU:HD11	1.75	0.69
1:A:151:ALA:HA	3:A:1286:PEG:H42	1.77	0.67
1:B:173:ARG:HB2	1:B:174:PRO:HD3	1.77	0.67
1:D:96:LEU:HD22	1:D:271:LEU:HD11	1.77	0.66
1:D:167:ALA:HB3	1:D:190:HIS:HB3	1.78	0.64
1:C:167:ALA:HB2	1:C:188:VAL:CG1	2.29	0.63
1:D:39:ILE:HG21	1:D:79:LEU:HD11	1.81	0.63
1:B:2:THR:HG22	1:B:2:THR:O	2.00	0.62
1:D:19:ILE:HD11	1:D:273:GLU:HG3	1.83	0.60
1:A:120:VAL:HG23	1:A:271:LEU:CD2	2.31	0.60
1:A:167:ALA:HB3	1:A:190:HIS:HB3	1.84	0.59
1:D:117:ASP:HA	1:D:137:LEU:HD11	1.84	0.59
1:A:283:HIS:O	1:A:284:ASP:HB2	2.03	0.58
1:D:173:ARG:HB2	1:D:174:PRO:HD3	1.86	0.58
1:A:271:LEU:C	1:A:271:LEU:HD13	2.25	0.57
1:A:157:LEU:HD22	3:A:1286:PEG:H22	1.86	0.57
1:A:106:ASP:O	1:A:110:LEU:HD23	2.04	0.57
1:A:240:LEU:HD12	1:A:240:LEU:N	2.19	0.57
1:C:150:LEU:HD21	1:C:205:LEU:CD1	2.35	0.55
1:D:200:VAL:HG11	1:D:220:TRP:CE3	2.42	0.53
1:A:3:ALA:HB2	1:A:253:ALA:O	2.08	0.53
1:D:57:ASP:HB3	1:D:272:LEU:HD11	1.91	0.53
1:A:104:HIS:HB3	1:D:72:ALA:HB3	1.91	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:80:LEU:CD1	1:B:110:LEU:HD23	2.39	0.52
1:B:167:ALA:HB1	1:B:173:ARG:NH1	2.25	0.52
1:D:7:ASP:OD2	1:D:10:ALA:HB2	2.10	0.52
1:A:51:VAL:HG23	1:A:98:GLN:OE1	2.09	0.51
1:C:76:GLN:HG3	1:C:110:LEU:HD11	1.91	0.51
1:D:93:ASP:OD1	1:D:283:HIS:ND1	2.43	0.51
1:C:170:ILE:H	1:C:170:ILE:HD12	1.76	0.51
1:A:157:LEU:HD22	3:A:1286:PEG:C3	2.41	0.50
1:C:3:ALA:HB2	1:C:253:ALA:O	2.11	0.50
1:D:42:GLY:HA2	1:D:72:ALA:HB2	1.93	0.49
1:C:102:PRO:HD2	1:C:105:LEU:HD12	1.94	0.49
1:A:98:GLN:HE21	1:A:99[B]:LEU:H	1.60	0.49
1:A:194:ARG:NH2	1:B:204:ASP:OD2	2.45	0.49
1:C:66:GLN:NE2	1:C:92:ILE:HD11	2.28	0.49
1:A:98:GLN:HE21	1:A:99[A]:LEU:H	1.60	0.48
1:C:114:ILE:HG22	1:C:115:HIS:O	2.13	0.48
1:A:283:HIS:O	1:A:284:ASP:CB	2.62	0.48
1:A:28:GLN:HE21	1:A:29:GLN:HG3	1.78	0.48
1:D:170:ILE:H	1:D:170:ILE:HD12	1.79	0.48
1:B:25:GLU:HA	1:B:28:GLN:HE21	1.79	0.47
1:C:167:ALA:HB3	1:C:190:HIS:HB3	1.97	0.47
1:D:6:ILE:HG22	1:D:149:LEU:HD22	1.97	0.47
1:D:96:LEU:HD22	1:D:271:LEU:CD1	2.45	0.46
1:C:206:VAL:HG12	1:C:208:VAL:HG23	1.97	0.46
1:B:2:THR:O	1:B:2:THR:CG2	2.63	0.45
1:D:70:LEU:HD23	1:D:74:THR:HG21	1.98	0.45
1:B:96:LEU:CD1	1:B:271:LEU:HD11	2.47	0.45
1:A:157:LEU:HB2	3:A:1286:PEG:C2	2.43	0.45
1:A:76:GLN:HG3	1:A:110:LEU:HD21	1.98	0.45
1:A:19:ILE:HG22	1:A:61:VAL:HG22	1.99	0.45
1:C:99:LEU:HD21	1:C:111:LEU:HD11	1.99	0.45
1:C:97:VAL:HB	1:C:111:LEU:HD21	1.98	0.45
1:D:34:PRO:HB3	1:D:279:ALA:HB2	1.98	0.45
1:D:132:ALA:HA	1:D:177:LEU:HD22	1.98	0.44
1:A:43:THR:HG22	1:A:43:THR:O	2.18	0.44
1:A:219:GLU:O	1:A:252:ARG:NH2	2.50	0.44
1:D:70:LEU:CD2	1:D:74:THR:HG21	2.48	0.44
1:A:27:ARG:CZ	1:A:33:VAL:HG21	2.47	0.44
1:A:169:ASN:HD22	1:A:232:ILE:HG22	1.83	0.43
1:B:70:LEU:HD23	1:B:74:THR:HG21	1.99	0.43
1:D:196:LEU:O	1:D:200:VAL:HG12	2.18	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:181:LEU:HD23	1:D:181:LEU:O	2.19	0.43
1:D:167:ALA:HB1	1:D:173:ARG:NH1	2.33	0.43
1:C:169:ASN:HD22	1:C:171:VAL:HG22	1.83	0.43
1:A:201:SER:HA	1:A:220:TRP:O	2.19	0.43
1:D:6:ILE:HG21	1:D:149:LEU:HB3	2.01	0.43
1:A:142:THR:HB	1:A:143:PRO:HD3	2.00	0.43
1:B:68:TYR:CE2	1:B:82:LEU:HD21	2.54	0.42
1:A:170:ILE:HD12	1:A:170:ILE:H	1.84	0.42
1:D:6:ILE:CG2	1:D:149:LEU:HD22	2.49	0.42
1:A:240:LEU:N	1:A:240:LEU:CD1	2.82	0.42
1:D:170:ILE:N	1:D:170:ILE:HD12	2.34	0.42
1:B:169:ASN:HD22	1:B:171:VAL:HB	1.85	0.42
1:D:19:ILE:HG23	1:D:276:LEU:HD22	2.02	0.42
1:B:80:LEU:HD12	1:B:110:LEU:HD23	2.01	0.41
1:B:211:GLY:N	1:B:232:ILE:HG23	2.36	0.41
1:C:125:PRO:O	1:D:129:GLY:HA3	2.20	0.41
1:D:25:GLU:HA	1:D:28:GLN:HE21	1.85	0.41
1:B:3:ALA:HB2	1:B:253:ALA:O	2.21	0.41
1:B:170:ILE:N	1:B:170:ILE:HD12	2.36	0.41
1:A:99[B]:LEU:HD23	1:A:99[B]:LEU:H	1.86	0.41
1:D:131:LEU:HD21	1:D:178:GLU:HG2	2.03	0.41
1:C:109:LEU:HD13	1:C:109:LEU:O	2.20	0.41
1:D:80:LEU:HD13	1:D:110:LEU:HD23	2.02	0.41
1:A:42:GLY:HA2	1:A:72:ALA:HB2	2.01	0.41
1:B:142:THR:HB	1:B:143:PRO:HD3	2.03	0.41
1:A:66:GLN:NE2	1:A:68:TYR:OH	2.54	0.41
1:C:134:ARG:O	1:C:135:MET:SD	2.79	0.41
1:B:19:ILE:O	1:B:23:VAL:HG23	2.20	0.40
1:C:188:VAL:HB	1:D:180:LEU:HD21	2.03	0.40
1:C:142:THR:HB	1:C:143:PRO:HD3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	285/286 (100%)	280 (98%)	5 (2%)	0	100	100
1	B	283/286 (99%)	276 (98%)	6 (2%)	1 (0%)	39	42
1	C	281/286 (98%)	268 (95%)	10 (4%)	3 (1%)	17	14
1	D	269/286 (94%)	258 (96%)	9 (3%)	2 (1%)	26	25
All	All	1118/1144 (98%)	1082 (97%)	30 (3%)	6 (0%)	34	35

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	48	GLN
1	B	166	GLY
1	C	245	GLU
1	D	168	SER
1	D	245	GLU
1	C	170	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	225/228 (99%)	220 (98%)	5 (2%)	60	72
1	B	228/228 (100%)	223 (98%)	5 (2%)	60	72
1	C	226/228 (99%)	224 (99%)	2 (1%)	84	92
1	D	218/228 (96%)	212 (97%)	6 (3%)	51	63
All	All	897/912 (98%)	879 (98%)	18 (2%)	63	76

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	GLN
1	A	61	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	64	LEU
1	A	79	LEU
1	A	157	LEU
1	B	21	GLN
1	B	64	LEU
1	B	65	SER
1	B	80	LEU
1	B	82	LEU
1	C	2	THR
1	C	271	LEU
1	D	22	ARG
1	D	79	LEU
1	D	111	LEU
1	D	171	VAL
1	D	198	ASP
1	D	275	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	GLN
1	A	28	GLN
1	A	48	GLN
1	A	66	GLN
1	B	28	GLN
1	B	48	GLN
1	B	66	GLN
1	B	169	ASN
1	B	281	HIS
1	C	17	GLN
1	C	66	GLN
1	C	169	ASN
1	D	14	ASN
1	D	17	GLN
1	D	28	GLN
1	D	133	GLN
1	D	281	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.



## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GOL	A	1285	-	5,5,5	0.35	0	5,5,5	0.34	0
3	PEG	A	1286	-	6,6,6	0.48	0	5,5,5	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	1285	-	-	0/4/4/4	0/0/0/0
3	PEG	A	1286	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1286	PEG	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	283/286 (98%)	0.18	6 (2%) 67 65	21, 39, 70, 87	0
1	B	284/286 (99%)	0.14	8 (2%) 56 55	23, 43, 65, 73	9 (3%)
1	C	283/286 (98%)	0.62	37 (13%) 5 4	29, 62, 125, 168	28 (9%)
1	D	273/286 (95%)	0.76	34 (12%) 5 5	26, 69, 106, 133	26 (9%)
All	All	1123/1144 (98%)	0.42	85 (7%) 17 16	21, 50, 110, 168	63 (5%)

All (85) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	262	GLY	6.3
1	C	81	ALA	5.5
1	C	41	VAL	5.1
1	D	30	GLY	5.0
1	C	23	VAL	5.0
1	D	71	PRO	5.0
1	C	42	GLY	4.8
1	C	101	LEU	4.7
1	B	71	PRO	4.6
1	D	41	VAL	4.6
1	C	13	ALA	4.6
1	D	31	LEU	4.5
1	D	261	GLY	4.2
1	D	168	SER	4.0
1	D	50	TYR	3.9
1	D	2	THR	3.8
1	D	167	ALA	3.7
1	A	72	ALA	3.6
1	D	45	PRO	3.6
1	D	67	ALA	3.6
1	C	284	ASP	3.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	83	ILE	3.4
1	B	168	SER	3.4
1	D	35	GLY	3.3
1	A	41	VAL	3.3
1	C	104	HIS	3.2
1	B	167	ALA	3.2
1	C	46	ALA	3.1
1	D	43	THR	3.1
1	B	104	HIS	3.1
1	D	192	PHE	3.0
1	A	45	PRO	3.0
1	D	103	ALA	2.9
1	C	52	ALA	2.9
1	D	94	GLY	2.9
1	D	83	ILE	2.9
1	C	19	ILE	2.8
1	C	113	ARG	2.8
1	C	40	LEU	2.8
1	C	2	THR	2.8
1	D	23	VAL	2.7
1	D	72	ALA	2.7
1	C	37	ALA	2.7
1	D	92	ILE	2.7
1	C	44	ASP	2.7
1	C	24	THR	2.7
1	A	103	ALA	2.6
1	D	70	LEU	2.6
1	C	250	ALA	2.6
1	C	54	LYS	2.6
1	D	25	GLU	2.5
1	D	28	GLN	2.5
1	C	68	TYR	2.5
1	C	12	ALA	2.4
1	C	168	SER	2.4
1	B	40	LEU	2.4
1	C	70	LEU	2.4
1	D	32	ARG	2.4
1	D	255	TRP	2.4
1	C	77	ASP	2.3
1	C	247	GLU	2.3
1	C	111	LEU	2.3
1	C	49	VAL	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	3	ALA	2.3
1	B	247	GLU	2.3
1	C	47	SER	2.3
1	A	79	LEU	2.3
1	D	102	PRO	2.2
1	C	20	ALA	2.2
1	D	59	GLU	2.2
1	C	35	GLY	2.2
1	C	62	GLY	2.2
1	B	1	MET	2.2
1	C	82	LEU	2.1
1	B	43	THR	2.1
1	D	79	LEU	2.1
1	C	71	PRO	2.1
1	C	16	ARG	2.1
1	C	53	HIS	2.1
1	C	283	HIS	2.1
1	D	263	VAL	2.1
1	A	168	SER	2.0
1	D	38	VAL	2.0
1	D	4	GLN	2.0
1	D	68	TYR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PEG	A	1286	7/7	0.82	0.25	5.04	38,40,41,41	0
2	GOL	A	1285	6/6	0.88	0.26	4.95	61,62,63,63	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.