



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:18 AM GMT

PDB ID : 3A8T
Title : Plant adenylate isopentenyltransferase in complex with ATP
Authors : Chu, H.-M.; Ko, T.-P.; Wang, A.H.-J.
Deposited on : 2009-10-09
Resolution : 2.37 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4019 (2.40-2.36)
Clashscore	102246	4595 (2.40-2.36)
Ramachandran outliers	100387	4520 (2.40-2.36)
Sidechain outliers	100360	4522 (2.40-2.36)
RSRZ outliers	91569	4034 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	339	<div> <div>5%</div> <div>60%</div> <div>22%</div> <div>• •</div> <div>15%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2511 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenylate isopentenyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	289	Total	C	N	O	S	0	0	0
			2277	1428	405	435	9			

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	MET	-	INITIATING METHIONINE	UNP Q5GHF7
A	-8	ARG	-	EXPRESSION TAG	UNP Q5GHF7
A	-7	GLY	-	EXPRESSION TAG	UNP Q5GHF7
A	-6	SER	-	EXPRESSION TAG	UNP Q5GHF7
A	-5	HIS	-	EXPRESSION TAG	UNP Q5GHF7
A	-4	HIS	-	EXPRESSION TAG	UNP Q5GHF7
A	-3	HIS	-	EXPRESSION TAG	UNP Q5GHF7
A	-2	HIS	-	EXPRESSION TAG	UNP Q5GHF7
A	-1	HIS	-	EXPRESSION TAG	UNP Q5GHF7
A	0	HIS	-	EXPRESSION TAG	UNP Q5GHF7

- Molecule 2 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	198	Total 198	O 198	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	45.42Å 75.89Å 81.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.47 – 2.37 25.47 – 2.34	Depositor EDS
% Data completeness (in resolution range)	93.6 (25.47-2.37) 92.2 (25.47-2.34)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.30 (at 2.33Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.181 , 0.235 0.180 , 0.232	Depositor DCC
R_{free} test set	581 reflections (5.22%)	DCC
Wilson B-factor (Å ²)	28.2	Xtriage
Anisotropy	0.014	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 48.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 11382 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2511	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	1/2317 (0.0%)	0.96	6/3118 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	317	GLU	CG-CD	6.09	1.61	1.51

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	215	ARG	NE-CZ-NH2	-6.62	116.99	120.30
1	A	215	ARG	NE-CZ-NH1	6.53	123.57	120.30
1	A	82	ARG	NE-CZ-NH2	-6.18	117.21	120.30
1	A	189	LEU	CA-CB-CG	5.66	128.32	115.30
1	A	281	GLY	N-CA-C	-5.58	99.15	113.10
1	A	191	LEU	CA-CB-CG	5.25	127.39	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2277	0	2268	80	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	31	0	12	1	0
3	A	5	0	0	0	0
4	A	198	0	0	18	0
All	All	2511	0	2280	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:266:CYS:HB3	4:A:687:HOH:O	1.71	0.91
1:A:240:GLU:HB3	1:A:244:PRO:HG3	1.54	0.89
1:A:246:ARG:O	1:A:249:VAL:HG12	1.76	0.85
1:A:195:ASP:OD2	4:A:671:HOH:O	1.93	0.85
1:A:328:ASP:HB3	1:A:329:GLU:OE1	1.78	0.83
1:A:276:ILE:HA	1:A:279:LEU:CD1	2.10	0.82
1:A:68:LYS:HE2	1:A:94:ASP:HA	1.65	0.79
1:A:225:PRO:HD2	1:A:226:GLU:OE1	1.83	0.78
1:A:97:ARG:HH11	1:A:97:ARG:HG2	1.47	0.78
1:A:238:ASP:OD1	4:A:683:HOH:O	2.06	0.74
1:A:36:MET:HE1	1:A:286:LEU:HD22	1.70	0.74
1:A:92:GLU:HB2	4:A:620:HOH:O	1.89	0.73
1:A:89:LEU:HD12	1:A:105:PHE:CE1	2.24	0.71
1:A:169:TRP:CZ3	1:A:288:ARG:HD3	2.27	0.70
1:A:300:THR:C	1:A:302:ASP:H	1.94	0.69
1:A:36:MET:HE1	1:A:286:LEU:CD2	2.23	0.68
1:A:317:GLU:HG2	1:A:318:PRO:HD3	1.74	0.68
1:A:97:ARG:CG	1:A:97:ARG:HH11	2.07	0.67
1:A:232:GLU:OE2	4:A:682:HOH:O	2.13	0.67
1:A:115:GLU:OE2	4:A:698:HOH:O	2.11	0.67
1:A:231:PHE:O	1:A:235:ARG:HD3	1.95	0.66
1:A:92:GLU:OE2	4:A:574:HOH:O	2.13	0.66
1:A:246:ARG:NH1	4:A:683:HOH:O	2.25	0.65
1:A:36:MET:CE	1:A:286:LEU:HD22	2.26	0.65
1:A:210:GLU:O	1:A:215:ARG:HD2	1.98	0.63
1:A:234:PHE:HZ	1:A:249:VAL:HG13	1.63	0.63
1:A:279:LEU:O	1:A:284:TRP:HB2	2.02	0.60
1:A:271:ARG:HG3	4:A:686:HOH:O	1.99	0.60
1:A:281:GLY:C	1:A:283:GLY:H	2.07	0.58
1:A:291:ALA:O	1:A:295:PHE:HD2	1.87	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:89:LEU:HD12	1:A:105:PHE:HE1	1.68	0.57
1:A:317:GLU:HG2	1:A:318:PRO:CD	2.35	0.57
1:A:60:ASN:HD21	1:A:62:ASP:HB3	1.69	0.57
1:A:234:PHE:CE2	1:A:250:ARG:HA	2.40	0.56
1:A:244:PRO:C	1:A:246:ARG:N	2.59	0.56
1:A:97:ARG:CG	1:A:97:ARG:NH1	2.64	0.55
1:A:244:PRO:C	1:A:246:ARG:H	2.08	0.55
1:A:240:GLU:CB	1:A:244:PRO:HG3	2.33	0.54
1:A:305:GLU:O	1:A:309:GLU:HG3	2.07	0.54
1:A:28:ARG:HG2	1:A:28:ARG:HH11	1.73	0.53
1:A:275:LYS:O	1:A:279:LEU:HD12	2.10	0.52
1:A:204:GLU:CG	1:A:204:GLU:O	2.57	0.52
1:A:300:THR:C	1:A:302:ASP:N	2.63	0.51
1:A:70:LEU:HB3	1:A:73:THR:HB	1.93	0.50
1:A:240:GLU:OE1	1:A:244:PRO:HG3	2.12	0.50
1:A:320:VAL:O	1:A:324:SER:HB2	2.12	0.50
1:A:276:ILE:HA	1:A:279:LEU:HD12	1.88	0.49
1:A:177:LEU:HG	1:A:181:LEU:HD22	1.94	0.49
1:A:291:ALA:HB1	1:A:295:PHE:CE2	2.47	0.49
1:A:68:LYS:HG2	1:A:91:GLY:O	2.13	0.48
1:A:279:LEU:H	1:A:279:LEU:HD12	1.77	0.48
1:A:75:ASN:ND2	1:A:188:MET:SD	2.86	0.48
1:A:77:ILE:HG23	1:A:82:ARG:HE	1.79	0.48
1:A:244:PRO:HB2	1:A:246:ARG:HB2	1.96	0.47
1:A:87:HIS:HE1	4:A:514:HOH:O	1.96	0.47
1:A:60:ASN:ND2	4:A:565:HOH:O	2.40	0.47
1:A:216:THR:HG22	4:A:583:HOH:O	2.13	0.47
1:A:92:GLU:N	4:A:620:HOH:O	2.44	0.47
1:A:275:LYS:O	1:A:279:LEU:CD1	2.63	0.46
1:A:215:ARG:NH2	1:A:224:VAL:HG12	2.29	0.46
1:A:291:ALA:HB1	1:A:295:PHE:HE2	1.80	0.46
1:A:259:ARG:HD2	4:A:606:HOH:O	2.15	0.46
1:A:224:VAL:N	1:A:225:PRO:CD	2.80	0.45
1:A:28:ARG:HG2	1:A:28:ARG:NH1	2.32	0.44
1:A:281:GLY:C	1:A:283:GLY:N	2.71	0.43
1:A:281:GLY:O	1:A:283:GLY:N	2.51	0.43
1:A:55:PRO:HA	4:A:548:HOH:O	2.17	0.43
1:A:211:ASP:OD1	4:A:603:HOH:O	2.21	0.43
1:A:287:ARG:NE	4:A:667:HOH:O	2.46	0.42
1:A:36:MET:HE2	1:A:36:MET:HB2	1.74	0.42
1:A:68:LYS:HE3	1:A:68:LYS:HB2	1.66	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:116:ILE:HG22	1:A:121:LYS:O	2.19	0.42
1:A:276:ILE:CA	1:A:279:LEU:HD12	2.48	0.42
1:A:259:ARG:CD	4:A:606:HOH:O	2.68	0.41
1:A:181:LEU:HG	1:A:265:THR:CG2	2.50	0.41
1:A:43:LYS:HE2	1:A:43:LYS:HB2	1.75	0.41
1:A:276:ILE:O	1:A:279:LEU:HD12	2.21	0.41
1:A:160:GLU:HG2	1:A:161:LEU:N	2.36	0.41
1:A:181:LEU:HG	1:A:265:THR:HG22	2.03	0.41
1:A:144:SER:O	1:A:145:GLY:C	2.58	0.41
2:A:800:ATP:O1G	2:A:800:ATP:O2B	2.39	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	285/339 (84%)	268 (94%)	12 (4%)	5 (2%)	11 12

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	284	TRP
1	A	282	ALA
1	A	301	SER
1	A	281	GLY
1	A	244	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	245/287 (85%)	230 (94%)	15 (6%)	23	34

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	43	LYS
1	A	45	ARG
1	A	89	LEU
1	A	139	ASP
1	A	143	SER
1	A	162	ARG
1	A	181	LEU
1	A	189	LEU
1	A	191	LEU
1	A	243	ASP
1	A	256	GLU
1	A	278	ARG
1	A	296	ARG
1	A	317	GLU
1	A	328	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	60	ASN
1	A	87	HIS
1	A	130	ASN
1	A	314	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ATP	A	800	-	24,33,33	1.74	4 (16%)	31,52,52	2.24	6 (19%)
3	PO4	A	900	-	4,4,4	1.06	0	6,6,6	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ATP	A	800	-	-	0/18/38/38	0/3/3/3
3	PO4	A	900	-	-	0/0/0/0	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	800	ATP	PB-O1B	-2.08	1.43	1.51
2	A	800	ATP	C5-C4	2.20	1.45	1.40
2	A	800	ATP	PG-O2G	2.44	1.63	1.54
2	A	800	ATP	O4'-C1'	5.89	1.48	1.41

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	800	ATP	N3-C2-N1	-7.98	122.78	128.89
2	A	800	ATP	PB-O3B-PG	-5.23	115.13	132.67
2	A	800	ATP	PA-O3A-PB	-4.94	118.86	132.73

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	800	ATP	C4-C5-N7	-3.03	106.69	109.48
2	A	800	ATP	C2'-C1'-N9	-2.39	110.64	114.29
2	A	800	ATP	O3A-PA-O5'	3.15	111.28	102.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	800	ATP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	289/339 (85%)	-0.18	16 (5%)	29 33	16, 28, 67, 88	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	244	PRO	5.6
1	A	303	SER	4.6
1	A	146	PRO	4.1
1	A	302	ASP	3.6
1	A	243	ASP	3.2
1	A	145	GLY	3.2
1	A	28	ARG	3.2
1	A	241	GLY	2.9
1	A	301	SER	2.4
1	A	242	GLU	2.3
1	A	305	GLU	2.3
1	A	245	GLY	2.3
1	A	329	GLU	2.3
1	A	300	THR	2.2
1	A	283	GLY	2.0
1	A	282	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	PO4	A	900	5/5	0.99	0.14	0.30	28,29,33,33	0
2	ATP	A	800	31/31	0.98	0.10	-0.98	23,26,48,48	0

6.5 Other polymers [i](#)

There are no such residues in this entry.