



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 09:26 PM GMT

PDB ID : 5ABE
Title : Structure of GH84 with ligand
Authors : Bergeron-Brlek, M.; Goodwin-Tindall, J.; Cekic, N.; Varghese, V.; Zandberg, W.F.; Shan, X.; Roth, C.; Chan, S.; Davies, G.J.; Vocadlo, D.J.; Britton, R.
Deposited on : 2015-08-05
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	726	
1	B	726	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	XQO	A	1716	-	-	-	X
2	XQO	B	1717	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11263 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called O-GLCNACASE BT_4395.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	681	Total	C	N	O	S	0	1	0
			5527	3541	934	1033	19			
1	B	670	Total	C	N	O	S	0	1	0
			5456	3497	924	1017	18			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	MET	-	EXPRESSION TAG	UNP Q89ZI2
A	-8	GLY	-	EXPRESSION TAG	UNP Q89ZI2
A	-7	SER	-	EXPRESSION TAG	UNP Q89ZI2
A	-6	SER	-	EXPRESSION TAG	UNP Q89ZI2
A	-5	HIS	-	EXPRESSION TAG	UNP Q89ZI2
A	-4	HIS	-	EXPRESSION TAG	UNP Q89ZI2
A	-3	HIS	-	EXPRESSION TAG	UNP Q89ZI2
A	-2	HIS	-	EXPRESSION TAG	UNP Q89ZI2
A	-1	HIS	-	EXPRESSION TAG	UNP Q89ZI2
A	0	HIS	-	EXPRESSION TAG	UNP Q89ZI2
B	-9	MET	-	EXPRESSION TAG	UNP Q89ZI2
B	-8	GLY	-	EXPRESSION TAG	UNP Q89ZI2
B	-7	SER	-	EXPRESSION TAG	UNP Q89ZI2
B	-6	SER	-	EXPRESSION TAG	UNP Q89ZI2
B	-5	HIS	-	EXPRESSION TAG	UNP Q89ZI2
B	-4	HIS	-	EXPRESSION TAG	UNP Q89ZI2
B	-3	HIS	-	EXPRESSION TAG	UNP Q89ZI2
B	-2	HIS	-	EXPRESSION TAG	UNP Q89ZI2
B	-1	HIS	-	EXPRESSION TAG	UNP Q89ZI2
B	0	HIS	-	EXPRESSION TAG	UNP Q89ZI2

- Molecule 2 is 2-[(2S,3S,4R,5R)-5-(HYDROXYMETHYL)-3,4-BIS(OXIDANYL)-1-PENTYL-PYRROLIDIN-2-YL]-N-METHYL-ETHANAMIDE (three-letter code: XQO) (formula: C₁₃H₂₆N₂O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			19	13	2	4		
2	B	1	Total	C	N	O	0	0
			19	13	2	4		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Ca	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	125	Total	O	0	0
			125	125		
4	B	116	Total	O	0	0
			116	116		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: O-GLCNACASE BT_4395



• Molecule 1: O-GLCNACASE BT_4395



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2 ₁	Depositor
Cell constants a, b, c, α , β , γ	51.57Å 159.60Å 223.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	129.81 – 2.00 129.81 – 2.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (129.81-2.00) 89.1 (129.81-2.00)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.75 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0124	Depositor
R, R_{free}	0.207 , 0.244 0.210 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	35.7	Xtriage
Anisotropy	0.648	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 37.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	4 of 125589 reflections (0.003%)	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11263	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.42 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 8.7184e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, XQO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.76	1/5662 (0.0%)	0.80	7/7669 (0.1%)
1	B	0.80	0/5589	0.83	6/7566 (0.1%)
All	All	0.78	1/11251 (0.0%)	0.81	13/15235 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	438	GLU	CD-OE1	5.14	1.31	1.25

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	235	ARG	NE-CZ-NH1	6.80	123.70	120.30
1	B	235	ARG	NE-CZ-NH1	6.49	123.54	120.30
1	B	219	ARG	NE-CZ-NH1	6.11	123.36	120.30
1	A	72	ARG	NE-CZ-NH1	5.90	123.25	120.30
1	B	168	ASP	CB-CG-OD1	5.84	123.56	118.30
1	B	348	ASP	CB-CG-OD1	5.48	123.23	118.30
1	A	567	ARG	NE-CZ-NH1	5.48	123.04	120.30
1	B	408	ARG	CG-CD-NE	5.46	123.26	111.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	398	ASP	CB-CG-OD1	5.35	123.11	118.30
1	A	567	ARG	NE-CZ-NH2	-5.17	117.71	120.30
1	A	474	ARG	NE-CZ-NH1	5.13	122.87	120.30
1	B	427	ASP	CB-CG-OD1	5.05	122.84	118.30
1	A	235	ARG	NE-CZ-NH2	-5.03	117.78	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	628	GLY	Peptide
1	B	457	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5527	0	5446	24	1
1	B	5456	0	5389	21	0
2	A	19	0	26	2	0
2	B	19	0	26	0	0
3	B	1	0	0	0	0
4	A	125	0	0	0	0
4	B	116	0	0	0	0
All	All	11263	0	10887	45	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:355:VAL:O	1:B:399:THR:HG23	1.90	0.71
1:A:355:VAL:O	1:A:399:THR:HG23	1.94	0.67
1:A:438:GLU:OE1	1:A:441:MET:HG3	1.96	0.65

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:452:LYS:N	1:A:452:LYS:HD3	2.18	0.58
1:A:673:LEU:HD22	1:A:680:LEU:HB3	1.88	0.56
1:B:263:GLU:HA	1:B:267:GLN:HG2	1.90	0.54
1:A:557:ALA:HB1	1:A:561:ILE:HB	1.89	0.54
1:B:70:TYR:OH	1:B:89:GLU:OE2	2.26	0.50
1:A:661:ILE:HD13	1:A:671:VAL:HG21	1.94	0.49
1:A:644:ILE:HD11	1:A:661:ILE:HD11	1.95	0.48
1:A:591:MET:HG3	1:A:593:HIS:O	2.13	0.48
1:B:354:PRO:HB2	1:B:399:THR:HG22	1.95	0.48
1:B:62[A]:LYS:NZ	1:B:71:SER:O	2.45	0.48
1:A:137:TYR:CE2	2:A:1716:XQO:H4	2.49	0.48
1:A:137:TYR:HE2	2:A:1716:XQO:H4	1.78	0.48
1:A:354:PRO:HB2	1:A:399:THR:HG22	1.96	0.47
1:A:252:GLN:HE22	1:A:300:LYS:HE2	1.79	0.47
1:B:310:THR:HA	1:B:337:TRP:O	2.14	0.47
1:B:536:GLN:HG2	1:B:590:TYR:CD1	2.50	0.47
1:B:616:ILE:HD12	1:B:709:PHE:CG	2.50	0.47
1:A:125:PRO:HB3	1:A:392:TRP:CE3	2.50	0.46
1:A:310:THR:HA	1:A:337:TRP:O	2.16	0.46
1:A:308:MET:HA	1:A:335:TYR:O	2.17	0.45
1:B:562:LYS:HB3	1:B:563:PRO:HD3	1.99	0.45
1:A:131:GLY:O	1:A:370:VAL:HA	2.17	0.45
1:B:356:TYR:HB3	1:B:399:THR:HG21	1.98	0.45
1:B:708:GLN:HG2	1:B:710:VAL:HG23	1.99	0.45
1:B:86:ASN:HA	1:B:118:GLU:HG3	1.98	0.45
1:B:557:ALA:HB1	1:B:561:ILE:HB	1.99	0.44
1:A:562:LYS:HB3	1:A:563:PRO:HD3	2.00	0.43
1:A:625:TRP:CZ2	1:A:706:LEU:N	2.87	0.43
1:B:468:LEU:HD23	1:B:468:LEU:HA	1.88	0.43
1:A:238:ALA:HA	1:A:276:VAL:O	2.19	0.42
1:B:676:LYS:C	1:B:676:LYS:HE3	2.39	0.42
1:B:354:PRO:HB2	1:B:399:THR:CG2	2.50	0.41
1:A:423[A]:MET:HB2	1:A:423[A]:MET:HE3	1.88	0.41
1:B:33:ALA:HB2	1:B:60:GLY:HA2	2.02	0.41
1:B:131:GLY:HA3	1:B:160:THR:O	2.21	0.41
1:A:644:ILE:CD1	1:A:661:ILE:HD11	2.51	0.41
1:A:283:ASN:HB3	1:A:310:THR:OG1	2.20	0.41
1:A:170:TYR:HB2	1:A:180:TYR:CE2	2.55	0.41
1:B:308:MET:HA	1:B:335:TYR:O	2.21	0.40
1:A:72:ARG:HG2	1:A:72:ARG:HH11	1.87	0.40
1:B:483:LEU:HG	1:B:499:VAL:HG11	2.04	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:81:TYR:CZ	1:B:123:ASP:HB3	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:170:TYR:OH	1:A:442:ASP:OD1[1_655]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	670/726 (92%)	644 (96%)	25 (4%)	1 (0%)	56	53
1	B	659/726 (91%)	635 (96%)	23 (4%)	1 (0%)	52	48
All	All	1329/1452 (92%)	1279 (96%)	48 (4%)	2 (0%)	52	48

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	629	ASN
1	B	291	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	599/639 (94%)	578 (96%)	21 (4%)	43	40
1	B	592/639 (93%)	575 (97%)	17 (3%)	50	49
All	All	1191/1278 (93%)	1153 (97%)	38 (3%)	47	44

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	52	LYS
1	A	53	LYS
1	A	65	LYS
1	A	68	ARG
1	A	72	ARG
1	A	183	LYS
1	A	371	THR
1	A	423[A]	MET
1	A	423[B]	MET
1	A	452	LYS
1	A	519	ARG
1	A	594	LYS
1	A	615	LEU
1	A	660	GLU
1	A	666	LYS
1	A	671	VAL
1	A	689	VAL
1	A	695	THR
1	A	706	LEU
1	A	707	ARG
1	A	715	LYS
1	B	15	GLN
1	B	48	LYS
1	B	112	LYS
1	B	113	ASP
1	B	258	LEU
1	B	299	ASP
1	B	371	THR
1	B	423	MET
1	B	574	LYS
1	B	594	LYS
1	B	644	ILE
1	B	661	ILE
1	B	666	LYS
1	B	672	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	676	LYS
1	B	693	ARG
1	B	694	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	16	ASN
1	A	252	GLN
1	A	444	GLN
1	A	578	GLN
1	A	608	GLN
1	A	696	ASN
1	B	2	ASN
1	B	252	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	XQO	A	1716	-	19,19,19	1.06	2 (10%)	16,25,25	2.72	4 (25%)
2	XQO	B	1717	-	19,19,19	1.10	2 (10%)	16,25,25	2.27	3 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XQO	A	1716	-	-	0/13/33/33	0/1/1/1
2	XQO	B	1717	-	-	0/13/33/33	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1716	XQO	C4-C3	-2.79	1.49	1.53
2	B	1717	XQO	C4-C3	-2.16	1.50	1.53
2	A	1716	XQO	C1-N2	2.79	1.50	1.46
2	B	1717	XQO	C1-N2	3.41	1.51	1.46

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1716	XQO	C3-C11-C12	-8.73	100.04	112.58
2	B	1717	XQO	C3-C11-C12	-7.56	101.73	112.58
2	A	1716	XQO	O19-C18-C6	-3.69	103.53	111.45
2	B	1717	XQO	O13-C12-C11	-2.25	117.86	121.36
2	A	1716	XQO	O13-C12-C11	-2.08	118.12	121.36
2	B	1717	XQO	C4-C5-C6	3.23	108.21	103.00
2	A	1716	XQO	C4-C5-C6	3.59	108.79	103.00

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1716	XQO	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	681/726 (93%)	0.03	9 (1%) 79 80	32, 48, 92, 116	0
1	B	670/726 (92%)	0.15	24 (3%) 46 48	31, 50, 95, 122	0
All	All	1351/1452 (93%)	0.09	33 (2%) 62 63	31, 49, 95, 122	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	706	LEU	4.7
1	B	673	LEU	4.5
1	B	685	GLN	4.3
1	A	680	LEU	3.8
1	B	605	LEU	3.6
1	B	709	PHE	3.1
1	B	659	LEU	3.1
1	B	51	SER	3.0
1	B	658	ARG	3.0
1	B	46	SER	2.9
1	A	618	PRO	2.9
1	B	715	LYS	2.9
1	B	52	LYS	2.8
1	A	709	PHE	2.7
1	B	48	LYS	2.7
1	B	689	VAL	2.7
1	B	290	ASN	2.6
1	B	667	GLU	2.6
1	B	692	VAL	2.5
1	A	707	ARG	2.5
1	A	49	GLN	2.5
1	B	286	TRP	2.4
1	B	519	ARG	2.3
1	B	638	ILE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	661	ILE	2.3
1	B	454	PHE	2.2
1	A	674	LYS	2.2
1	B	216	LYS	2.2
1	B	670	THR	2.1
1	A	682	ALA	2.1
1	A	648	PHE	2.1
1	B	214	TRP	2.0
1	B	680	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	XQO	A	1716	19/19	0.88	0.26	10.07	42,59,67,69	0
2	XQO	B	1717	19/19	0.79	0.30	4.28	46,56,66,66	19
3	CA	B	1716	1/1	0.97	0.10	-	42,42,42,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.