



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 09:33 PM GMT

PDB ID : 5ADK
Title : Structure of bovine endothelial nitric oxide synthase heme domain in complex with 7-((3-(Dimethylamino)methyl)phenoxy)methyl)quinolin-2- amine
Authors : Li, H.; Poulos, T.L.
Deposited on : 2015-08-20
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	443	<div> <div>9%</div> <div>85%</div> <div>7% • 6%</div> </div>
1	B	443	<div> <div>5%</div> <div>82%</div> <div>9% 9%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	7M3	B	800	-	-	-	X
5	ACT	A	860	-	-	-	X
5	ACT	A	861	-	-	-	X
5	ACT	B	860	-	-	-	X
6	GOL	A	880	-	-	-	X

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NITRIC OXIDE SYNTHASE, ENDOTHELIAL.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	416	Total 3305	As 1	C 2101	N 584	O 603	S 16	0	0	0
1	B	404	Total 3221	As 1	C 2048	N 568	O 588	S 16	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	100	ARG	CYS	CONFLICT	UNP P29473
B	100	ARG	CYS	CONFLICT	UNP P29473

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	Fe	N	O	
2	B	1	43	34	1	4	4	0

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	17	9	5	3	0	0
3	B	1	17	9	5	3	0	0

- Molecule 4 is 7-[[3-[(DIMETHYLAMINO)METHYL]PHENOXY]METHYL]QUINOLIN-2-AMINE (three-letter code: 7M3) (formula: $C_{19}H_{21}N_3O$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			23	19	3	1		
4	B	1	Total	C	N	O	0	0
			23	19	3	1		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Zn	0	0
			1	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	291	Total	O	0	0
			291	291		
8	B	211	Total	O	0	0
			211	211		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NITRIC OXIDE SYNTHASE, ENDOTHELIAL



• Molecule 1: NITRIC OXIDE SYNTHASE, ENDOTHELIAL



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.96Å 106.38Å 157.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.19 – 1.80 39.19 – 1.81	Depositor EDS
% Data completeness (in resolution range)	97.9 (39.19-1.80) 97.9 (39.19-1.81)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.29 (at 1.81Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.164 , 0.193 0.171 , 0.198	Depositor DCC
R_{free} test set	4369 reflections (5.22%)	DCC
Wilson B-factor (Å ²)	33.7	Xtriage
Anisotropy	0.131	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 46.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 88064 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7223	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, H4B, CAS, 7M3, ACT, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/3390	0.53	0/4620
1	B	0.38	0/3300	0.52	0/4491
All	All	0.37	0/6690	0.52	0/9111

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3305	0	3215	24	1
1	B	3221	0	3130	21	0
2	A	43	0	30	1	0
2	B	43	0	30	1	0
3	A	17	0	15	1	0
3	B	17	0	15	0	0
4	A	23	0	21	3	0
4	B	23	0	21	3	0
5	A	8	0	6	0	0
5	B	8	0	6	1	0
6	A	6	0	8	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	6	0	8	0	0
7	A	1	0	0	0	0
8	A	291	0	0	2	1
8	B	211	0	0	1	0
All	All	7223	0	6505	42	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:72:ARG:NH1	1:A:83:TYR:OH	2.23	0.72
1:A:99:ARG:NH1	1:B:93:ASP:OD1	2.26	0.68
1:A:240:ARG:NH2	8:A:2157:HOH:O	2.19	0.65
1:A:108:PRO:HG2	1:A:111:LEU:HG	1.85	0.57
2:B:500:HEM:HHC	2:B:500:HEM:HBB2	1.87	0.56
1:A:240:ARG:HD3	1:A:298:PRO:HB3	1.88	0.56
1:A:249:GLN:OE1	4:A:800:7M3:H24	2.06	0.55
1:B:71:PRO:HG2	1:B:84:ASP:HB3	1.88	0.55
1:A:342:LEU:HD11	1:A:349:GLU:HB3	1.88	0.55
1:B:355:PHE:HE1	4:B:800:7M3:H06	1.73	0.53
1:A:77:GLU:HG3	1:B:372:PRO:HG2	1.91	0.52
1:A:105:LEU:HD22	1:B:465:GLU:HB3	1.91	0.51
1:B:281:TRP:HB2	1:B:304:LEU:HD21	1.91	0.51
1:A:455:SER:OG	1:B:398:ASP:OD2	2.19	0.50
1:B:252:ARG:NH2	5:B:861:ACT:O	2.44	0.50
1:A:458:LEU:HD22	1:B:378:LEU:HD21	1.94	0.49
1:B:338:VAL:HG21	4:B:800:7M3:C07	2.42	0.49
1:B:378:LEU:HB2	8:B:2170:HOH:O	2.12	0.49
1:A:355:PHE:HE1	4:A:800:7M3:H06	1.78	0.49
1:A:338:VAL:HG21	4:A:800:7M3:C07	2.43	0.48
2:A:500:HEM:HHC	2:A:500:HEM:HBB2	1.96	0.47
1:B:249:GLN:OE1	4:B:800:7M3:H24	2.15	0.46
1:B:366:THR:O	1:B:370:CYS:HB2	2.15	0.46
1:A:465:GLU:HB3	1:B:105:LEU:HD22	1.98	0.46
1:A:72:ARG:HD2	8:A:2020:HOH:O	2.15	0.46
3:A:600:H4B:O4	6:A:880:GOL:O3	2.33	0.45
1:A:259:GLN:HG2	1:A:259:GLN:H	1.47	0.45
1:A:370:CYS:SG	1:A:378:LEU:HD13	2.57	0.45
1:A:126:LEU:HA	1:A:126:LEU:HD12	1.74	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:240:ARG:HD3	1:A:298:PRO:CB	2.45	0.45
1:B:121:PRO:HB2	1:B:122:ALA:H	1.61	0.44
1:B:370:CYS:SG	1:B:378:LEU:HD13	2.58	0.43
1:B:126:LEU:O	1:B:130:ARG:HG3	2.17	0.43
6:A:880:GOL:HO3	6:A:880:GOL:HO2	1.50	0.42
1:A:70:PHE:HB3	1:A:84:ASP:O	2.20	0.42
1:A:423:HIS:HB2	1:B:392:THR:HB	2.02	0.41
1:B:322:LEU:HD13	1:B:324:TRP:CZ2	2.55	0.41
1:A:100:ARG:NH1	1:A:102:LEU:HD22	2.36	0.41
1:A:246:TRP:HB2	1:A:294:LEU:HB3	2.03	0.40
1:A:308:PRO:HA	1:A:309:PRO:HD3	1.99	0.40
1:B:150:GLU:O	1:B:154:GLU:HG3	2.21	0.40
1:B:180:TRP:CE3	1:B:192:TRP:HA	2.57	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:386:ASP:OD2	8:A:2073:HOH:O[3_555]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	413/443 (93%)	403 (98%)	10 (2%)	0	100	100
1	B	399/443 (90%)	390 (98%)	9 (2%)	0	100	100
All	All	812/886 (92%)	793 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	353/375 (94%)	341 (97%)	12 (3%)	44	26
1	B	343/375 (92%)	329 (96%)	14 (4%)	37	19
All	All	696/750 (93%)	670 (96%)	26 (4%)	41	23

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	69	LYS
1	A	100	ARG
1	A	102	LEU
1	A	105	LEU
1	A	114	ARG
1	A	125	LEU
1	A	146	GLN
1	A	259	GLN
1	A	293	LEU
1	A	294	LEU
1	A	328	LEU
1	A	331	ARG
1	B	69	LYS
1	B	91	GLN
1	B	92	GLN
1	B	100	ARG
1	B	102	LEU
1	B	124	GLN
1	B	190	ILE
1	B	258	GLN
1	B	278	GLN
1	B	330	LEU
1	B	331	ARG
1	B	378	LEU
1	B	389	THR
1	B	458	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CAS	A	384	1	4,8,9	1.05	0	2,9,11	1.28	0
1	CAS	B	384	1	4,8,9	1.11	0	2,9,11	1.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CAS	A	384	1	-	0/0/7/9	0/0/0/0
1	CAS	B	384	1	-	0/0/7/9	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 1 is monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	A	500	1	24,50,50	2.43	8 (33%)	16,82,82	1.69	2 (12%)
3	H4B	A	600	-	15,18,18	0.90	0	12,26,26	2.66	5 (41%)
4	7M3	A	800	-	25,25,25	0.80	0	34,34,34	1.15	3 (8%)
5	ACT	A	860	-	0,3,3	0.00	-	0,3,3	0.00	-
5	ACT	A	861	-	0,3,3	0.00	-	0,3,3	0.00	-
6	GOL	A	880	-	5,5,5	0.34	0	5,5,5	0.39	0
2	HEM	B	500	1	24,50,50	2.27	6 (25%)	16,82,82	2.19	3 (18%)
3	H4B	B	600	-	15,18,18	0.94	1 (6%)	12,26,26	2.56	5 (41%)
4	7M3	B	800	-	25,25,25	0.89	0	34,34,34	1.16	3 (8%)
5	ACT	B	860	-	0,3,3	0.00	-	0,3,3	0.00	-
5	ACT	B	861	-	0,3,3	0.00	-	0,3,3	0.00	-
6	GOL	B	880	-	5,5,5	0.36	0	5,5,5	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	500	1	-	0/6/54/54	0/0/8/8
3	H4B	A	600	-	-	0/8/17/17	0/2/2/2
4	7M3	A	800	-	-	0/9/9/9	0/3/3/3
5	ACT	A	860	-	-	0/0/0/0	0/0/0/0
5	ACT	A	861	-	-	0/0/0/0	0/0/0/0
6	GOL	A	880	-	-	0/4/4/4	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	B	500	1	-	0/6/54/54	0/0/8/8
3	H4B	B	600	-	-	0/8/17/17	0/2/2/2
4	7M3	B	800	-	-	0/9/9/9	0/3/3/3
5	ACT	B	860	-	-	0/0/0/0	0/0/0/0
5	ACT	B	861	-	-	0/0/0/0	0/0/0/0
6	GOL	B	880	-	-	0/4/4/4	0/0/0/0

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	HEM	C3B-C2B	-4.83	1.34	1.40
2	A	500	HEM	C3C-C2C	-4.64	1.34	1.40
2	B	500	HEM	C3B-C2B	-4.45	1.34	1.40
2	B	500	HEM	C3C-C2C	-4.26	1.34	1.40
3	B	600	H4B	C4-C4A	-2.30	1.38	1.41
2	B	500	HEM	CMD-C2D	2.18	1.56	1.51
2	A	500	HEM	C1B-NB	2.25	1.39	1.36
2	A	500	HEM	CAD-C3D	2.63	1.55	1.52
2	A	500	HEM	C4D-ND	2.94	1.40	1.36
2	B	500	HEM	C3B-CAB	3.42	1.55	1.47
2	A	500	HEM	C3B-CAB	3.48	1.55	1.47
2	A	500	HEM	C3C-CAC	3.83	1.55	1.47
2	B	500	HEM	C3C-CAC	3.95	1.56	1.47
2	A	500	HEM	C3D-C2D	4.97	1.52	1.37
2	B	500	HEM	C3D-C2D	5.16	1.52	1.37

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	500	HEM	CBA-CAA-C2A	-6.51	101.05	112.49
2	A	500	HEM	CBA-CAA-C2A	-4.92	103.85	112.49
3	A	600	H4B	N3-C2-N1	-3.51	119.78	125.51
3	B	600	H4B	N3-C2-N1	-3.30	120.13	125.51
2	B	500	HEM	CBD-CAD-C3D	-3.06	107.09	112.47
2	B	500	HEM	CMA-C3A-C4A	-2.51	124.04	128.31
2	A	500	HEM	CBD-CAD-C3D	-2.44	108.19	112.47
4	A	800	7M3	C21-C27-N28	-2.41	108.19	112.85
4	B	800	7M3	O12-C11-C08	-2.06	103.30	109.17
4	A	800	7M3	C02-N01-C10	2.25	120.36	117.60
3	B	600	H4B	N2-C2-N1	2.27	120.94	117.20
3	B	600	H4B	C2-N1-C8A	2.48	120.20	114.63
3	A	600	H4B	C4A-C8A-N8	2.52	120.93	118.26

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	800	7M3	N02-C02-N01	2.67	119.68	118.03
4	B	800	7M3	N02-C02-N01	2.71	119.71	118.03
4	B	800	7M3	C02-N01-C10	2.80	121.02	117.60
3	A	600	H4B	C2-N1-C8A	2.85	121.03	114.63
3	A	600	H4B	C4-N3-C2	4.51	121.17	115.88
3	B	600	H4B	C4-N3-C2	4.52	121.18	115.88
3	A	600	H4B	C4-C4A-C8A	5.25	119.23	114.61
3	B	600	H4B	C4-C4A-C8A	5.52	119.47	114.61

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	HEM	1	0
3	A	600	H4B	1	0
4	A	800	7M3	3	0
6	A	880	GOL	2	0
2	B	500	HEM	1	0
4	B	800	7M3	3	0
5	B	861	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	415/443 (93%)	0.31	38 (9%) 11 8	21, 32, 61, 112	1 (0%)
1	B	403/443 (90%)	0.13	20 (4%) 32 27	23, 36, 61, 106	0
All	All	818/886 (92%)	0.22	58 (7%) 19 15	21, 34, 62, 112	1 (0%)

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	119	PRO	9.0
1	A	117	PRO	8.7
1	B	259	GLN	7.8
1	A	116	SER	7.3
1	A	118	GLY	6.8
1	A	67	GLY	6.3
1	B	260	ASP	6.1
1	A	115	PRO	5.9
1	A	259	GLN	5.4
1	A	160	ALA	5.1
1	A	120	PRO	4.8
1	B	110	LYS	4.5
1	A	239	GLY	4.5
1	A	260	ASP	4.1
1	A	121	PRO	4.1
1	A	123	GLU	4.0
1	B	122	ALA	4.0
1	A	114	ARG	3.9
1	B	121	PRO	3.8
1	A	69	LYS	3.7
1	B	323	GLU	3.7
1	B	123	GLU	3.6
1	A	122	ALA	3.5
1	A	124	GLN	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	338	VAL	3.2
1	B	223	ARG	3.1
1	A	238	PRO	3.0
1	B	239	GLY	3.0
1	A	126	LEU	2.9
1	A	355	PHE	2.7
1	A	156	GLU	2.7
1	A	146	GLN	2.7
1	A	153	GLN	2.6
1	A	261	GLY	2.6
1	A	109	ARG	2.6
1	A	230	ILE	2.6
1	A	157	ALA	2.5
1	B	146	GLN	2.5
1	A	299	ASP	2.5
1	B	141	LYS	2.4
1	B	258	GLN	2.4
1	A	390	ARG	2.4
1	B	149	GLU	2.4
1	A	161	SER	2.4
1	B	70	PHE	2.3
1	B	91	GLN	2.3
1	B	108	PRO	2.3
1	A	111	LEU	2.3
1	A	91	GLN	2.2
1	B	261	GLY	2.2
1	A	130	ARG	2.2
1	B	142	ARG	2.2
1	B	262	SER	2.1
1	A	99	ARG	2.1
1	A	113	THR	2.1
1	A	449	TRP	2.0
1	A	285	ASN	2.0
1	B	280	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy

less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	CAS	A	384	9/10	0.91	0.10	-	29,34,58,106	0
1	CAS	B	384	9/10	0.93	0.12	-	45,49,79,138	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	ACT	B	860	4/4	0.94	0.13	3.95	44,45,47,55	0
4	7M3	B	800	23/23	0.91	0.24	3.06	24,44,90,91	0
5	ACT	A	861	4/4	0.87	0.19	2.75	36,41,41,43	0
5	ACT	A	860	4/4	0.89	0.15	2.64	42,45,49,55	0
6	GOL	A	880	6/6	0.84	0.21	2.10	55,61,67,71	0
6	GOL	B	880	6/6	0.93	0.15	1.48	50,57,57,58	0
4	7M3	A	800	23/23	0.93	0.19	1.00	25,39,81,82	0
2	HEM	A	500	43/43	0.98	0.16	0.81	19,23,39,48	0
2	HEM	B	500	43/43	0.98	0.13	0.48	20,24,43,52	0
5	ACT	B	861	4/4	0.90	0.16	0.43	41,47,48,48	0
3	H4B	B	600	17/17	0.94	0.10	-0.77	27,31,36,37	0
3	H4B	A	600	17/17	0.96	0.10	-1.04	24,32,36,36	0
7	ZN	A	900	1/1	1.00	0.04	-4.54	30,30,30,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.