



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 8, 2016 – 01:11 PM EST

PDB ID : 5AEG  
Title : A bacterial protein structure in glycoside hydrolase family 31.  
Authors : Jin, Y.; Speciale, G.; Davies, G.J.; Williams, S.J.; Goddard-Borger, E.D.  
Deposited on : 2015-08-30  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20028442  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20028442


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1745 (1.86-1.86)
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	686	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 98%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> <span>%</span> <span>90%</span> <span>7%</span> <span>.</span> </div> </div>
1	B	686	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 3%, orange 1%, yellow 1%, green 95%);"></div> <div style="display: flex; justify-content: space-between; width: 89%; margin: 0 auto;"> <span>3%</span> <span>89%</span> <span>8%</span> <span>.</span> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MPD	A	1682	-	-	-	X
4	MPD	A	1683	-	-	-	X
4	MPD	B	1680	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11534 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ALPHA-GLUCOSIDASE YIHQ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	667	Total	C	N	O	S	0	1	0
			5371	3445	894	1004	28			
1	B	670	Total	C	N	O	S	0	1	0
			5398	3458	902	1010	28			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	679	LEU	-	EXPRESSION TAG	UNP P32138
A	680	GLU	-	EXPRESSION TAG	UNP P32138
A	681	HIS	-	EXPRESSION TAG	UNP P32138
A	682	HIS	-	EXPRESSION TAG	UNP P32138
A	683	HIS	-	EXPRESSION TAG	UNP P32138
A	684	HIS	-	EXPRESSION TAG	UNP P32138
A	685	HIS	-	EXPRESSION TAG	UNP P32138
A	686	HIS	-	EXPRESSION TAG	UNP P32138
B	679	LEU	-	EXPRESSION TAG	UNP P32138
B	680	GLU	-	EXPRESSION TAG	UNP P32138
B	681	HIS	-	EXPRESSION TAG	UNP P32138
B	682	HIS	-	EXPRESSION TAG	UNP P32138
B	683	HIS	-	EXPRESSION TAG	UNP P32138
B	684	HIS	-	EXPRESSION TAG	UNP P32138
B	685	HIS	-	EXPRESSION TAG	UNP P32138
B	686	HIS	-	EXPRESSION TAG	UNP P32138


- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Ca	0	0
			1	1		
2	A	2	Total	Ca	0	0
			2	2		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			8	6	2		
4	A	1	Total	C	O	0	0
			8	6	2		
4	B	1	Total	C	O	0	0
			8	6	2		

- Molecule 5 is SUGAR ((2R,3R,4R,5S,6R)-6-FLUORANYL-6-(HYDROXYMETHYL)OXANE-2,3,4,5-TETROL) (three-letter code: B9D) (formula: C<sub>6</sub>H<sub>11</sub>FO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	F	O	0	0
			12	6	1	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	465	Total	O	0	0
			465	465		
6	B	259	Total	O	0	0
			259	259		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: ALPHA-GLUCOSIDASE YIHQ



#### • Molecule 1: ALPHA-GLUCOSIDASE YIHQ



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.67Å 112.64Å 111.88Å 90.00° 109.52° 90.00°	Depositor
Resolution (Å)	105.45 – 1.85 49.68 – 1.85	Depositor EDS
% Data completeness (in resolution range)	98.3 (105.45-1.85) 98.4 (49.68-1.85)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.57 (at 1.86Å)	Xtriage
Refinement program	REFMAC 5.8.0095	Depositor
R, $R_{free}$	0.160 , 0.193 0.172 , 0.200	Depositor DCC
$R_{free}$ test set	7578 reflections (5.18%)	DCC
Wilson B-factor (Å <sup>2</sup> )	27.1	Xtriage
Anisotropy	0.048	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 47.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.018 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	11534	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.77% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, B9D, CA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.11	3/5533 (0.1%)	1.03	25/7519 (0.3%)
1	B	0.89	2/5562 (0.0%)	0.92	10/7560 (0.1%)
All	All	1.01	5/11095 (0.0%)	0.98	35/15079 (0.2%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	266	GLU	CG-CD	5.66	1.60	1.51
1	A	611	ARG	CZ-NH1	5.66	1.40	1.33
1	B	492	SER	CA-CB	5.25	1.60	1.52
1	A	472	ASP	CB-CG	5.15	1.62	1.51
1	B	43	SER	CB-OG	-5.08	1.35	1.42

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	472	ASP	CB-CG-OD2	10.27	127.55	118.30
1	A	472	ASP	CB-CG-OD1	-9.87	109.41	118.30
1	B	88	ASP	CB-CG-OD1	8.99	126.39	118.30
1	A	481	ASP	CB-CG-OD1	8.61	126.05	118.30
1	B	50	MET	CG-SD-CE	-8.26	86.98	100.20
1	A	455	ARG	NE-CZ-NH1	7.93	124.27	120.30
1	A	295	MET	CG-SD-CE	-7.16	88.75	100.20
1	A	663	ARG	NE-CZ-NH2	-6.96	116.82	120.30
1	A	317	ASP	CB-CG-OD2	-6.91	112.08	118.30
1	B	611	ARG	NE-CZ-NH1	6.89	123.75	120.30
1	A	116	ARG	NE-CZ-NH1	6.79	123.69	120.30
1	A	625	ASP	CB-CG-OD1	6.74	124.37	118.30
1	B	623	ARG	NE-CZ-NH1	6.59	123.59	120.30
1	A	88	ASP	CB-CG-OD2	6.48	124.14	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	516	ARG	NE-CZ-NH1	6.36	123.48	120.30
1	B	663	ARG	NE-CZ-NH1	6.22	123.41	120.30
1	B	319	ARG	NE-CZ-NH1	5.98	123.29	120.30
1	A	405	ASP	CB-CG-OD1	5.97	123.68	118.30
1	A	394	MET	CG-SD-CE	-5.86	90.83	100.20
1	A	623	ARG	NE-CZ-NH1	5.81	123.20	120.30
1	A	140	ASP	CB-CG-OD1	5.66	123.39	118.30
1	A	341	ASP	CB-CG-OD1	5.59	123.33	118.30
1	B	481	ASP	CB-CG-OD1	5.50	123.25	118.30
1	A	357	ASP	CB-CG-OD2	-5.28	113.54	118.30
1	A	176	ASP	CB-CG-OD1	5.25	123.03	118.30
1	A	317	ASP	CB-CG-OD1	5.19	122.97	118.30
1	A	611	ARG	NE-CZ-NH2	-5.19	117.71	120.30
1	B	276	ARG	NE-CZ-NH2	-5.13	117.74	120.30
1	A	266	GLU	OE1-CD-OE2	-5.11	117.17	123.30
1	A	611	ARG	NE-CZ-NH1	5.09	122.84	120.30
1	A	198	ASP	CB-CG-OD1	5.08	122.88	118.30
1	A	527	PHE	CB-CG-CD1	5.04	124.33	120.80
1	A	59	ASP	CB-CG-OD2	-5.04	113.76	118.30
1	A	633	ASP	CB-CG-OD2	5.03	122.83	118.30
1	B	527	PHE	CB-CG-CD1	5.01	124.31	120.80

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5371	0	5090	8	0
1	B	5398	0	5107	25	0
2	A	2	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	16	0	28	0	0
4	B	8	0	14	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	12	0	9	0	0
6	A	465	0	0	3	0
6	B	259	0	0	1	0
All	All	11534	0	10248	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:512:PHE:O	1:B:513:GLU:CB	2.37	0.71
1:B:13:PHE:CD1	1:B:72:VAL:HG21	2.29	0.68
1:B:13:PHE:CD1	1:B:72:VAL:CG2	2.77	0.68
1:B:13:PHE:CE1	1:B:72:VAL:HG23	2.31	0.66
1:B:301:ARG:NH1	6:B:2163:HOH:O	2.27	0.63
1:B:251:ASP:OD1	1:B:573:LYS:NZ	2.28	0.62
1:B:512:PHE:O	1:B:513:GLU:HB2	2.00	0.60
1:B:13:PHE:CG	1:B:72:VAL:HG21	2.37	0.59
1:A:21:THR:HG22	1:A:31:THR:HG22	1.88	0.56
1:B:57:ILE:HD11	1:B:149:TRP:CD1	2.41	0.55
1:A:52:ARG:HD2	1:A:171:GLU:OE2	2.08	0.53
1:A:619:HIS:HD2	6:A:2412:HOH:O	1.92	0.53
1:A:648:GLU:HG2	6:A:2197:HOH:O	2.08	0.52
1:B:13:PHE:CZ	1:B:72:VAL:HG23	2.45	0.52
1:B:331:LEU:HA	1:B:401:GLY:O	2.10	0.51
1:B:494:ALA:HB1	1:B:585:PRO:HG3	1.95	0.48
1:B:301:ARG:NH2	1:B:367:PHE:O	2.47	0.47
1:A:64:LYS:HE3	1:A:118:TRP:CE2	2.50	0.47
1:B:13:PHE:CE1	1:B:72:VAL:CG2	2.98	0.46
1:B:13:PHE:CD1	1:B:72:VAL:HG23	2.48	0.46
1:B:515:LYS:HG2	1:B:544:ASP:HB3	1.98	0.46
1:B:13:PHE:CG	1:B:72:VAL:CG2	2.99	0.45
1:B:388:GLU:OE1	1:B:392:LYS:HE3	2.17	0.44
1:A:366:GLU:HG2	1:A:371:TYR:CE2	2.52	0.44
1:B:557:HIS:CE1	1:B:655:ILE:HG22	2.52	0.44
1:A:300:LYS:HE3	6:A:2263:HOH:O	2.18	0.43
1:A:8:LEU:HD23	1:A:9:LEU:HD12	2.01	0.42
1:B:194:TYR:CZ	1:B:227:GLU:HB2	2.55	0.42
1:B:75:SER:OG	1:B:78:GLY:O	2.29	0.42
1:B:453:PHE:HA	1:B:467:MET:O	2.21	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:468:MET:HG2	1:B:498:HIS:CE1	2.56	0.41
1:B:304:TRP:HA	1:B:306:TRP:CH2	2.56	0.40
1:B:72:VAL:HG12	1:B:73:SER:N	2.36	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	664/686 (97%)	646 (97%)	18 (3%)	0	100	100
1	B	669/686 (98%)	644 (96%)	23 (3%)	2 (0%)	46	29
All	All	1333/1372 (97%)	1290 (97%)	41 (3%)	2 (0%)	52	36

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	171	GLU
1	B	475	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	562/580 (97%)	549 (98%)	13 (2%)	58	41

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	565/580 (97%)	549 (97%)	16 (3%)	51	33
All	All	1127/1160 (97%)	1098 (97%)	29 (3%)	54	36

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	LEU
1	A	13	PHE
1	A	34	LYS
1	A	77	ASP
1	A	94	ASN
1	A	107	LEU
1	A	181	PHE
1	A	182	PHE
1	A	301	ARG
1	A	392	LYS
1	A	469	TRP
1	A	524	TRP
1	A	628	LEU
1	B	9	LEU
1	B	31	THR
1	B	58	LYS
1	B	94	ASN
1	B	99	ASP
1	B	182	PHE
1	B	221	LYS
1	B	307	LYS
1	B	392	LYS
1	B	393	ASN
1	B	432	LEU
1	B	469	TRP
1	B	513	GLU
1	B	581	LYS
1	B	596	ASP
1	B	628	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	GLN
1	B	15	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 5 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	MPD	A	1682	-	6,7,7	0.36	0	6,10,10	0.78	0
4	MPD	A	1683	-	6,7,7	0.42	0	6,10,10	1.25	0
4	MPD	B	1680	-	6,7,7	0.55	0	6,10,10	1.01	0
5	B9D	B	1681	1	12,12,13	1.11	1 (8%)	14,18,20	1.47	2 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MPD	A	1682	-	-	0/5/5/5	0/0/0/0
4	MPD	A	1683	-	-	0/5/5/5	0/0/0/0
4	MPD	B	1680	-	-	0/5/5/5	0/0/0/0
5	B9D	B	1681	1	-	0/2/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	1681	B9D	O-C5	2.48	1.47	1.39

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
5	B	1681	B9D	O-C1-C2	-2.14	107.60	111.02
5	B	1681	B9D	O3-C3-C2	2.71	114.97	110.01

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	667/686 (97%)	-0.58	8 (1%) 81 81	16, 26, 53, 87	1 (0%)
1	B	670/686 (97%)	-0.17	19 (2%) 56 54	24, 41, 64, 97	0
All	All	1337/1372 (97%)	-0.37	27 (2%) 68 67	16, 33, 61, 97	1 (0%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	13	PHE	5.8
1	B	76	PRO	5.3
1	B	298	PHE	4.9
1	A	99	ASP	4.0
1	B	99	ASP	3.9
1	B	75	SER	3.9
1	A	13	PHE	3.4
1	B	79	TRP	3.3
1	A	76	PRO	3.2
1	B	77	ASP	3.1
1	B	358	ALA	2.9
1	B	469	TRP	2.9
1	B	359	SER	2.7
1	B	11	PHE	2.7
1	B	97	ALA	2.6
1	A	298	PHE	2.5
1	A	77	ASP	2.4
1	A	73	SER	2.4
1	B	74	GLN	2.4
1	B	78	GLY	2.3
1	A	79	TRP	2.3
1	B	8	LEU	2.3
1	A	98	ASP	2.3
1	B	302	VAL	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	181	PHE	2.2
1	B	456	ALA	2.1
1	B	80	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	MPD	B	1680	8/8	0.88	0.25	6.29	68,72,76,77	0
4	MPD	A	1683	8/8	0.86	0.14	2.48	59,65,66,75	0
4	MPD	A	1682	8/8	0.93	0.12	2.43	45,53,62,69	0
2	CA	B	1678	1/1	1.00	0.11	0.10	27,27,27,27	0
2	CA	A	1680	1/1	1.00	0.07	0.02	18,18,18,18	0
5	B9D	B	1681	12/13	0.94	0.11	-1.57	32,38,46,50	0
3	CL	A	1681	1/1	0.99	0.03	-	28,28,28,28	0
2	CA	A	1679	1/1	0.99	0.07	-	33,33,33,33	0
3	CL	B	1679	1/1	0.99	0.03	-	39,39,39,39	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.