



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:27 AM GMT

PDB ID : 2AHV
Title : Crystal Structure of Acyl-CoA transferase from E. coli O157:H7 (YdiF)-thioester complex with CoA- 1
Authors : Rangarajan, E.S.; Li, Y.; Ajamian, E.; Iannuzzi, P.; Kernaghan, S.D.; Fraser, M.E.; Cygler, M.; Matte, A.; Montreal-Kingston Bacterial Structural Genomics Initiative (BSGI)
Deposited on : 2005-07-28
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	531	<div> <div>%</div> <div> <div></div> <div>88%</div> <div>8%</div> <div>.</div> </div> </div>
1	B	531	<div> <div>2%</div> <div> <div></div> <div>86%</div> <div>10%</div> <div>.</div> </div> </div>
1	C	531	<div> <div>%</div> <div> <div></div> <div>87%</div> <div>9%</div> <div>.</div> </div> </div>
1	D	531	<div> <div>%</div> <div> <div></div> <div>88%</div> <div>8%</div> <div>.</div> </div> </div>

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 17284 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called putative enzyme YdiF.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	512	Total	C	N	O	S	0	0	0
			3905	2486	666	737	16			
1	B	512	Total	C	N	O	S	0	0	0
			3902	2484	666	736	16			
1	C	512	Total	C	N	O	S	0	0	0
			3905	2486	666	737	16			
1	D	512	Total	C	N	O	S	0	0	0
			3906	2486	666	738	16			

- Molecule 2 is COENZYME A (three-letter code: COA) (formula: $C_{21}H_{36}N_7O_{16}P_3S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P S	0	0
			48	21	7	16	3 1		
2	D	1	Total	C	N	O	P S	0	0
			48	21	7	16	3 1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		
2	C	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	391	Total	O	0	0
			391	391		
3	B	323	Total	O	0	0
			323	323		
3	C	397	Total	O	0	0
			397	397		
3	D	363	Total	O	0	0
			363	363		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: putative enzyme YdiF




- Molecule 1: putative enzyme YdiF



- Molecule 1: putative enzyme YdiF



- Molecule 1: putative enzyme YdiF



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	81.08Å 140.24Å 112.68Å 90.00° 108.22° 90.00°	Depositor
Resolution (Å)	50.00 – 2.00 43.23 – 2.00	Depositor EDS
% Data completeness (in resolution range)	97.2 (50.00-2.00) 97.2 (43.23-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.182 , 0.224 0.184 , 0.225	Depositor DCC
R_{free} test set	14361 reflections (10.11%)	DCC
Wilson B-factor (Å ²)	23.2	Xtriage
Anisotropy	0.312	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.7	EDS
Estimated twinning fraction	0.017 for h,-k,-h-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 156371 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	17284	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: COA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.44	0/3971	0.74	10/5375 (0.2%)
1	B	0.45	0/3968	0.73	7/5372 (0.1%)
1	C	0.46	0/3971	0.74	8/5375 (0.1%)
1	D	0.45	0/3972	0.74	8/5377 (0.1%)
All	All	0.45	0/15882	0.74	33/21499 (0.2%)

There are no bond length outliers.

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	490	ASP	CB-CG-OD2	6.34	124.01	118.30
1	B	276	ASP	CB-CG-OD2	6.17	123.86	118.30
1	D	276	ASP	CB-CG-OD2	5.96	123.67	118.30
1	C	358	ASP	CB-CG-OD2	5.93	123.64	118.30
1	C	490	ASP	CB-CG-OD2	5.91	123.62	118.30
1	D	248	ASP	CB-CG-OD2	5.90	123.61	118.30
1	B	490	ASP	CB-CG-OD2	5.73	123.46	118.30
1	D	500	ASP	CB-CG-OD2	5.70	123.43	118.30
1	B	494	ASP	CB-CG-OD2	5.65	123.38	118.30
1	D	182	ASP	CB-CG-OD2	5.62	123.36	118.30
1	A	146	ASP	CB-CG-OD2	5.57	123.31	118.30
1	B	253	ASP	CB-CG-OD2	5.49	123.25	118.30
1	C	404	ASP	CB-CG-OD2	5.46	123.21	118.30
1	C	248	ASP	CB-CG-OD2	5.46	123.21	118.30
1	A	358	ASP	CB-CG-OD2	5.42	123.18	118.30
1	A	519	ASP	CB-CG-OD2	5.42	123.18	118.30
1	C	255	ASP	CB-CG-OD2	5.37	123.14	118.30
1	D	358	ASP	CB-CG-OD2	5.37	123.14	118.30
1	A	404	ASP	CB-CG-OD2	5.36	123.13	118.30
1	C	478	ASP	CB-CG-OD2	5.36	123.13	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	513	ASP	CB-CG-OD2	5.36	123.12	118.30
1	D	519	ASP	CB-CG-OD2	5.28	123.05	118.30
1	B	327	ASP	CB-CG-OD2	5.27	123.04	118.30
1	A	182	ASP	CB-CG-OD2	5.22	123.00	118.30
1	A	500	ASP	CB-CG-OD2	5.22	123.00	118.30
1	D	429	ASP	CB-CG-OD2	5.18	122.97	118.30
1	C	276	ASP	CB-CG-OD2	5.17	122.95	118.30
1	C	372	ASP	CB-CG-OD2	5.14	122.92	118.30
1	B	513	ASP	CB-CG-OD2	5.12	122.91	118.30
1	A	429	ASP	CB-CG-OD2	5.09	122.88	118.30
1	D	327	ASP	CB-CG-OD2	5.05	122.84	118.30
1	A	372	ASP	CB-CG-OD2	5.04	122.83	118.30
1	B	372	ASP	CB-CG-OD2	5.03	122.83	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3905	0	3963	17	0
1	B	3902	0	3959	26	0
1	C	3905	0	3963	21	0
1	D	3906	0	3963	19	0
2	A	48	0	31	1	0
2	B	48	0	32	1	0
2	C	48	0	31	0	0
2	D	48	0	32	1	0
3	A	391	0	0	0	0
3	B	323	0	0	2	0
3	C	397	0	0	1	0
3	D	363	0	0	2	0
All	All	17284	0	15974	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:184:ALA:HB2	1:D:215:VAL:HG21	1.72	0.71
1:C:495:ILE:HG22	1:C:499:MET:CE	2.22	0.70
1:B:416:GLY:O	1:B:468:THR:HA	1.96	0.64
1:C:120:VAL:HG13	1:C:135:ILE:HD11	1.78	0.64
1:C:495:ILE:HG22	1:C:499:MET:HE2	1.81	0.63
1:A:416:GLY:O	1:A:468:THR:HA	1.98	0.62
1:B:184:ALA:HB2	1:B:215:VAL:HG21	1.81	0.61
1:B:34:LEU:HD21	1:B:41:LEU:HD13	1.85	0.58
1:B:496:LEU:HA	1:B:499:MET:HE2	1.86	0.57
1:C:333:GLU:HG2	1:C:405:ILE:HD11	1.87	0.57
1:C:184:ALA:HB2	1:C:215:VAL:HG21	1.87	0.56
1:D:496:LEU:HA	1:D:499:MET:HE2	1.86	0.56
1:D:300:ARG:HD3	3:D:2845:HOH:O	2.05	0.56
1:A:184:ALA:HB2	1:A:215:VAL:HG21	1.87	0.56
1:A:136:ILE:HD11	1:D:136:ILE:HD11	1.88	0.55
1:C:495:ILE:HG22	1:C:499:MET:HE1	1.87	0.55
1:D:416:GLY:O	1:D:468:THR:HA	2.07	0.55
1:C:374:CYS:SG	1:C:405:ILE:HG22	2.47	0.54
1:B:472:VAL:HB	1:B:484:GLU:HB2	1.90	0.54
1:C:416:GLY:O	1:C:468:THR:HA	2.07	0.53
1:A:495:ILE:HG22	1:A:499:MET:CE	2.39	0.53
1:A:412:ILE:HD11	1:A:462:LEU:HD13	1.91	0.52
1:C:382:ASP:HA	1:C:444:ILE:O	2.10	0.52
1:B:266:ASN:HB3	1:B:269:ILE:HD12	1.93	0.51
1:B:437:GLU:HG2	1:B:498:LYS:HE3	1.93	0.51
1:B:156:GLU:O	1:B:159:LYS:HE3	2.11	0.50
1:B:410:LYS:N	1:B:410:LYS:HD2	2.27	0.50
1:A:305:GLY:HA2	1:A:373:VAL:O	2.12	0.50
1:D:444:ILE:HD11	1:D:447:LEU:HD23	1.94	0.49
1:B:307:VAL:CG1	1:B:313:ASP:HA	2.42	0.49
1:C:122:THR:HG23	1:C:360:THR:HG23	1.93	0.49
1:D:472:VAL:HB	1:D:484:GLU:HB2	1.95	0.49
1:C:384:HIS:NE2	1:C:446:GLU:OE2	2.43	0.49
1:B:489:VAL:HG13	1:B:494:ASP:HB2	1.94	0.48
1:B:410:LYS:HD3	3:B:2770:HOH:O	2.13	0.48
1:B:377:SER:HB3	2:B:2600:COA:H122	1.96	0.48
1:C:472:VAL:HB	1:C:484:GLU:HB2	1.95	0.47
1:A:401:GLY:O	1:A:405:ILE:HD12	2.14	0.47
1:C:489:VAL:HG13	1:C:494:ASP:HB2	1.98	0.46
1:A:112:ILE:HG23	1:A:158:THR:HG23	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:495:ILE:HG22	1:A:499:MET:HE2	1.98	0.45
1:D:420:ALA:HB2	1:D:442:LYS:HD2	1.99	0.45
1:C:376:LEU:CD1	1:C:406:SER:HB3	2.46	0.45
1:A:376:LEU:CD1	1:A:406:SER:HB3	2.47	0.45
1:A:122:THR:HG23	1:A:360:THR:HG23	1.98	0.45
1:B:299:MET:SD	1:B:303:ALA:HB3	2.57	0.45
1:B:499:MET:HE3	1:B:503:PRO:HD3	1.98	0.44
1:B:6:PRO:HD3	1:B:24:TYR:CE1	2.53	0.44
1:D:477:GLU:HG2	3:D:2885:HOH:O	2.17	0.44
1:B:444:ILE:HA	1:B:500:ASP:HB2	1.99	0.44
1:A:333:GLU:HG2	1:A:405:ILE:HD11	2.00	0.43
1:D:377:SER:HB3	2:D:2601:COA:H122	2.01	0.43
3:C:2903:HOH:O	1:D:265:VAL:HG11	2.18	0.43
1:C:124:THR:HG22	1:C:135:ILE:HG12	2.00	0.43
1:D:311:ILE:HG21	1:D:416:GLY:HA2	2.00	0.43
1:C:64:SER:HB3	1:C:90:TRP:CE3	2.54	0.43
1:B:387:VAL:C	1:B:447:LEU:HD21	2.39	0.43
1:D:496:LEU:HD23	1:D:499:MET:HE1	2.01	0.43
1:A:243:PRO:HD2	1:A:246:LEU:HD12	2.01	0.43
1:B:383:GLN:NE2	3:B:2795:HOH:O	2.35	0.43
1:A:405:ILE:HD13	2:A:2602:COA:S1P	2.59	0.42
1:C:431:LYS:HB2	1:C:431:LYS:HE3	1.77	0.42
1:B:412:ILE:HD11	1:B:462:LEU:HD13	2.01	0.42
1:A:495:ILE:HG22	1:A:499:MET:HE1	2.01	0.42
1:D:385:GLY:HA2	1:D:480:LEU:HD11	2.02	0.42
1:B:136:ILE:HD11	1:C:136:ILE:HD11	2.02	0.41
1:C:34:LEU:HD21	1:C:41:LEU:HD13	2.01	0.41
1:D:305:GLY:HA2	1:D:373:VAL:O	2.20	0.41
1:D:300:ARG:HG3	1:D:303:ALA:HB2	2.02	0.41
1:A:120:VAL:HG13	1:A:135:ILE:HD11	2.02	0.41
1:B:382:ASP:HB3	1:B:447:LEU:HD13	2.03	0.41
1:D:294:ARG:O	1:D:297:PHE:HB2	2.20	0.41
1:B:243:PRO:HD2	1:B:246:LEU:HD12	2.03	0.41
1:B:112:ILE:HG23	1:B:158:THR:HG23	2.02	0.41
1:C:120:VAL:O	1:C:124:THR:HG23	2.21	0.41
1:B:8:ARG:NH2	1:B:244:GLY:O	2.54	0.41
1:C:412:ILE:HD11	1:C:462:LEU:HD13	2.02	0.41
1:D:6:PRO:HD3	1:D:24:TYR:CE1	2.55	0.41
1:A:332:VAL:HB	1:A:336:PRO:HD2	2.04	0.40
1:B:180:ALA:HA	1:B:181:PRO:HD3	1.96	0.40
1:D:496:LEU:HD21	1:D:505:ILE:HD11	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	506/531 (95%)	492 (97%)	13 (3%)	1 (0%)	52	48
1	B	506/531 (95%)	491 (97%)	14 (3%)	1 (0%)	52	48
1	C	506/531 (95%)	494 (98%)	12 (2%)	0	100	100
1	D	506/531 (95%)	494 (98%)	11 (2%)	1 (0%)	52	48
All	All	2024/2124 (95%)	1971 (97%)	50 (2%)	3 (0%)	56	53

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	75	ALA
1	B	75	ALA
1	D	75	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	416/431 (96%)	409 (98%)	7 (2%)	68	71
1	B	415/431 (96%)	408 (98%)	7 (2%)	68	71
1	C	416/431 (96%)	407 (98%)	9 (2%)	60	62
1	D	416/431 (96%)	406 (98%)	10 (2%)	57	58

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1663/1724 (96%)	1630 (98%)	33 (2%)	63 65

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	101	ARG
1	A	115	ASN
1	A	171	LYS
1	A	374	CYS
1	A	410	LYS
1	A	451	THR
1	A	465	ARG
1	B	101	ARG
1	B	115	ASN
1	B	170	ASN
1	B	171	LYS
1	B	410	LYS
1	B	435	VAL
1	B	465	ARG
1	C	101	ARG
1	C	115	ASN
1	C	157	VAL
1	C	171	LYS
1	C	374	CYS
1	C	410	LYS
1	C	418	LEU
1	C	439	ARG
1	C	465	ARG
1	D	4	VAL
1	D	101	ARG
1	D	115	ASN
1	D	171	LYS
1	D	300	ARG
1	D	309	VAL
1	D	410	LYS
1	D	418	LEU
1	D	451	THR
1	D	465	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	10	ASN
1	A	287	GLN
1	A	350	ASN
1	B	287	GLN
1	B	350	ASN
1	C	287	GLN
1	C	350	ASN
1	C	383	GLN
1	C	436	GLN
1	D	287	GLN
1	D	350	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	COA	A	2602	1	40,50,50	0.85	1 (2%)	50,75,75	1.62	7 (14%)
2	COA	B	2600	-	40,50,50	1.73	3 (7%)	50,75,75	1.83	2 (4%)
2	COA	C	2603	1	40,50,50	0.88	1 (2%)	50,75,75	1.55	6 (12%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	COA	D	2601	-	40,50,50	1.73	3 (7%)	50,75,75	1.90	3 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	COA	A	2602	1	-	0/44/64/64	0/3/3/3
2	COA	B	2600	-	-	0/44/64/64	0/3/3/3
2	COA	C	2603	1	-	0/44/64/64	0/3/3/3
2	COA	D	2601	-	-	0/44/64/64	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	2601	COA	C2A-N1A	2.41	1.38	1.33
2	B	2600	COA	C2A-N1A	2.50	1.38	1.33
2	A	2602	COA	C5A-C4A	3.12	1.47	1.40
2	C	2603	COA	C5A-C4A	3.24	1.47	1.40
2	D	2601	COA	C2A-N3A	3.64	1.38	1.32
2	B	2600	COA	C2A-N3A	3.73	1.38	1.32
2	B	2600	COA	O9P-C9P	9.28	1.41	1.23
2	D	2601	COA	O9P-C9P	9.37	1.41	1.23

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2601	COA	N3A-C2A-N1A	-11.90	119.78	128.89
2	B	2600	COA	N3A-C2A-N1A	-11.64	119.98	128.89
2	A	2602	COA	N3A-C2A-N1A	-7.93	122.82	128.89
2	C	2603	COA	N3A-C2A-N1A	-6.82	123.67	128.89
2	C	2603	COA	C2B-C1B-N9A	-3.98	108.21	114.29
2	A	2602	COA	C2B-C1B-N9A	-2.74	110.11	114.29
2	C	2603	COA	C4A-C5A-N7A	-2.69	107.00	109.48
2	A	2602	COA	C4A-C5A-N7A	-2.65	107.04	109.48
2	D	2601	COA	P2A-O3A-P1A	-2.57	125.52	132.73
2	B	2600	COA	P2A-O3A-P1A	-2.30	126.28	132.73
2	C	2603	COA	C1B-N9A-C4A	-2.16	123.69	126.94
2	A	2602	COA	P2A-O3A-P1A	-2.04	127.01	132.73
2	C	2603	COA	C6P-C7P-N8P	2.00	116.27	111.88
2	A	2602	COA	O4B-C1B-N9A	2.12	112.54	108.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2602	COA	C2A-N1A-C6A	2.23	122.75	118.77
2	D	2601	COA	O4B-C1B-N9A	2.24	112.79	108.10
2	A	2602	COA	O6A-CCP-CBP	2.44	114.48	110.55
2	C	2603	COA	O4B-C1B-N9A	2.74	113.83	108.10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2602	COA	1	0
2	B	2600	COA	1	0
2	D	2601	COA	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	512/531 (96%)	-0.25	5 (0%) 84 84	13, 22, 35, 43	0
1	B	512/531 (96%)	-0.05	9 (1%) 71 72	15, 27, 40, 47	0
1	C	512/531 (96%)	-0.27	4 (0%) 87 88	13, 22, 34, 45	0
1	D	512/531 (96%)	-0.27	4 (0%) 87 88	13, 24, 39, 51	0
All	All	2048/2124 (96%)	-0.21	22 (1%) 82 83	13, 24, 38, 51	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	429	ASP	4.2
1	B	4	VAL	3.9
1	C	156	GLU	3.7
1	D	429	ASP	3.5
1	B	428	ALA	3.5
1	D	435	VAL	3.0
1	A	429	ASP	2.9
1	B	309	VAL	2.8
1	D	4	VAL	2.8
1	B	170	ASN	2.7
1	A	393	ASN	2.5
1	B	507	PRO	2.4
1	D	422	SER	2.4
1	B	429	ASP	2.3
1	A	428	ALA	2.3
1	A	361	SER	2.3
1	A	9	ILE	2.2
1	C	507	PRO	2.2
1	C	56	GLN	2.2
1	B	445	ARG	2.1
1	B	169	ASP	2.1
1	B	514	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	COA	D	2601	48/48	0.86	0.15	0.78	53,58,63,64	0
2	COA	B	2600	48/48	0.80	0.17	0.71	60,63,73,74	0
2	COA	A	2602	48/48	0.91	0.12	-0.05	40,44,52,54	0
2	COA	C	2603	48/48	0.95	0.11	-0.19	27,33,45,46	0

6.5 Other polymers [i](#)

There are no such residues in this entry.