



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:21 AM GMT

PDB ID : 3AHQ  
Title : hyperactive human Ero1  
Authors : Inaba, K.; Sitia, R.; Suzuki, M.  
Deposited on : 2010-04-26  
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1352 (2.38-2.34)
Clashscore	102246	1456 (2.38-2.34)
Ramachandran outliers	100387	1435 (2.38-2.34)
Sidechain outliers	100360	1436 (2.38-2.34)
RSRZ outliers	91569	1358 (2.38-2.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	465	<div> <div>11%</div> <div>63%</div> <div>11%</div> <div>23%</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3091 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ERO1-like protein alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	356	Total	C	N	O	S	0	0	0
			2973	1914	508	538	13			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	4	MET	-	EXPRESSION TAG	UNP Q96HE7
A	5	GLY	-	EXPRESSION TAG	UNP Q96HE7
A	6	SER	-	EXPRESSION TAG	UNP Q96HE7
A	7	SER	-	EXPRESSION TAG	UNP Q96HE7
A	8	HIS	-	EXPRESSION TAG	UNP Q96HE7
A	9	HIS	-	EXPRESSION TAG	UNP Q96HE7
A	10	HIS	-	EXPRESSION TAG	UNP Q96HE7
A	11	HIS	-	EXPRESSION TAG	UNP Q96HE7
A	12	HIS	-	EXPRESSION TAG	UNP Q96HE7
A	13	HIS	-	EXPRESSION TAG	UNP Q96HE7
A	14	SER	-	EXPRESSION TAG	UNP Q96HE7
A	15	SER	-	EXPRESSION TAG	UNP Q96HE7
A	16	GLY	-	EXPRESSION TAG	UNP Q96HE7
A	17	HIS	-	EXPRESSION TAG	UNP Q96HE7
A	18	ILE	-	EXPRESSION TAG	UNP Q96HE7
A	19	GLU	-	EXPRESSION TAG	UNP Q96HE7
A	20	GLY	-	EXPRESSION TAG	UNP Q96HE7
A	21	ARG	-	EXPRESSION TAG	UNP Q96HE7
A	104	ALA	CYS	ENGINEERED MUTATION	UNP Q96HE7
A	131	ALA	CYS	ENGINEERED MUTATION	UNP Q96HE7
A	166	ALA	CYS	ENGINEERED MUTATION	UNP Q96HE7

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	65	Total	O	0	0
			65	65		

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Chain A: 

## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.96 Å   140.73 Å   144.85 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	36.27 – 2.35 36.21 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.2 (36.27-2.35) 99.2 (36.21-2.35)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.59 (at 2.34 Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.237 , 0.282 0.231 , 0.276	Depositor DCC
$R_{free}$ test set	1221 reflections (5.43%)	DCC
Wilson B-factor (Å <sup>2</sup> )	51.0	Xtriage
Anisotropy	0.064	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 44.7	EDS
Estimated twinning fraction	0.000 for -h,-l,-k	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 23792 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3091	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.69% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.88	2/3046 (0.1%)	0.79	1/4103 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	400	TRP	CE3-CZ3	5.84	1.48	1.38
1	A	294	GLU	CG-CD	5.02	1.59	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	187	ARG	NE-CZ-NH2	-5.47	117.56	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2973	0	2901	37	0
2	A	53	0	31	0	0
3	A	65	0	0	1	0
All	All	3091	0	2932	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:383:ARG:HG2	1:A:383:ARG:HH11	1.29	0.94
1:A:36:PHE:CZ	1:A:359:ASP:OD1	2.25	0.90
1:A:202:VAL:HG12	1:A:388:ILE:CD1	2.11	0.80
1:A:383:ARG:NH1	1:A:383:ARG:HG2	1.96	0.80
1:A:202:VAL:HG12	1:A:388:ILE:HD13	1.66	0.74
1:A:210:LYS:CE	1:A:373:LYS:HD2	2.21	0.71
1:A:270:GLU:HG3	1:A:275:LYS:HG2	1.73	0.69
1:A:134:ALA:HA	1:A:137:LEU:HB2	1.75	0.67
1:A:383:ARG:CG	1:A:383:ARG:HH11	2.02	0.67
1:A:281:ILE:HD11	1:A:463:LEU:HB3	1.84	0.59
1:A:424:MET:HE2	1:A:433:PHE:HB3	1.86	0.57
1:A:400:TRP:CZ3	1:A:449:ARG:HD3	2.41	0.56
1:A:210:LYS:HE2	1:A:373:LYS:HD2	1.87	0.55
1:A:207:ASN:ND2	1:A:243:GLU:HB2	2.23	0.53
1:A:383:ARG:NH1	1:A:384:ASN:OD1	2.43	0.50
1:A:36:PHE:HZ	1:A:359:ASP:OD1	1.92	0.50
1:A:424:MET:HB2	1:A:424:MET:HE3	1.71	0.48
1:A:162:SER:O	1:A:164:ASN:N	2.45	0.47
1:A:210:LYS:HE3	1:A:373:LYS:HD2	1.96	0.47
1:A:266:TYR:O	1:A:277:TRP:HA	2.14	0.47
1:A:461:ARG:HH11	1:A:461:ARG:HG3	1.81	0.46
1:A:306:PHE:O	1:A:310:ILE:HG12	2.16	0.46
1:A:76:TYR:HA	1:A:179:VAL:O	2.16	0.45
1:A:38:GLN:HG3	1:A:40:SER:H	1.81	0.45
1:A:64:ARG:HH11	1:A:64:ARG:HG3	1.83	0.44
1:A:304:LEU:HD11	1:A:453:SER:HB3	2.00	0.44
1:A:164:ASN:HB2	3:A:517:HOH:O	2.18	0.44
1:A:202:VAL:HG12	1:A:388:ILE:HD11	1.95	0.44
1:A:419:LYS:HE3	1:A:419:LYS:HB2	1.89	0.43
1:A:323:GLU:HG2	1:A:345:LEU:HD22	2.00	0.42
1:A:60:ARG:HB3	1:A:60:ARG:HE	1.46	0.42
1:A:83:ARG:HD3	1:A:390:ASP:OD2	2.20	0.42
1:A:424:MET:HA	1:A:425:PRO:HD2	1.81	0.41
1:A:207:ASN:CG	1:A:243:GLU:HB2	2.40	0.41
1:A:312:LEU:HG	1:A:351:ILE:HG23	2.01	0.41
1:A:313:ARG:HG3	1:A:415:LEU:HD21	2.02	0.41
1:A:246:ALA:O	1:A:250:LEU:HG	2.21	0.40



There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	348/465 (75%)	330 (95%)	17 (5%)	1 (0%)	46 55

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	163	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	324/415 (78%)	306 (94%)	18 (6%)	26 31

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	GLN
1	A	44	ASP
1	A	60	ARG
1	A	162	SER
1	A	163	ASP
1	A	208	CYS
1	A	245	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	272	TRP
1	A	273	LEU
1	A	329	LEU
1	A	353	SER
1	A	362	SER
1	A	363	PHE
1	A	368	LYS
1	A	373	LYS
1	A	380	LEU
1	A	383	ARG
1	A	449	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	57	ASN
1	A	158	HIS
1	A	333	ASN
1	A	361	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FAD	A	1	-	48,58,58	1.33	6 (12%)	54,89,89	2.68	15 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	1	-	-	0/30/50/50	0/6/6/6

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	FAD	C1'-N10	2.23	1.50	1.48
2	A	1	FAD	C5X-N5	2.28	1.38	1.35
2	A	1	FAD	C4-N3	2.35	1.37	1.33
2	A	1	FAD	C2A-N3A	3.42	1.38	1.32
2	A	1	FAD	C2A-N1A	3.47	1.40	1.33
2	A	1	FAD	C4X-N5	4.07	1.39	1.33

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	FAD	N3A-C2A-N1A	-14.17	118.05	128.89
2	A	1	FAD	C1B-N9A-C4A	-3.31	121.95	126.94
2	A	1	FAD	O3B-C3B-C4B	-2.74	102.83	111.05
2	A	1	FAD	C2B-C1B-N9A	-2.46	110.54	114.29
2	A	1	FAD	C9A-C5X-N5	-2.44	118.74	122.36
2	A	1	FAD	C4X-C4-N3	-2.44	120.25	123.59
2	A	1	FAD	C4-C4X-C10	-2.40	118.41	119.94
2	A	1	FAD	O3B-C3B-C2B	-2.03	105.21	111.83
2	A	1	FAD	C4-C4X-N5	2.17	121.36	118.72
2	A	1	FAD	C2A-N1A-C6A	2.51	123.26	118.77
2	A	1	FAD	C4X-N5-C5X	2.84	120.03	116.76
2	A	1	FAD	O3P-P-O5'	3.15	111.28	102.94
2	A	1	FAD	C1'-N10-C9A	4.16	123.53	118.86
2	A	1	FAD	C5X-C9A-N10	5.08	121.48	117.62

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	FAD	C4-N3-C2	5.78	120.25	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	356/465 (76%)	0.74	49 (13%) 4 8	33, 56, 87, 99	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	272	TRP	10.3
1	A	173	SER	6.9
1	A	369	LYS	5.6
1	A	335	ILE	5.3
1	A	240	LEU	5.3
1	A	239	GLY	5.2
1	A	42	TYR	4.6
1	A	165	PHE	4.4
1	A	241	CYS	4.4
1	A	430	SER	3.9
1	A	211	PRO	3.8
1	A	342	MET	3.7
1	A	250	LEU	3.3
1	A	325	PRO	3.2
1	A	425	PRO	3.2
1	A	43	LEU	3.2
1	A	155	TRP	3.1
1	A	431	TYR	3.1
1	A	366	GLY	3.0
1	A	408	LEU	3.0
1	A	429	PRO	2.8
1	A	307	LEU	2.8
1	A	423	ASN	2.7
1	A	164	ASN	2.7
1	A	304	LEU	2.7
1	A	447	PHE	2.7
1	A	150	LYS	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	324	ARG	2.6
1	A	210	LYS	2.6
1	A	315	LEU	2.6
1	A	271	THR	2.6
1	A	426	GLU	2.5
1	A	407	GLY	2.5
1	A	251	ILE	2.5
1	A	146	GLU	2.5
1	A	370	GLU	2.5
1	A	40	SER	2.4
1	A	89	ASN	2.3
1	A	412	LEU	2.3
1	A	443	LEU	2.3
1	A	36	PHE	2.2
1	A	314	ALA	2.2
1	A	310	ILE	2.2
1	A	208	CYS	2.2
1	A	414	ILE	2.1
1	A	444	PHE	2.1
1	A	273	LEU	2.1
1	A	135	GLU	2.1
1	A	411	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	FAD	A	1	53/53	0.98	0.12	-0.58	28,35,43,47	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.