



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:50 PM GMT

PDB ID : 4AM0
Title : Structure of Dengue virus strain 4 DIII in complex with Fab 2H12
Authors : Midgley, C.M.; Flanagan, A.; Mongkolsapaya, J.; Grimes, J.M.; Screaton, G.R.
Deposited on : 2012-03-06
Resolution : 3.02 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	2117 (3.04-3.00)
Ramachandran outliers	100387	2050 (3.04-3.00)
Sidechain outliers	100360	2053 (3.04-3.00)
RSRZ outliers	91569	1788 (3.04-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	217	
1	C	217	
1	E	217	
1	H	217	
2	B	212	
2	D	212	
2	F	212	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	L	212	<div><div></div><div>83%</div><div>14%</div><div></div></div>
3	Q	101	<div><div></div><div>61%</div><div>27%</div><div></div><div>•</div><div>10%</div></div>
3	R	101	<div><div></div><div>68%</div><div>22%</div><div></div><div>5%</div><div>5%</div></div>
3	S	101	<div><div></div><div>63%</div><div>26%</div><div></div><div>•</div><div>9%</div></div>
3	T	101	<div><div></div><div>71%</div><div>18%</div><div></div><div>5%</div><div>6%</div></div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 15830 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FAB 2H12, HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	211	Total	C	N	O	S	0	0	0
			1580	996	255	321	8			
1	C	211	Total	C	N	O	S	0	0	0
			1580	996	255	321	8			
1	E	212	Total	C	N	O	S	0	0	0
			1584	998	256	322	8			
1	H	214	Total	C	N	O	S	0	0	0
			1596	1005	258	325	8			

- Molecule 2 is a protein called FAB 2H12, LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	212	Total	C	N	O	S	0	0	0
			1655	1031	283	334	7			
2	D	212	Total	C	N	O	S	0	0	0
			1655	1031	283	334	7			
2	F	212	Total	C	N	O	S	0	0	0
			1655	1031	283	334	7			
2	L	212	Total	C	N	O	S	0	0	0
			1655	1031	283	334	7			


- Molecule 3 is a protein called ENVELOPE PROTEIN,.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	Q	91	Total	C	N	O	S	0	0	0
			696	443	114	135	4			
3	R	96	Total	C	N	O	S	0	0	0
			736	468	122	142	4			
3	S	92	Total	C	N	O	S	0	0	0
			706	451	116	135	4			
3	T	95	Total	C	N	O	S	0	0	0
			732	466	121	141	4			


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: FAB 2H12, HEAVY CHAIN




• Molecule 1: FAB 2H12, HEAVY CHAIN




• Molecule 1: FAB 2H12, HEAVY CHAIN




• Molecule 1: FAB 2H12, HEAVY CHAIN




- Molecule 2: FAB 2H12, LIGHT CHAIN




- Molecule 2: FAB 2H12, LIGHT CHAIN




- Molecule 2: FAB 2H12, LIGHT CHAIN



- Molecule 2: FAB 2H12, LIGHT CHAIN




- Molecule 3: ENVELOPE PROTEIN,




- Molecule 3: ENVELOPE PROTEIN,






● Molecule 3: ENVELOPE PROTEIN,



● Molecule 3: ENVELOPE PROTEIN,



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	69.93 Å 92.80 Å 96.59 Å 118.55° 90.33° 104.05°	Depositor
Resolution (Å)	41.99 – 3.02 41.99 – 3.02	Depositor EDS
% Data completeness (in resolution range)	95.7 (41.99-3.02) 79.5 (41.99-3.02)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 3.01 Å)	Xtriage
Refinement program	BUSTER 2.11.2	Depositor
R, R_{free}	0.196 , 0.252 0.215 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	38.7	Xtriage
Anisotropy	0.866	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 17.4	EDS
Estimated twinning fraction	0.097 for -h,-k-l,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	0 of 38469 reflections	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	15830	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/1622	0.78	0/2217
1	C	0.57	0/1622	0.79	0/2217
1	E	0.55	0/1626	0.79	0/2222
1	H	0.55	0/1638	0.79	0/2239
2	B	0.51	0/1693	0.72	0/2299
2	D	0.47	0/1693	0.69	0/2299
2	F	0.48	0/1693	0.70	0/2299
2	L	0.48	0/1693	0.69	0/2299
3	Q	0.57	0/710	0.82	0/963
3	R	0.56	0/751	0.81	0/1016
3	S	0.56	0/720	0.76	0/974
3	T	0.56	0/747	0.85	1/1011 (0.1%)
All	All	0.53	0/16208	0.76	1/22055 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	T	345	GLU	C-N-CA	6.18	137.15	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1580	0	1526	27	0
1	C	1580	0	1526	27	0
1	E	1584	0	1529	35	0
1	H	1596	0	1538	30	0
2	B	1655	0	1588	15	0
2	D	1655	0	1588	13	0
2	F	1655	0	1588	17	0
2	L	1655	0	1588	16	0
3	Q	696	0	685	15	0
3	R	736	0	734	14	0
3	S	706	0	705	15	0
3	T	732	0	731	12	0
All	All	15830	0	15326	221	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 221 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:208:THR:HG22	1:H:208:THR:HG22	1.20	1.18
1:C:208:THR:HG22	1:E:208:THR:HG22	1.38	1.06
1:E:97:THR:HG21	1:E:104:MET:HG3	1.37	1.06
1:E:97:THR:HG21	1:E:104:MET:CG	1.95	0.95
2:F:134:CYS:HG	2:F:194:CYS:HG	0.91	0.87

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	207/217 (95%)	193 (93%)	11 (5%)	3 (1%)	14 50

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	207/217 (95%)	194 (94%)	10 (5%)	3 (1%)	14	50
1	E	208/217 (96%)	193 (93%)	11 (5%)	4 (2%)	10	41
1	H	210/217 (97%)	195 (93%)	11 (5%)	4 (2%)	10	41
2	B	210/212 (99%)	195 (93%)	13 (6%)	2 (1%)	19	59
2	D	210/212 (99%)	196 (93%)	12 (6%)	2 (1%)	19	59
2	F	210/212 (99%)	197 (94%)	12 (6%)	1 (0%)	34	75
2	L	210/212 (99%)	197 (94%)	12 (6%)	1 (0%)	34	75
3	Q	87/101 (86%)	83 (95%)	3 (3%)	1 (1%)	17	56
3	R	94/101 (93%)	87 (93%)	7 (7%)	0	100	100
3	S	88/101 (87%)	82 (93%)	5 (6%)	1 (1%)	17	56
3	T	93/101 (92%)	88 (95%)	4 (4%)	1 (1%)	17	56
All	All	2034/2120 (96%)	1900 (93%)	111 (6%)	23 (1%)	17	56

5 of 23 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	103	ALA
2	D	83	LEU
1	H	103	ALA
3	T	346	LYS
1	A	189	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/186 (98%)	169 (93%)	13 (7%)	18	53
1	C	182/186 (98%)	168 (92%)	14 (8%)	16	48
1	E	182/186 (98%)	166 (91%)	16 (9%)	12	41
1	H	183/186 (98%)	168 (92%)	15 (8%)	14	45
2	B	189/189 (100%)	172 (91%)	17 (9%)	12	40

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	D	189/189 (100%)	175 (93%)	14 (7%)	17	50
2	F	189/189 (100%)	175 (93%)	14 (7%)	17	50
2	L	189/189 (100%)	174 (92%)	15 (8%)	15	47
3	Q	78/86 (91%)	73 (94%)	5 (6%)	22	58
3	R	82/86 (95%)	74 (90%)	8 (10%)	10	35
3	S	79/86 (92%)	73 (92%)	6 (8%)	16	49
3	T	82/86 (95%)	76 (93%)	6 (7%)	17	51
All	All	1806/1844 (98%)	1663 (92%)	143 (8%)	15	47

5 of 143 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	72	ILE
2	F	116	SER
3	S	322	VAL
1	E	97	THR
1	E	196	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	102	HIS
1	E	50	ASN
3	R	316	GLN
1	C	59	ASN
3	Q	316	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	211/217 (97%)	-0.69	0	100	100	14, 39, 71, 106	0
1	C	211/217 (97%)	-0.70	0	100	100	12, 39, 68, 103	0
1	E	212/217 (97%)	-0.66	1 (0%)	91	76	12, 38, 75, 125	0
1	H	214/217 (98%)	-0.66	0	100	100	16, 38, 76, 113	0
2	B	212/212 (100%)	-0.68	0	100	100	19, 46, 73, 109	0
2	D	212/212 (100%)	-0.65	0	100	100	20, 53, 71, 91	0
2	F	212/212 (100%)	-0.61	0	100	100	25, 51, 72, 102	0
2	L	212/212 (100%)	-0.59	0	100	100	19, 48, 78, 90	0
3	Q	91/101 (90%)	-0.75	0	100	100	24, 36, 56, 60	0
3	R	96/101 (95%)	-0.75	0	100	100	19, 39, 53, 58	0
3	S	92/101 (91%)	-0.76	0	100	100	20, 37, 53, 68	0
3	T	95/101 (94%)	-0.73	0	100	100	24, 36, 56, 66	0
All	All	2070/2120 (97%)	-0.67	1 (0%)	100	100	12, 43, 73, 125	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	217	ARG	2.2

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.