



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:09 PM GMT

PDB ID : 5AMO
Title : Structure of a mouse Olfactomedin-1 disulfide-linked dimer of the Olfactomedin domain and part of the coiled coil
Authors : Pronker, M.F.; Bos, T.G.A.A.; Sharp, T.H.; Thies-Weesie, D.M.; Janssen, B.J.C.
Deposited on : 2015-03-11
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	471	
1	B	471	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	1473	-	-	-	X
3	NAG	A	1394	-	-	-	X
3	NAG	B	1473	-	-	-	X
4	GOL	B	1482	-	-	-	X
4	GOL	B	1483	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4304 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NOELIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	253	Total	C	N	O	S	0	0	0
			2044	1301	347	386	10			
1	B	257	Total	C	N	O	S	0	0	0
			2075	1319	356	390	10			


There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	479	ALA	-	EXPRESSION TAG	UNP O88998
A	480	ALA	-	EXPRESSION TAG	UNP O88998
A	481	ALA	-	EXPRESSION TAG	UNP O88998
A	482	HIS	-	EXPRESSION TAG	UNP O88998
A	483	HIS	-	EXPRESSION TAG	UNP O88998
A	484	HIS	-	EXPRESSION TAG	UNP O88998
A	485	HIS	-	EXPRESSION TAG	UNP O88998
A	486	HIS	-	EXPRESSION TAG	UNP O88998
A	487	HIS	-	EXPRESSION TAG	UNP O88998
A	329	THR	ALA	VARIANT	UNP O88998
B	479	ALA	-	EXPRESSION TAG	UNP O88998
B	480	ALA	-	EXPRESSION TAG	UNP O88998
B	481	ALA	-	EXPRESSION TAG	UNP O88998
B	482	HIS	-	EXPRESSION TAG	UNP O88998
B	483	HIS	-	EXPRESSION TAG	UNP O88998
B	484	HIS	-	EXPRESSION TAG	UNP O88998
B	485	HIS	-	EXPRESSION TAG	UNP O88998
B	486	HIS	-	EXPRESSION TAG	UNP O88998
B	487	HIS	-	EXPRESSION TAG	UNP O88998
B	329	THR	ALA	VARIANT	UNP O88998

- Molecule 2 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	2	Total	C	N	O	0	0
			28	16	2	10		
2	A	2	Total	C	N	O	0	0
			28	16	2	10		
2	B	2	Total	C	N	O	0	0
			28	16	2	10		
2	B	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Cl	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	13	Total	O	0	0
			13	13		
6	B	13	Total	O	0	0
			13	13		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NOELIN



• Molecule 1: NOELIN





4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	160.22Å 43.94Å 104.06Å 90.00° 114.17° 90.00°	Depositor
Resolution (Å)	50.32 – 2.40 50.32 – 2.40	Depositor EDS
% Data completeness (in resolution range)	94.7 (50.32-2.40) 94.3 (50.32-2.40)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.32 (at 2.39Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.237 , 0.258 0.245 , 0.251	Depositor DCC
R_{free} test set	1252 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	49.9	Xtriage
Anisotropy	0.864	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 24878 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4304	wwPDB-VP
Average B, all atoms (Å ²)	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.21	0/2098	0.38	0/2851
1	B	0.21	0/2130	0.38	0/2894
All	All	0.21	0/4228	0.38	0/5745

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2044	0	1953	16	0
1	B	2075	0	1983	21	0
2	A	56	0	50	0	0
2	B	56	0	50	0	0
3	A	14	0	13	0	0
3	B	14	0	13	0	0
4	B	18	0	24	1	0
5	B	1	0	0	0	0
6	A	13	0	0	0	0
6	B	13	0	0	0	0
All	All	4304	0	4086	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:264:TYR:HB3	1:A:275:ARG:HB2	1.61	0.81
1:B:228:GLY:HA2	1:B:478:HIS:HB3	1.66	0.77
1:A:461:LEU:HB2	1:A:472:TYR:HB2	1.70	0.72
1:B:461:LEU:HB2	1:B:472:TYR:HB2	1.73	0.70
1:B:299:SER:HB2	1:B:315:PHE:HA	1.74	0.69
1:B:252:ASP:HB3	1:B:255:ALA:HB2	1.74	0.68
1:B:264:TYR:HB3	1:B:275:ARG:HB2	1.79	0.64
1:A:394:ASN:OD1	1:A:394:ASN:N	2.31	0.63
1:B:423:LYS:HE2	1:B:441:PRO:HB3	1.80	0.62
1:A:226:ALA:O	1:A:459:ARG:NH2	2.32	0.62
1:B:360:ASP:OD1	1:B:365:TRP:NE1	2.34	0.60
1:A:448:HIS:O	1:A:465:ASN:ND2	2.35	0.59
1:A:252:ASP:HB3	1:A:255:ALA:HB2	1.88	0.56
1:B:269:HIS:ND1	1:B:270:ASN:OD1	2.36	0.52
1:A:304:VAL:HG11	1:A:358:MET:HA	1.91	0.51
1:B:239:VAL:HG13	1:B:284:MET:HG2	1.94	0.50
1:B:304:VAL:HG11	1:B:358:MET:HA	1.94	0.49
1:A:327:THR:HB	1:A:329:THR:HG22	1.94	0.49
1:B:408:ILE:HB	1:B:413:TYR:HE2	1.77	0.49
1:B:414:VAL:HB	1:B:426:TYR:HB3	1.96	0.48
1:B:376:ASN:HA	1:B:398:PRO:HA	1.94	0.47
1:B:334:ARG:HH12	1:B:388:GLN:HG2	1.80	0.47
1:A:360:ASP:OD1	1:A:365:TRP:NE1	2.43	0.47
1:A:371:ASN:N	1:A:371:ASN:OD1	2.41	0.47
1:A:411:THR:HG22	1:A:429:GLN:HA	1.96	0.46
1:B:423:LYS:HG2	1:B:441:PRO:HA	1.97	0.46
1:B:236:PRO:HG2	1:B:445:LYS:HD2	1.99	0.44
1:A:368:TYR:O	1:A:378:VAL:N	2.32	0.43
1:A:332:LYS:HA	1:A:332:LYS:HD2	1.82	0.43
1:B:234:SER:OG	1:B:235:ASP:N	2.52	0.42
1:A:373:ASN:O	1:A:375:GLY:N	2.52	0.42
1:A:459:ARG:NH2	1:A:476:LEU:HD12	2.35	0.41
1:B:320:ILE:HD13	1:B:357:LEU:HD21	2.02	0.41
1:B:267:GLY:HA3	4:B:1482:GOL:H11	2.02	0.41
1:A:320:ILE:HD13	1:A:357:LEU:HD21	2.01	0.41
1:B:310:ILE:HG23	1:B:323:PHE:HB3	2.04	0.40
1:B:224:LYS:O	1:B:227:CYS:HB2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	249/471 (53%)	241 (97%)	8 (3%)	0	100	100
1	B	253/471 (54%)	244 (96%)	9 (4%)	0	100	100
All	All	502/942 (53%)	485 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/416 (54%)	216 (96%)	8 (4%)	42	63
1	B	226/416 (54%)	218 (96%)	8 (4%)	43	64
All	All	450/832 (54%)	434 (96%)	16 (4%)	42	63

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	286	THR
1	A	358	MET
1	A	371	ASN
1	A	373	ASN
1	A	394	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	425	HIS
1	A	458	ASP
1	A	465	ASN
1	B	227	CYS
1	B	282	ASP
1	B	287	ASP
1	B	303	GLN
1	B	356	ASP
1	B	425	HIS
1	B	453	ASP
1	B	469	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	469	GLN
1	B	223	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	1307	1,2	14,14,15	0.21	0	15,19,21	0.28	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	1473	1,2	14,14,15	0.28	0	15,19,21	0.40	0
2	NAG	A	2307	2	14,14,15	0.24	0	15,19,21	0.35	0
2	NAG	A	2473	2	14,14,15	0.20	0	15,19,21	0.22	0
2	NAG	B	1307	1,2	14,14,15	0.35	0	15,19,21	0.22	0
2	NAG	B	1394	1,2	14,14,15	0.42	0	15,19,21	0.65	1 (6%)
2	NAG	B	2307	2	14,14,15	0.25	0	15,19,21	0.41	0
2	NAG	B	2394	2	14,14,15	0.27	0	15,19,21	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1307	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	1473	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	2307	2	-	0/6/23/26	0/1/1/1
2	NAG	A	2473	2	-	0/6/23/26	0/1/1/1
2	NAG	B	1307	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	1394	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2307	2	-	0/6/23/26	0/1/1/1
2	NAG	B	2394	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1394	NAG	C1-O5-C5	2.32	115.19	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	1394	1	14,14,15	0.24	0	15,19,21	0.32	0
3	NAG	B	1473	1	14,14,15	0.20	0	15,19,21	0.33	0
4	GOL	B	1481	-	5,5,5	0.34	0	5,5,5	0.22	0
4	GOL	B	1482	-	5,5,5	0.26	0	5,5,5	0.71	0
4	GOL	B	1483	-	5,5,5	0.33	0	5,5,5	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1394	1	-	0/6/23/26	0/1/1/1
3	NAG	B	1473	1	-	0/6/23/26	0/1/1/1
4	GOL	B	1481	-	-	0/4/4/4	0/0/0/0
4	GOL	B	1482	-	-	0/4/4/4	0/0/0/0
4	GOL	B	1483	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1482	GOL	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	253/471 (53%)	0.43	15 (5%) 26 26	44, 67, 93, 133	0
1	B	257/471 (54%)	0.62	17 (6%) 22 22	48, 67, 98, 119	0
All	All	510/942 (54%)	0.52	32 (6%) 23 24	44, 67, 96, 133	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	421	GLY	6.6
1	A	211	VAL	6.1
1	B	478	HIS	5.8
1	B	214	LEU	5.4
1	A	212	SER	4.8
1	B	418	TYR	4.6
1	A	477	PHE	4.0
1	B	338	TYR	3.8
1	A	216	GLU	3.6
1	B	257	GLU	3.6
1	B	211	VAL	3.4
1	B	258	GLY	3.4
1	A	338	TYR	3.3
1	A	213	ASN	3.3
1	B	400	ARG	3.1
1	B	477	PHE	3.1
1	A	214	LEU	3.1
1	B	216	GLU	3.1
1	A	400	ARG	2.7
1	A	281	VAL	2.6
1	B	447	SER	2.5
1	B	219	ARG	2.5
1	A	219	ARG	2.5
1	B	479	ALA	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	387	LEU	2.3
1	A	286	THR	2.3
1	A	474	VAL	2.2
1	B	370	THR	2.2
1	B	371	ASN	2.1
1	A	285	ASN	2.0
1	A	215	GLU	2.0
1	B	278	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NAG	A	1473	14/15	0.86	0.25	2.32	73,77,80,86	0
2	NAG	A	1307	14/15	0.79	0.25	0.33	67,72,77,84	0
2	NAG	A	2473	14/15	0.76	0.37	-	92,97,100,101	0
2	NAG	B	2394	14/15	0.76	0.24	-	85,88,91,92	0
2	NAG	B	1307	14/15	0.78	0.25	-	68,72,76,83	0
2	NAG	B	1394	14/15	0.77	0.22	-	70,74,78,82	0
2	NAG	B	2307	14/15	0.69	0.47	-	89,96,98,99	0
2	NAG	A	2307	14/15	0.81	0.29	-	89,91,93,94	0

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NAG	B	1473	14/15	0.82	0.36	10.30	68,74,77,77	0
3	NAG	A	1394	14/15	0.78	0.50	8.58	69,78,81,84	0
4	GOL	B	1482	6/6	0.63	0.34	6.56	95,95,99,100	0
4	GOL	B	1483	6/6	0.47	0.41	3.09	106,107,108,108	0
4	GOL	B	1481	6/6	0.87	0.20	1.58	71,72,73,73	0
5	CL	B	1484	1/1	0.81	0.16	-	91,91,91,91	0

6.5 Other polymers [i](#)

There are no such residues in this entry.