



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 11:10 PM GMT

PDB ID : 5AMQ
Title : Structure of the La Crosse Bunyavirus polymerase in complex with the 3' and 5' viral RNA
Authors : Reguera, J.; Gerlach, P.; Cusack, S.
Deposited on : 2015-03-12
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)
RNA backbone	2183	1036 (3.40-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	2263	<div> <div>70%</div> <div>26%</div> </div>
2	B	16	<div> <div>75%</div> <div>25%</div> </div>
3	C	10	<div> <div>60%</div> <div>40%</div> </div>
4	c	8	<div> <div>63%</div> <div>38%</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 14334 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA POLYMERASE L.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1666	Total	C	N	O	S	0	0	0
			13618	8719	2269	2542	88			

- Molecule 2 is a RNA chain called RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	16	Total	C	N	O	P	0	0	0
			334	151	56	112	15			

- Molecule 3 is a RNA chain called RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	10	Total	C	N	O	P	0	0	0
			216	96	39	71	10			

- Molecule 4 is a RNA chain called RNA.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	c	8	Total	C	N	O	P	0	0	0
			166	76	31	52	7			

LEU
LEU
ILE
GLU
ARG
GLU
THR
LYS
ILE
PHE
GLU
GLU
ALA
PHE
THR
PHE
SER
GLU
ASN
GLY
PHE
ILE
SER
SER
PRO
GLU
ASN
GLY
LEU
MET
GLY
CYS
LEU
GLU
ALA
VAL
VAL
SER
SER
LEU
ILE
LYS
LYS
LEU
LEU
THR
THR
GLU
TRP
ASN
TRP
SER
THR
VAL
ILE
ASP
LYS
CYS
ILE
HIS
CYS
ILE
LEU
ILE
LYS
ASN

GLY
MET
ASP
HIS
MET
TYR
HIS
SER
PHE
PHE
ASP
VAL
PRO
LYS
CYS
PHE
MET
MET
GLY
ASN
PRO
ILE
THR
ARG
SER
SER
ILE
ASN
TRP
VAL
MET
MET
PHE
ARG
GLU
PHE
ILE
ASN
SER
SER
LEU
PRO
GLY
THR
ASP
ILE
LYS
PRO
PRO
TRP
ASN
VAL
SER
MET
THR
THR
GLU
ASN
PHE
LYS
LYS
LYS
CYS
ILE
ALA
LEU
ILE
LYS
ASN

SER
LYS
PHE
THR
GLN
ARG
ASP
PHE
SER
GLU
PHE
THR
LYS
LEU
MET
LYS
LYS
GLU
GLY
GLY
ARG
SER
SER
ASN
ILE
GLU
PHE
ASP

● Molecule 2: RNA



● Molecule 3: RNA



● Molecule 4: RNA



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	103.00Å 141.10Å 165.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.17 – 3.00 49.17 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.0 (49.17-3.00) 99.1 (49.17-3.00)	Depositor EDS
R_{merge}	0.01	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.72 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.192 , 0.252 0.194 , 0.252	Depositor DCC
R_{free} test set	2407 reflections (5.22%)	DCC
Wilson B-factor (Å ²)	80.9	Xtriage
Anisotropy	0.475	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 47.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 48549 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	14334	wwPDB-VP
Average B, all atoms (Å ²)	90.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/13890	0.61	3/18721 (0.0%)
2	B	0.33	0/372	0.69	0/577
3	C	0.70	1/241 (0.4%)	0.64	0/372
4	c	0.25	0/185	0.85	2/286 (0.7%)
All	All	0.40	1/14688 (0.0%)	0.61	5/19956 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1	A	OP3-P	-9.73	1.49	1.61

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	568	GLU	N-CA-C	13.17	146.55	111.00
1	A	569	GLU	N-CA-CB	-7.70	96.74	110.60
4	c	9	G	C5'-C4'-O4'	6.69	117.12	109.10
1	A	568	GLU	CB-CA-C	-5.68	99.04	110.40
4	c	9	G	C5'-C4'-C3'	5.60	124.96	116.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	13618	0	13692	52	0
2	B	334	0	172	0	0
3	C	216	0	108	0	0
4	c	166	0	89	0	0
All	All	14334	0	14061	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 52 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1639:GLY:O	1:A:1640:ALA:O	1.78	1.00
1:A:1063:LYS:HD3	1:A:1067:GLN:OE1	1.76	0.86
1:A:84:ILE:HD11	1:A:148:LEU:HD21	1.60	0.83
1:A:1063:LYS:HD3	1:A:1067:GLN:CD	1.99	0.83
1:A:1063:LYS:CD	1:A:1067:GLN:OE1	2.33	0.75

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1646/2263 (73%)	1568 (95%)	74 (4%)	4 (0%)	52 88

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1638	GLN
1	A	548	ALA
1	A	872	ARG
1	A	1175	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	1538/2087 (74%)	1527 (99%)	11 (1%)	88 96

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	699	ASP
1	A	732	PHE
1	A	1342	SER
1	A	390	ASN
1	A	1192	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	15/16 (93%)	4 (26%)	1 (6%)
3	C	9/10 (90%)	3 (33%)	0
4	c	7/8 (87%)	2 (28%)	0
All	All	31/34 (91%)	9 (29%)	1 (3%)

5 of 9 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	7	G
2	B	9	A
2	B	10	C
2	B	14	A
3	C	6	U

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	B	10	C

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1666/2263 (73%)	-0.11	19 (1%) 82 58	53, 84, 143, 189	0
2	B	16/16 (100%)	-0.39	0 100 100	76, 83, 93, 95	0
3	C	10/10 (100%)	-0.31	0 100 100	67, 76, 81, 84	0
4	c	8/8 (100%)	-0.68	0 100 100	85, 89, 97, 100	0
All	All	1700/2297 (74%)	-0.12	19 (1%) 82 58	53, 84, 143, 189	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1035	TYR	3.4
1	A	1045	LEU	3.4
1	A	1625	LEU	2.7
1	A	1645	MET	2.7
1	A	1040	GLY	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.