



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 10, 2016 – 02:14 PM EST

PDB ID : 5AMT  
Title : Intracellular growth locus protein E  
Authors : Robb, C.S.; Nano, F.E.; Boraston, A.B.  
Deposited on : 2015-09-01  
Resolution : 1.62 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20028320  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20028320


## 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION



The reported resolution of this entry is 1.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



<b>Metric</b>	<b>Whole archive (#Entries)</b>	<b>Similar resolution (#Entries, resolution range(Å))</b>
$R_{free}$	91344	3202 (1.64-1.60)
Clashscore	102246	3500 (1.64-1.60)
Ramachandran outliers	100387	3411 (1.64-1.60)
Sidechain outliers	100360	3410 (1.64-1.60)
RSRZ outliers	91569	3207 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	125	 73%10%16%
1	B	125	 70%10%18%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	BR	A	1127	-	-	X	-
3	BR	A	1128	-	-	-	X
3	BR	A	1130	-	-	-	X
3	BR	A	1131	-	-	-	X
3	BR	B	1126	-	-	X	X
3	BR	B	1128	-	-	-	X
3	BR	B	1129	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2005 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IGLE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	105	Total	C	N	O	S	0	0	0
			855	547	138	169	1			
1	B	103	Total	C	N	O	S	0	2	0
			850	544	136	169	1			

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	EXPRESSION TAG	UNP A0Q7H2
A	2	GLY	-	EXPRESSION TAG	UNP A0Q7H2
A	3	SER	-	EXPRESSION TAG	UNP A0Q7H2
A	4	SER	-	EXPRESSION TAG	UNP A0Q7H2
A	5	HIS	-	EXPRESSION TAG	UNP A0Q7H2
A	6	HIS	-	EXPRESSION TAG	UNP A0Q7H2
A	7	HIS	-	EXPRESSION TAG	UNP A0Q7H2
A	8	HIS	-	EXPRESSION TAG	UNP A0Q7H2
A	9	HIS	-	EXPRESSION TAG	UNP A0Q7H2
A	10	HIS	-	EXPRESSION TAG	UNP A0Q7H2
A	11	SER	-	EXPRESSION TAG	UNP A0Q7H2
A	12	SER	-	EXPRESSION TAG	UNP A0Q7H2
A	13	GLY	-	EXPRESSION TAG	UNP A0Q7H2
A	14	LEU	-	EXPRESSION TAG	UNP A0Q7H2
A	15	VAL	-	EXPRESSION TAG	UNP A0Q7H2
A	16	PRO	-	EXPRESSION TAG	UNP A0Q7H2
A	17	ARG	-	EXPRESSION TAG	UNP A0Q7H2
A	18	GLY	-	EXPRESSION TAG	UNP A0Q7H2
A	19	SER	-	EXPRESSION TAG	UNP A0Q7H2
A	20	HIS	-	EXPRESSION TAG	UNP A0Q7H2
A	21	GLY	-	EXPRESSION TAG	UNP A0Q7H2
A	22	SER	-	EXPRESSION TAG	UNP A0Q7H2
A	23	HIS	-	EXPRESSION TAG	UNP A0Q7H2
A	24	MET	-	EXPRESSION TAG	UNP A0Q7H2
B	1	MET	-	EXPRESSION TAG	UNP A0Q7H2

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	2	GLY	-	EXPRESSION TAG	UNP A0Q7H2
B	3	SER	-	EXPRESSION TAG	UNP A0Q7H2
B	4	SER	-	EXPRESSION TAG	UNP A0Q7H2
B	5	HIS	-	EXPRESSION TAG	UNP A0Q7H2
B	6	HIS	-	EXPRESSION TAG	UNP A0Q7H2
B	7	HIS	-	EXPRESSION TAG	UNP A0Q7H2
B	8	HIS	-	EXPRESSION TAG	UNP A0Q7H2
B	9	HIS	-	EXPRESSION TAG	UNP A0Q7H2
B	10	HIS	-	EXPRESSION TAG	UNP A0Q7H2
B	11	SER	-	EXPRESSION TAG	UNP A0Q7H2
B	12	SER	-	EXPRESSION TAG	UNP A0Q7H2
B	13	GLY	-	EXPRESSION TAG	UNP A0Q7H2
B	14	LEU	-	EXPRESSION TAG	UNP A0Q7H2
B	15	VAL	-	EXPRESSION TAG	UNP A0Q7H2
B	16	PRO	-	EXPRESSION TAG	UNP A0Q7H2
B	17	ARG	-	EXPRESSION TAG	UNP A0Q7H2
B	18	GLY	-	EXPRESSION TAG	UNP A0Q7H2
B	19	SER	-	EXPRESSION TAG	UNP A0Q7H2
B	20	HIS	-	EXPRESSION TAG	UNP A0Q7H2
B	21	GLY	-	EXPRESSION TAG	UNP A0Q7H2
B	22	SER	-	EXPRESSION TAG	UNP A0Q7H2
B	23	HIS	-	EXPRESSION TAG	UNP A0Q7H2
B	24	MET	-	EXPRESSION TAG	UNP A0Q7H2

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0

- Molecule 3 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	5	Total Br 5 5	0	0
3	A	5	Total Br 5 5	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	150	Total O 150 150	0	0
4	B	136	Total O 136 136	0	0



- Molecule 1: IGLE



- Molecule 1: IGLE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 <sub>1</sub> 2 <sub>1</sub> 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	43.31Å 89.61Å 47.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.02 – 1.62 38.99 – 1.62	Depositor EDS
% Data completeness (in resolution range)	99.9 (39.02-1.62) 99.9 (38.99-1.62)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.95 (at 1.62Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.165 , 0.194 0.164 , 0.194	Depositor DCC
$R_{free}$ test set	1243 reflections (5.38%)	DCC
Wilson B-factor (Å <sup>2</sup> )	13.9	Xtriage
Anisotropy	0.157	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 45.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	2005	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 56.22 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.8254e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, BR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.15	2/872 (0.2%)	1.16	4/1177 (0.3%)
1	B	1.28	5/873 (0.6%)	1.17	2/1180 (0.2%)
All	All	1.21	7/1745 (0.4%)	1.16	6/2357 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	41	GLU	CD-OE1	-6.78	1.18	1.25
1	B	103	LYS	CB-CG	-6.20	1.35	1.52
1	B	113	GLU	CG-CD	5.52	1.60	1.51
1	B	87	TYR	CZ-OH	5.31	1.46	1.37
1	A	108	SER	CB-OG	-5.22	1.35	1.42

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	119	ASP	CB-CG-OD1	7.11	124.70	118.30
1	A	24	MET	CG-SD-CE	-7.03	88.96	100.20
1	A	81	ASP	CB-CG-OD1	6.02	123.72	118.30
1	B	24	MET	CG-SD-CE	-5.67	91.13	100.20
1	B	63	ASP	CB-CG-OD2	5.31	123.08	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	124	LYS	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	855	0	840	7	0
1	B	850	0	835	12	0
2	A	4	0	6	0	0
3	A	5	0	0	3	0
3	B	5	0	0	6	0
4	A	150	0	0	3	1
4	B	136	0	0	7	1
All	All	2005	0	1681	21	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 21 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:55:GLN:CB	4:A:2065:HOH:O	2.06	1.03
1:B:70:ASN:OD1	4:B:2031:HOH:O	1.93	0.86
3:B:1129:BR:BR	4:B:2112:HOH:O	2.57	0.76
1:B:113:GLU:CD	4:B:2128:HOH:O	2.25	0.75
4:A:2053:HOH:O	3:B:1126:BR:BR	2.62	0.72

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:2066:HOH:O	4:B:2008:HOH:O[1_556]	2.11	0.09

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	103/125 (82%)	101 (98%)	2 (2%)	0	100	100
1	B	103/125 (82%)	102 (99%)	1 (1%)	0	100	100
All	All	206/250 (82%)	203 (98%)	3 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	98/118 (83%)	97 (99%)	1 (1%)	82	66
1	B	99/118 (84%)	96 (97%)	3 (3%)	48	18
All	All	197/236 (84%)	193 (98%)	4 (2%)	61	35

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	MET
1	B	24	MET
1	B	45	ASP
1	B	125	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	23	HIS
1	A	118	ASN
1	B	55	GLN
1	B	96	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 10 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	EDO	A	1126	-	3,3,3	0.52	0	2,2,2	1.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	1126	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	105/125 (84%)	-0.42	0 100 100	8, 13, 25, 30	0
1	B	103/125 (82%)	-0.45	0 100 100	7, 12, 23, 34	2 (1%)
All	All	208/250 (83%)	-0.44	0 100 100	7, 13, 24, 34	2 (0%)

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	BR	B	1129	1/1	0.90	0.27	29.94	51,51,51,51	0
3	BR	B	1128	1/1	0.92	0.28	11.11	58,58,58,58	0
3	BR	A	1131	1/1	0.98	0.17	10.50	57,57,57,57	0
3	BR	A	1130	1/1	0.95	0.27	8.22	49,49,49,49	0
3	BR	A	1128	1/1	0.96	0.22	5.22	27,27,27,27	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	BR	B	1126	1/1	0.99	0.13	2.96	11,11,11,11	1
2	EDO	A	1126	4/4	0.91	0.12	1.95	24,28,29,33	0
3	BR	B	1127	1/1	0.98	0.31	-	39,39,39,39	0
3	BR	B	1130	1/1	0.94	0.29	-	49,49,49,49	0
3	BR	A	1129	1/1	0.94	0.32	-	37,37,37,37	0
3	BR	A	1127	1/1	0.99	0.13	-	14,14,14,14	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.