



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:55 PM GMT

PDB ID : 4AQU
Title : Crystal structure of I-CreI complexed with its target methylated at position plus 2 (in the b strand) in the presence of calcium
Authors : Valton, J.; Daboussi, F.; Leduc, S.; Redondo, P.; Macmaster, R.; Molina, R.; Montoya, G.; Duchateau, P.
Deposited on : 2012-04-19
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	152	<div> <div>3%</div> <div> <div></div> <div>80%</div> <div>18%</div> <div>.</div> </div> </div>
1	B	152	<div> <div>6%</div> <div> <div></div> <div>81%</div> <div>15%</div> <div>.</div> </div> </div>
2	C	24	<div> <div></div> <div> <div>50%</div> <div>42%</div> <div>8%</div> </div> </div>
3	D	24	<div> <div></div> <div> <div>54%</div> <div>38%</div> <div>8%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	B	1354	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3567 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA ENDONUCLEASE I-CREI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	152	Total	C	N	O	S	0	0	0
			1236	796	210	229	1			
1	B	152	Total	C	N	O	S	0	0	0
			1236	796	210	229	1			

- Molecule 2 is a DNA chain called 5'-D(*DTP*CP*AP*AP*AP*AP*CP*GP*TP*CP*GP*TP*GP*DAP *GP*AP*CP*AP*GP*TP*TP*TP*GP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	24	Total	C	N	O	P	1	0	0
			494	236	94	141	23			

- Molecule 3 is a DNA chain called 5'-D(*DCP*CP*AP*AP*AP*CP*TP*GP*TP*CP*TP*CP*AP*5CMP *GP*AP*CP*GP*TP*TP*TP*TP*GP*A)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	24	Total	C	N	O	P	0	0	0
			485	234	85	143	23			

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Ca	0	0
			2	2		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	52	Total	O	0	0
			52	52		
6	B	23	Total	O	0	0
			23	23		
6	C	18	Total	O	0	0
			18	18		
6	D	9	Total	O	0	0
			9	9		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA ENDONUCLEASE I-CREI



- Molecule 1: DNA ENDONUCLEASE I-CREI



- Molecule 2: 5'-D(*DTP*CP*AP*AP*AP*AP*CP*GP*TP*CP*GP*TP*GP*DAP *GP*AP*C P*AP*GP*TP*TP*TP*GP*G)-3'



- Molecule 3: 5'-D(*DCP*CP*AP*AP*AP*CP*TP*GP*TP*CP*TP*CP*AP*5CMP *GP*AP* CP*GP*TP*TP*TP*TP*GP*A)-3'



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2 ₁	Depositor
Cell constants a, b, c, α , β , γ	71.58 Å 45.35 Å 172.33 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.80 – 2.30 44.80 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (44.80-2.30) 99.7 (44.80-2.30)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.63 (at 2.29 Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.196 , 0.249 0.195 , 0.244	Depositor DCC
R_{free} test set	2000 reflections (8.43%)	DCC
Wilson B-factor (Å ²)	43.0	Xtriage
Anisotropy	0.151	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 34.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 25711 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3567	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA, 5CM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/1259	0.50	0/1700
1	B	0.39	0/1259	0.48	0/1700
2	C	0.86	1/555 (0.2%)	1.52	7/856 (0.8%)
3	D	0.83	0/519	1.48	7/796 (0.9%)
All	All	0.57	1/3592 (0.0%)	0.95	14/5052 (0.3%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	412	DT	C5-C7	6.68	1.54	1.50

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	412	DT	C6-C5-C7	-9.08	117.45	122.90
2	C	413	DG	O4'-C1'-N9	-7.88	102.48	108.00
2	C	412	DT	C4-C5-C7	7.51	123.50	119.00
3	D	521	DT	O4'-C1'-N1	-7.29	102.89	108.00
2	C	403	DA	O4'-C1'-N9	-7.08	103.05	108.00
3	D	516	DA	O4'-C1'-N9	6.92	112.84	108.00
2	C	423	DG	O4'-C1'-N9	6.35	112.44	108.00
3	D	522	DT	O4'-C1'-N1	-6.29	103.59	108.00
3	D	515	DG	O4'-C1'-N9	-6.19	103.67	108.00
3	D	511	DT	C6-C5-C7	-5.78	119.43	122.90
3	D	501	DC	O4'-C1'-N1	5.75	112.02	108.00
2	C	422	DT	O4'-C1'-N1	-5.74	103.98	108.00
2	C	416	DA	O4'-C1'-N9	5.72	112.00	108.00
3	D	524	DA	O4'-C1'-N9	5.02	111.51	108.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1236	0	1270	20	0
1	B	1236	0	1270	15	0
2	C	494	0	272	8	0
3	D	485	0	275	6	0
4	A	2	0	0	0	0
5	A	6	0	8	0	0
5	B	6	0	8	0	0
6	A	52	0	0	3	0
6	B	23	0	0	0	0
6	C	18	0	0	0	0
6	D	9	0	0	0	0
All	All	3567	0	3103	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:324:GLU:O	1:B:327:THR:HG22	1.84	0.78
1:A:123:LEU:HD21	1:A:149:ARG:HG2	1.72	0.71
1:A:73:VAL:HG21	3:D:514:5CM:H5A1	1.75	0.68
1:B:281:ILE:HG23	1:B:282:LYS:HG3	1.74	0.68
3:D:518:DG:H2''	3:D:519:DT:H5'	1.79	0.64
1:A:62:ILE:HG22	1:A:64:VAL:HG12	1.83	0.60
1:A:52:ARG:NH1	1:A:69:ASP:OD1	2.35	0.60
6:A:2003:HOH:O	3:D:515:DG:OP2	2.17	0.58
1:B:228:LYS:NZ	2:C:419:DG:O6	2.34	0.56
2:C:401:DT:H2''	2:C:402:DC:H5''	1.88	0.56
2:C:418:DA:H2''	2:C:419:DG:H5'	1.88	0.55
1:A:151:VAL:C	1:A:153:ASP:H	2.10	0.54
1:B:330:ASP:OD1	1:B:341:ARG:NH2	2.40	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:124:GLU:O	1:A:127:THR:HB	2.12	0.49
1:B:235:PHE:HE2	1:B:316:LYS:HA	1.78	0.48
1:B:344:THR:H	1:B:347:THR:HG23	1.78	0.48
1:B:312:LEU:N	1:B:313:PRO:CD	2.78	0.46
1:B:318:SER:OG	1:B:321:LYS:N	2.38	0.46
1:A:142:LYS:HG3	3:D:518:DG:H5''	1.99	0.45
2:C:418:DA:H2''	2:C:419:DG:C5'	2.46	0.45
1:B:292:GLN:HE21	1:B:292:GLN:HB3	1.63	0.45
1:A:123:LEU:HD21	1:A:149:ARG:CG	2.43	0.44
1:A:130:ASP:OD1	1:A:141:ARG:NH2	2.48	0.44
1:A:130:ASP:HA	1:A:141:ARG:NH2	2.32	0.43
6:A:2052:HOH:O	2:C:415:DG:H5'	2.18	0.43
1:B:321:LYS:HE3	1:B:324:GLU:OE1	2.19	0.43
1:B:232:SER:O	3:D:501:DC:H2'	2.19	0.43
2:C:413:DG:H1'	2:C:414:DA:O4'	2.19	0.42
1:A:82:LYS:HB2	1:A:83:PRO:HD3	2.00	0.42
2:C:402:DC:H2''	2:C:403:DA:H5'	2.01	0.42
1:A:123:LEU:HD12	1:A:123:LEU:HA	1.72	0.42
1:B:314:SER:OG	1:B:314:SER:O	2.36	0.42
1:B:302:ALA:O	1:B:306:LEU:HB2	2.19	0.42
1:A:141:ARG:NH1	6:A:2008:HOH:O	2.34	0.42
1:A:22:SER:HB3	1:A:44:GLN:HG2	2.02	0.42
1:B:228:LYS:HA	1:B:229:PRO:HD2	1.83	0.41
1:A:33:TYR:OH	2:C:403:DA:N7	2.51	0.41
1:A:54:PHE:CD2	1:B:297:LEU:HD21	2.56	0.41
1:A:28:LYS:O	1:A:37:HIS:HA	2.21	0.40
1:A:151:VAL:C	1:A:153:ASP:N	2.74	0.40
1:A:127:THR:O	1:A:130:ASP:HB2	2.22	0.40
1:A:88:LEU:HD23	1:A:88:LEU:HA	1.95	0.40
3:D:513:DA:H1'	3:D:514:5CM:O4'	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	150/152 (99%)	143 (95%)	6 (4%)	1 (1%)	26	31
1	B	150/152 (99%)	137 (91%)	13 (9%)	0	100	100
All	All	300/304 (99%)	280 (93%)	19 (6%)	1 (0%)	46	57

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	152	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	139/139 (100%)	128 (92%)	11 (8%)	15	19
1	B	139/139 (100%)	126 (91%)	13 (9%)	11	12
All	All	278/278 (100%)	254 (91%)	24 (9%)	13	15

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	THR
1	A	41	LEU
1	A	48	LYS
1	A	68	ARG
1	A	81	ILE
1	A	118	SER
1	A	121	LYS
1	A	123	LEU
1	A	124	GLU
1	A	127	THR
1	A	141	ARG
1	B	203	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	204	LYS
1	B	257	LYS
1	B	268	ARG
1	B	269	ASP
1	B	292	GLN
1	B	307	LYS
1	B	314	SER
1	B	316	LYS
1	B	317	GLU
1	B	318	SER
1	B	344	THR
1	B	347	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	5CM	D	514	3,2,4	13,21,22	3.05	6 (46%)	17,30,33	1.72	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	5CM	D	514	3,2,4	-	0/3/21/22	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	514	5CM	C3'-C4'	-2.26	1.46	1.53
3	D	514	5CM	C6-C5	3.24	1.48	1.40
3	D	514	5CM	C4-N4	4.09	1.44	1.34
3	D	514	5CM	C5-C4	4.18	1.47	1.41
3	D	514	5CM	C4-N3	4.58	1.42	1.35
3	D	514	5CM	C6-N1	6.63	1.44	1.35

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	514	5CM	O5'-C5'-C4'	2.34	117.68	109.12
3	D	514	5CM	O4'-C1'-N1	3.72	114.16	107.72
3	D	514	5CM	C2'-C1'-N1	4.29	124.58	114.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	514	5CM	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GOL	A	1155	-	5,5,5	0.36	0	5,5,5	0.32	0
5	GOL	B	1354	-	5,5,5	0.34	0	5,5,5	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	1155	-	-	0/4/4/4	0/0/0/0
5	GOL	B	1354	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	152/152 (100%)	0.23	4 (2%) 59 68	19, 36, 65, 84	0
1	B	152/152 (100%)	0.39	9 (5%) 26 34	21, 44, 79, 99	0
2	C	24/24 (100%)	-0.15	0 100 100	27, 36, 51, 59	0
3	D	23/24 (95%)	-0.16	0 100 100	25, 37, 50, 54	0
All	All	351/352 (99%)	0.25	13 (3%) 45 54	19, 39, 75, 99	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	152	LEU	4.1
1	B	351	VAL	4.1
1	B	316	LYS	3.7
1	B	317	GLU	3.1
1	A	118	SER	3.0
1	B	319	PRO	2.8
1	A	117	GLU	2.8
1	B	320	ASP	2.7
1	B	323	LEU	2.6
1	B	326	CYS	2.4
1	B	327	THR	2.1
1	A	122	PHE	2.0
1	B	318	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy

less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	5CM	D	514	20/21	0.95	0.18	-	19,30,38,47	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	GOL	B	1354	6/6	0.85	0.19	2.60	48,53,56,56	0
5	GOL	A	1155	6/6	0.91	0.19	1.39	44,52,55,56	0
4	CA	A	1156	1/1	0.96	0.14	-0.68	25,25,25,25	0
4	CA	A	1154	1/1	0.97	0.12	-2.26	22,22,22,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.