



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 12:34 AM GMT

PDB ID : 2ASV
Title : X-Ray studies on protein complexes: Enzymatic catalysis in Crystals of E. coli
Maltodextrin Phosphorylase (MalP)
Authors : Geremia, S.; Campagnolo, M.
Deposited on : 2005-08-24
Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1833 (1.96-1.96)
Clashscore	102246	1953 (1.96-1.96)
Ramachandran outliers	100387	1936 (1.96-1.96)
Sidechain outliers	100360	1936 (1.96-1.96)
RSRZ outliers	91569	1835 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	796	<div> <div>4%</div> <div>86%</div> <div>13%</div> </div>
1	B	796	<div> <div>6%</div> <div>86%</div> <div>13%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLC	B	996	-	-	-	X
3	ASO	A	1998	-	-	-	X
3	ASO	B	2998	-	-	X	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 14136 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltodextrin phosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	796	Total	C	N	O	S	0	0	0
			6389	4079	1128	1162	20			
1	B	796	Total	C	N	O	S	0	0	0
			6389	4079	1128	1162	20			


There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	261	ALA	HIS	ENGINEERED	UNP P00490
A	262	PHE	THR	ENGINEERED	UNP P00490
A	263	GLU	ALA	ENGINEERED	UNP P00490
B	261	ALA	HIS	ENGINEERED	UNP P00490
B	262	PHE	THR	ENGINEERED	UNP P00490
B	263	GLU	ALA	ENGINEERED	UNP P00490

- Molecule 2 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	5	Total	C	O	0	0
			56	30	26		
2	B	5	Total	C	O	0	0
			56	30	26		

- Molecule 3 is SUGAR (D-GLUCOPYRANOSYLIUM) (three-letter code: ASO) (formula: C₆H₁₂O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	6	5		
3	B	1	Total	C	O	0	0
			11	6	5		

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 5 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: $C_8H_{10}NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
5	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	613	Total	O	0	0
			613	613		
6	B	571	Total	O	0	0
			571	571		

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Chain A:

4% 86% 13%

Chain B:

Category	Status
S1	Green
Q2	Green
P3	Red
I4	Green
D7	Orange
Q11	Yellow
S15	Yellow
W18	Yellow
Q19	Green
R20	Yellow
R47	Yellow
A48	Green
Q49	Yellow
P50	Yellow
F51	Red
A56	Red
N57	Red
E67	Yellow
Q85	Red
D89	Yellow
K92	Red
A93	Green
Y94	Yellow
D95	Red
I96	Green
N97	Yellow
D100	Yellow
P108	Yellow
A109	Yellow
L110	Green
G111	Yellow
N112	Yellow
L138	Yellow
N139	Yellow
G143	Yellow
D151	Red
D159	Yellow
D160	Green
W161	Yellow
F169	Yellow
N172	Yellow
D176	Yellow
T186	Red
K187	Red
D188	Red
G189	Red
R190	Red
W202	Yellow
Q215	Yellow
W220	Yellow
D229	Yellow
D237	Orange
F238	Green
L239	Yellow
R240	Yellow
D259	Yellow
M260	Yellow
Q275	Yellow
C278	Yellow
R286	Yellow
L289	Red
K293	Red
D299	Yellow
D307	Yellow
T308	Green
H309	Yellow
P310	Yellow
Q326	Yellow
W333	Yellow
N344	Yellow
H345	Yellow
L363	Yellow
K384	Red
K385	Yellow
F387	Yellow
G388	Yellow
D389	Yellow
E390	Yellow
K391	Yellow
D400	Orange
D401	Red
H424	Yellow
S425	Green
D426	Yellow
E435	Yellow
K443	Yellow
V447	Yellow
R454	Yellow
Q459	Yellow
L468	Yellow
L486	Yellow
E487	Yellow
K488	Yellow
F489	Yellow
A490	Yellow
D491	Red
D492	Yellow
A493	Green
K494	Red
F495	Yellow
R496	Yellow
Q497	Red
E500	Yellow
K515	Yellow
V516	Yellow
Q531	Yellow
I532	Yellow
K533	Yellow
R534	Orange
L535	Yellow
Y538	Yellow
Q541	Yellow
N544	Orange
I555	Orange
N558	Yellow
P559	Yellow
Q560	Yellow
A561	Yellow
D562	Yellow
R563	Yellow
V564	Yellow
Y572	Yellow
K732	Red
Y579	Red
K590	Yellow
D593	Yellow
V594	Red
D598	Yellow
P599	Red
L600	Yellow
V601	Yellow
F609	Yellow
L610	Yellow
L621	Yellow
P622	Yellow
P623	Yellow
D626	Yellow
T653	Yellow
V662	Yellow
E670	Red
I673	Yellow
E681	Red
K684	Yellow
A685	Yellow
I686	Yellow
L687	Yellow
A688	Yellow
K689	Yellow
G690	Yellow
Y691	Yellow
D692	Yellow
P693	Yellow
R694	Yellow
K695	Red
K698	Yellow
K699	Yellow
D700	Yellow
D717	Yellow
G718	Yellow
D719	Yellow
K720	Yellow
H721	Yellow
D724	Yellow
L727	Yellow

6%

86%

13%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	74.33Å 104.72Å 214.79Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 1.95 15.00 – 1.95	Depositor EDS
% Data completeness (in resolution range)	97.5 (15.00-1.95) 97.6 (15.00-1.95)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.42 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.179 , 0.226 0.191 , 0.187	Depositor DCC
R_{free} test set	6000 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	25.8	Xtriage
Anisotropy	0.346	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 55.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtriage
Outliers	0 of 119786 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	14136	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, PO4, BGC, ASO, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.90	2/6539 (0.0%)	0.91	16/8865 (0.2%)
1	B	0.90	1/6539 (0.0%)	0.92	25/8865 (0.3%)
All	All	0.90	3/13078 (0.0%)	0.92	41/17730 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	273	TYR	CE2-CZ	-5.94	1.30	1.38
1	A	578	TYR	CD2-CE2	-5.72	1.30	1.39
1	B	333	TRP	CB-CG	-5.14	1.41	1.50

The worst 5 of 41 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	299	ASP	CB-CG-OD2	7.45	125.00	118.30
1	B	307	ASP	CB-CG-OD2	7.20	124.78	118.30
1	B	626	ASP	CB-CG-OD2	6.74	124.37	118.30
1	A	259	ASP	CB-CG-OD2	6.49	124.14	118.30
1	B	500	ARG	NE-CZ-NH1	6.43	123.52	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6389	0	6333	52	0
1	B	6389	0	6333	43	0
2	A	56	0	48	2	0
2	B	56	0	48	3	0
3	A	11	0	11	3	0
3	B	11	0	12	6	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0
5	A	15	0	6	0	0
5	B	15	0	6	0	0
6	A	613	0	0	8	0
6	B	571	0	0	6	0
All	All	14136	0	12797	97	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 97 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:997:GLC:O4	3:A:1998:ASO:H11	1.47	1.15
2:B:997:GLC:O4	3:B:2998:ASO:H11	1.65	0.95
1:B:345:HIS:O	3:B:2998:ASO:H12	1.74	0.86
1:A:97:ASN:HD22	1:A:100:ASP:H	1.33	0.76
2:A:997:GLC:HO4	3:A:1998:ASO:H11	1.49	0.74

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	794/796 (100%)	770 (97%)	23 (3%)	1 (0%)	56 48

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	794/796 (100%)	770 (97%)	23 (3%)	1 (0%)	56	48
All	All	1588/1592 (100%)	1540 (97%)	46 (3%)	2 (0%)	56	48

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	533	LYS
1	B	533	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	667/667 (100%)	633 (95%)	34 (5%)	29	13
1	B	667/667 (100%)	635 (95%)	32 (5%)	31	15
All	All	1334/1334 (100%)	1268 (95%)	66 (5%)	31	15

5 of 66 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	727	LEU
1	B	139	ASN
1	B	689	LYS
1	A	732	LYS
1	A	795	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	544	ASN
1	B	57	ASN
1	B	531	GLN
1	A	678	HIS
1	B	9	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

10 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BGC	A	993	2	12,12,12	0.54	0	17,17,17	1.06	0
2	GLC	A	994	2	11,11,12	0.94	0	14,15,17	1.48	3 (21%)
2	GLC	A	995	2	11,11,12	0.89	0	14,15,17	1.21	1 (7%)
2	GLC	A	996	2	11,11,12	0.91	0	14,15,17	0.83	0
2	GLC	A	997	2	11,11,12	0.94	0	14,15,17	1.49	2 (14%)
2	BGC	B	993	2	12,12,12	0.63	0	17,17,17	1.03	1 (5%)
2	GLC	B	994	2	11,11,12	0.51	0	14,15,17	1.29	4 (28%)
2	GLC	B	995	2	11,11,12	0.82	1 (9%)	14,15,17	1.57	2 (14%)
2	GLC	B	996	2	11,11,12	0.75	0	14,15,17	1.40	3 (21%)
2	GLC	B	997	2	11,11,12	0.83	0	14,15,17	1.82	5 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	A	993	2	-	0/2/22/22	0/1/1/1
2	GLC	A	994	2	-	0/2/19/22	0/1/1/1
2	GLC	A	995	2	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	996	2	-	0/2/19/22	0/1/1/1
2	GLC	A	997	2	-	0/2/19/22	0/1/1/1
2	BGC	B	993	2	-	0/2/22/22	0/1/1/1
2	GLC	B	994	2	-	0/2/19/22	0/1/1/1
2	GLC	B	995	2	-	0/2/19/22	0/1/1/1
2	GLC	B	996	2	-	0/2/19/22	0/1/1/1
2	GLC	B	997	2	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	995	GLC	C2-C3	2.02	1.55	1.52

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	997	GLC	O3-C3-C2	-3.45	103.77	110.00
2	A	994	GLC	O5-C1-C2	-2.85	106.24	110.86
2	B	996	GLC	O2-C2-C3	-2.52	105.05	110.12
2	B	996	GLC	O3-C3-C2	-2.43	105.60	110.00
2	B	997	GLC	O4-C4-C5	-2.28	103.21	109.24

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	997	GLC	2	0
2	B	997	GLC	3	0

5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is

the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ASO	A	1998	-	11,11,11	3.40	3 (27%)	14,15,15	4.27	7 (50%)
5	PLP	A	900	1	15,15,16	2.03	3 (20%)	21,22,23	1.66	6 (28%)
4	PO4	A	999	-	4,4,4	0.60	0	6,6,6	0.29	0
4	PO4	B	1999	-	4,4,4	0.56	0	6,6,6	0.30	0
3	ASO	B	2998	-	11,11,11	3.28	4 (36%)	14,15,15	3.84	5 (35%)
5	PLP	B	900	1	15,15,16	2.14	3 (20%)	21,22,23	1.82	6 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ASO	A	1998	-	-	0/2/19/19	0/1/1/1
5	PLP	A	900	1	-	0/6/6/8	0/1/1/1
4	PO4	A	999	-	-	0/0/0/0	0/0/0/0
4	PO4	B	1999	-	-	0/0/0/0	0/0/0/0
3	ASO	B	2998	-	-	0/2/19/19	0/1/1/1
5	PLP	B	900	1	-	0/6/6/8	0/1/1/1

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1998	ASO	O5-C1	-9.94	1.27	1.43
3	B	2998	ASO	O5-C1	-9.48	1.27	1.43
5	B	900	PLP	C3-C2	-6.92	1.36	1.40
5	A	900	PLP	C3-C2	-5.71	1.36	1.40
3	A	1998	ASO	O2-C2	-3.33	1.35	1.43

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	900	PLP	C3-C2-N1	-3.51	115.76	120.61
5	B	900	PLP	C4A-C4-C3	-3.09	114.76	120.36
5	B	900	PLP	C6-C5-C4	-3.03	115.58	118.15
5	B	900	PLP	C3-C2-N1	-2.85	116.68	120.61
5	A	900	PLP	C4A-C4-C3	-2.82	115.25	120.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1998	ASO	3	0
3	B	2998	ASO	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	796/796 (100%)	0.35	35 (4%) 38 49	20, 31, 50, 67	0
1	B	796/796 (100%)	0.35	49 (6%) 24 34	20, 31, 49, 67	0
All	All	1592/1592 (100%)	0.35	84 (5%) 30 41	20, 31, 49, 67	0

The worst 5 of 84 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	187	LYS	8.8
1	B	57	ASN	7.7
1	A	51	PHE	6.9
1	B	559	PRO	6.8
1	B	562	ASP	5.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	GLC	B	996	11/12	0.91	0.14	4.00	31,32,36,36	0
2	GLC	A	997	11/12	0.93	0.13	0.93	27,28,31,35	0
2	GLC	B	995	11/12	0.92	0.14	0.70	30,34,37,39	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	GLC	A	995	11/12	0.91	0.11	0.27	30,32,34,35	0
2	GLC	B	997	11/12	0.95	0.10	0.01	26,30,31,34	0
2	GLC	B	994	11/12	0.93	0.12	-0.46	38,40,43,44	0
2	GLC	A	996	11/12	0.96	0.09	-0.66	28,30,32,35	0
2	GLC	A	994	11/12	0.96	0.09	-0.81	34,37,41,42	0
2	BGC	B	993	12/12	0.60	0.42	-	49,53,54,55	10
2	BGC	A	993	12/12	0.61	0.41	-	49,53,54,54	10

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	ASO	B	2998	11/11	0.86	0.20	4.16	34,41,45,47	0
3	ASO	A	1998	11/11	0.89	0.17	2.44	29,39,42,43	0
5	PLP	A	900	15/16	0.96	0.10	-0.12	21,24,32,32	0
5	PLP	B	900	15/16	0.96	0.09	-0.40	21,24,31,31	0
4	PO4	A	999	5/5	0.98	0.07	-1.24	31,33,38,38	0
4	PO4	B	1999	5/5	0.97	0.08	-1.80	30,33,36,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.