



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:27 AM GMT

PDB ID : 3AUJ
Title : Structure of diol dehydratase complexed with glycerol
Authors : Yamanishi, M.; Kinoshita, K.; Fukuoka, M.; Shibata, T.; Tobimatsu, T.; Toraya, T.
Deposited on : 2011-02-07
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	554	<div> <div>89%</div> <div>10%</div> <div>.</div> </div>
1	L	554	<div> <div>84%</div> <div>14%</div> <div>..</div> </div>
2	B	224	<div> <div>68%</div> <div>12%</div> <div>20%</div> </div>
2	E	224	<div> <div>71%</div> <div>9%</div> <div>20%</div> </div>
3	G	173	<div> <div>68%</div> <div>13%</div> <div>20%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	M	173	 A horizontal bar chart showing the quality of chain M. The bar is divided into three segments: a small red segment at the beginning labeled '2%', a large green segment labeled '68%', and a yellow segment labeled '11%'. A dot is located on the yellow segment, and a grey segment at the end is labeled '20%'.

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 14964 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Diol dehydrase alpha subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	552	Total	C	N	O	S	0	0	0
			4210	2625	728	828	29			
1	L	551	Total	C	N	O	S	0	0	0
			4201	2620	727	825	29			

- Molecule 2 is a protein called Diol dehydrase beta subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	179	Total	C	N	O	S	0	0	0
			1367	865	245	255	2			
2	E	180	Total	C	N	O	S	0	0	0
			1375	869	246	258	2			

- Molecule 3 is a protein called Diol dehydrase gamma subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	139	Total	C	N	O	S	0	0	0
			1110	690	200	217	3			
3	M	138	Total	C	N	O	S	0	0	0
			1104	687	199	215	3			

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	L	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Ca	0	0
			1	1		
5	L	1	Total	Ca	0	0
			1	1		

- Molecule 6 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	P	0	0
			5	4	1		
6	B	1	Total	O	P	0	0
			5	4	1		
6	G	1	Total	O	P	0	0
			5	4	1		
6	L	1	Total	O	P	0	0
			5	4	1		
6	E	1	Total	O	P	0	0
			5	4	1		
6	M	1	Total	O	P	0	0
			5	4	1		

- Molecule 7 is COBALAMIN (three-letter code: B12) (formula: $C_{62}H_{89}CoN_{13}O_{14}P$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
7	B	1	Total	C	Co	N	O	P	0	0
			91	62	1	13	14	1		
7	E	1	Total	C	Co	N	O	P	0	0
			91	62	1	13	14	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	417	Total	O	0	0
			417	417		
8	B	158	Total	O	0	0
			158	158		
8	G	116	Total	O	0	0
			116	116		
8	L	393	Total	O	0	0
			393	393		
8	E	178	Total	O	0	0
			178	178		
8	M	109	Total	O	0	0
			109	109		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Diol dehydrase alpha subunit



• Molecule 1: Diol dehydrase alpha subunit



• Molecule 2: Diol dehydrase beta subunit




• Molecule 2: Diol dehydrase beta subunit






• Molecule 3: Diol dehydrase gamma subunit



• Molecule 3: Diol dehydrase gamma subunit



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	190.61Å 196.80Å 77.36Å 90.00° 95.03° 90.00°	Depositor
Resolution (Å)	40.34 – 2.10 40.34 – 2.10	Depositor EDS
% Data completeness (in resolution range)	82.8 (40.34-2.10) 81.8 (40.34-2.10)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.81 (at 2.10Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.186 , 0.191 0.178 , 0.202	Depositor DCC
R_{free} test set	15624 reflections (12.85%)	DCC
Wilson B-factor (Å ²)	18.7	Xtriage
Anisotropy	0.724	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 48.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 156848 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	14964	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA, B12, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/4282	0.60	0/5799
1	L	0.31	0/4273	0.60	1/5787 (0.0%)
2	B	0.30	0/1389	0.58	0/1879
2	E	0.30	0/1397	0.58	0/1890
3	G	0.29	0/1125	0.52	0/1519
3	M	0.29	0/1119	0.52	0/1511
All	All	0.31	0/13585	0.58	1/18385 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	L	269	GLY	N-CA-C	5.22	126.14	113.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4210	0	4146	38	0
1	L	4201	0	4140	58	0
2	B	1367	0	1419	18	0
2	E	1375	0	1423	15	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	G	1110	0	1119	13	0
3	M	1104	0	1114	16	0
4	A	6	0	6	0	0
4	L	6	0	6	0	0
5	A	1	0	0	0	0
5	L	1	0	0	0	0
6	A	5	0	0	0	0
6	B	5	0	0	0	0
6	E	5	0	0	0	0
6	G	5	0	0	0	0
6	L	5	0	0	0	0
6	M	5	0	0	0	0
7	B	91	0	88	6	0
7	E	91	0	88	5	0
8	A	417	0	0	4	0
8	B	158	0	0	3	0
8	E	178	0	0	1	0
8	G	116	0	0	0	0
8	L	393	0	0	4	0
8	M	109	0	0	1	0
All	All	14964	0	13549	162	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 162 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:2:ARG:HB3	1:L:2:ARG:HH11	1.19	1.04
1:A:24:ILE:H	1:A:24:ILE:HD12	1.36	0.90
3:G:71:SER:OG	3:G:73:LYS:HG3	1.83	0.79
3:G:36:ARG:HD2	3:G:88:LEU:HD21	1.66	0.77
1:L:489:MET:O	1:L:492:ILE:HG22	1.87	0.74

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	550/554 (99%)	526 (96%)	22 (4%)	2 (0%)	39	37
1	L	549/554 (99%)	516 (94%)	31 (6%)	2 (0%)	39	37
2	B	177/224 (79%)	173 (98%)	4 (2%)	0	100	100
2	E	178/224 (80%)	175 (98%)	3 (2%)	0	100	100
3	G	137/173 (79%)	135 (98%)	2 (2%)	0	100	100
3	M	136/173 (79%)	132 (97%)	4 (3%)	0	100	100
All	All	1727/1902 (91%)	1657 (96%)	66 (4%)	4 (0%)	52	53

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	300	VAL
1	A	363	GLY
1	L	300	VAL
1	L	363	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	451/453 (100%)	443 (98%)	8 (2%)	66	72
1	L	450/453 (99%)	439 (98%)	11 (2%)	57	61
2	B	147/183 (80%)	146 (99%)	1 (1%)	88	92

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	E	148/183 (81%)	148 (100%)	0	100	100
3	G	118/141 (84%)	116 (98%)	2 (2%)	68	74
3	M	117/141 (83%)	115 (98%)	2 (2%)	68	74
All	All	1431/1554 (92%)	1407 (98%)	24 (2%)	68	74

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	G	131	SER
1	L	25	GLU
3	M	58	ASN
1	L	1	MET
1	L	2	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	L	35	ASN
1	L	182	ASN
2	E	167	GLN
3	G	58	ASN
2	E	204	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 12 ligands modelled in this entry, 2 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
6	PO4	A	1401	-	4,4,4	1.13	0	6,6,6	0.27	0
4	GOL	A	1602	5	5,5,5	0.15	0	5,5,5	0.13	0
6	PO4	B	1406	-	4,4,4	1.07	0	6,6,6	0.27	0
7	B12	B	1601	-	74,101,101	1.34	9 (12%)	111,166,166	1.13	9 (8%)
6	PO4	E	1405	-	4,4,4	1.04	0	6,6,6	0.27	0
7	B12	E	2601	-	74,101,101	1.31	9 (12%)	111,166,166	1.16	12 (10%)
6	PO4	G	1403	-	4,4,4	1.13	0	6,6,6	0.27	0
6	PO4	L	1402	-	4,4,4	1.14	0	6,6,6	0.27	0
4	GOL	L	2602	5	5,5,5	0.14	0	5,5,5	0.13	0
6	PO4	M	1404	-	4,4,4	1.13	0	6,6,6	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PO4	A	1401	-	-	0/0/0/0	0/0/0/0
4	GOL	A	1602	5	-	0/4/4/4	0/0/0/0
6	PO4	B	1406	-	-	0/0/0/0	0/0/0/0
7	B12	B	1601	-	-	0/51/223/223	0/3/11/11
6	PO4	E	1405	-	-	0/0/0/0	0/0/0/0
7	B12	E	2601	-	-	0/51/223/223	0/3/11/11
6	PO4	G	1403	-	-	0/0/0/0	0/0/0/0
6	PO4	L	1402	-	-	0/0/0/0	0/0/0/0
4	GOL	L	2602	5	-	0/4/4/4	0/0/0/0
6	PO4	M	1404	-	-	0/0/0/0	0/0/0/0

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	E	2601	B12	C5M-C5B	-5.50	1.39	1.51
7	B	1601	B12	C5M-C5B	-5.46	1.40	1.51
7	B	1601	B12	C11-C10	-3.52	1.34	1.41
7	E	2601	B12	C11-C10	-3.28	1.35	1.41
7	E	2601	B12	C1-C2	2.27	1.64	1.58

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	B	1601	B12	O58-C57-C56	-2.71	117.31	121.98
7	E	2601	B12	O58-C57-C56	-2.61	117.49	121.98
7	B	1601	B12	C4B-C9B-C8B	-2.41	118.64	121.10
7	E	2601	B12	C4B-C9B-C8B	-2.21	118.83	121.10
7	B	1601	B12	C3-C4-C5	-2.21	124.40	131.88

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	B	1601	B12	6	0
7	E	2601	B12	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	552/554 (99%)	-0.30	5 (0%) 85 88	17, 24, 36, 83	0
1	L	551/554 (99%)	-0.22	8 (1%) 76 81	17, 24, 37, 69	0
2	B	179/224 (79%)	-0.45	2 (1%) 82 86	18, 27, 38, 63	0
2	E	180/224 (80%)	-0.48	3 (1%) 73 78	20, 26, 39, 84	0
3	G	139/173 (80%)	-0.15	3 (2%) 65 71	21, 31, 47, 85	0
3	M	138/173 (79%)	-0.05	3 (2%) 65 71	23, 33, 55, 83	0
All	All	1739/1902 (91%)	-0.27	24 (1%) 78 82	17, 26, 42, 85	0

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	G	35	SER	8.2
2	B	224	LEU	6.2
2	E	45	ASP	4.8
1	A	1	MET	4.1
1	A	552	GLU	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	GOL	L	2602	6/6	0.95	0.21	0.68	20,24,28,34	0
7	B12	B	1601	91/91	0.97	0.14	0.48	16,20,24,25	0
7	B12	E	2601	91/91	0.97	0.14	0.39	15,20,24,26	0
6	PO4	G	1403	5/5	0.97	0.12	-0.07	61,61,61,62	0
4	GOL	A	1602	6/6	0.97	0.17	-0.17	15,20,26,33	0
5	CA	A	1603	1/1	1.00	0.10	-2.83	20,20,20,20	0
5	CA	L	2603	1/1	0.99	0.11	-3.21	21,21,21,21	0
6	PO4	L	1402	5/5	0.95	0.23	-	78,78,78,78	0
6	PO4	A	1401	5/5	0.95	0.14	-	74,74,75,75	0
6	PO4	B	1406	5/5	0.93	0.31	-	70,71,72,72	0
6	PO4	E	1405	5/5	0.91	0.30	-	70,71,72,72	0
6	PO4	M	1404	5/5	0.94	0.13	-	61,61,62,63	0

6.5 Other polymers [i](#)

There are no such residues in this entry.