



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 06:28 PM GMT

PDB ID : 1AWV
Title : CYPA COMPLEXED WITH HVGPIA
Authors : Vajdos, F.F.
Deposited on : 1997-10-05
Resolution : 2.34 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1406 (2.36-2.32)
Clashscore	102246	1509 (2.36-2.32)
Ramachandran outliers	100387	1490 (2.36-2.32)
Sidechain outliers	100360	1491 (2.36-2.32)
RSRZ outliers	91569	1412 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	164	<div> <div>8%</div> <div>71%</div> <div>27%</div> <div>.</div> </div>
1	B	164	<div> <div>54%</div> <div>50%</div> <div>42%</div> <div>8%</div> </div>
1	C	164	<div> <div>18%</div> <div>62%</div> <div>35%</div> <div>.</div> </div>
1	D	164	<div> <div>21%</div> <div>58%</div> <div>39%</div> <div>.</div> </div>
1	E	164	<div> <div>15%</div> <div>66%</div> <div>30%</div> <div>.</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	164	<div><div></div><div>51%</div><div>57%</div><div>40%</div><div></div></div>
2	G	6	<div><div></div><div>67%</div><div>33%</div><div></div></div>
2	H	6	<div><div></div><div>17%</div><div>67%</div><div>33%</div><div></div></div>
2	I	6	<div><div></div><div>50%</div><div>50%</div><div></div></div>
2	J	6	<div><div></div><div>17%</div><div>33%</div><div>67%</div><div></div></div>
2	K	6	<div><div></div><div>17%</div><div>67%</div><div>33%</div><div></div></div>
2	L	6	<div><div></div><div>33%</div><div>83%</div><div>17%</div><div></div></div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7975 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYCLOPHILIN A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	164	Total	C	N	O	S	0	0	0
			1258	797	217	236	8			
1	B	164	Total	C	N	O	S	0	0	0
			1258	797	217	236	8			
1	C	164	Total	C	N	O	S	0	0	0
			1258	797	217	236	8			
1	D	164	Total	C	N	O	S	0	0	0
			1258	797	217	236	8			
1	E	164	Total	C	N	O	S	0	0	0
			1258	797	217	236	8			
1	F	164	Total	C	N	O	S	0	0	0
			1258	797	217	236	8			

- Molecule 2 is a protein called PEPTIDE FROM THE HIV-1 CAPSID PROTEIN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	G	6	Total	C	N	O	11	0	0
			42	27	8	7			
2	H	6	Total	C	N	O	0	0	0
			42	27	8	7			
2	I	6	Total	C	N	O	0	0	0
			42	27	8	7			
2	J	6	Total	C	N	O	0	0	0
			42	27	8	7			
2	K	6	Total	C	N	O	11	0	0
			42	27	8	7			
2	L	6	Total	C	N	O	0	0	0
			42	27	8	7			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	32	Total 32	O 32	0	0
3	B	22	Total 22	O 22	0	0
3	C	31	Total 31	O 31	0	0
3	D	20	Total 20	O 20	0	0
3	E	28	Total 28	O 28	0	0
3	F	29	Total 29	O 29	0	0
3	G	3	Total 3	O 3	0	0
3	H	1	Total 1	O 1	0	0
3	I	2	Total 2	O 2	0	0
3	J	2	Total 2	O 2	0	0
3	K	3	Total 3	O 3	0	0
3	L	2	Total 2	O 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: CYCLOPHILIN A




• Molecule 1: CYCLOPHILIN A




• Molecule 1: CYCLOPHILIN A




• Molecule 1: CYCLOPHILIN A




• Molecule 1: CYCLOPHILIN A




• Molecule 1: CYCLOPHILIN A



• Molecule 2: PEPTIDE FROM THE HIV-1 CAPSID PROTEIN



• Molecule 2: PEPTIDE FROM THE HIV-1 CAPSID PROTEIN




- Molecule 2: PEPTIDE FROM THE HIV-1 CAPSID PROTEIN

Chain I: 



- Molecule 2: PEPTIDE FROM THE HIV-1 CAPSID PROTEIN

Chain J: 




- Molecule 2: PEPTIDE FROM THE HIV-1 CAPSID PROTEIN

Chain K: 



- Molecule 2: PEPTIDE FROM THE HIV-1 CAPSID PROTEIN

Chain L: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	74.00 Å 74.00 Å 190.40 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.34 14.98 – 2.35	Depositor EDS
% Data completeness (in resolution range)	83.1 (15.00-2.34) 84.1 (14.98-2.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.04 (at 2.34 Å)	Xtriage
Refinement program	X-PLOR 3.843	Depositor
R, R_{free}	0.358 , 0.460 0.355 , 0.449	Depositor DCC
R_{free} test set	1781 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	16.6	Xtriage
Anisotropy	0.687	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 63.0	EDS
Estimated twinning fraction	0.048 for h,-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	4 of 41416 reflections (0.010%)	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	7975	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 85.32 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.3041e-07. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/1286	0.69	0/1723
1	B	0.51	0/1286	0.71	0/1723
1	C	0.50	1/1286 (0.1%)	0.68	0/1723
1	D	0.48	0/1286	0.67	0/1723
1	E	0.51	0/1286	0.70	0/1723
1	F	0.49	0/1286	0.71	0/1723
2	G	0.45	0/43	0.80	0/57
2	H	0.50	0/43	0.58	0/57
2	I	0.61	0/43	0.92	0/57
2	J	0.67	0/43	0.86	0/57
2	K	0.60	0/43	0.78	0/57
2	L	0.53	0/43	0.76	0/57
All	All	0.50	1/7974 (0.0%)	0.70	0/10680

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	1115	CYS	CB-SG	-5.18	1.73	1.81

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1258	0	1225	28	0
1	B	1258	0	1225	65	0
1	C	1258	0	1225	42	0
1	D	1258	0	1225	48	0
1	E	1258	0	1225	39	0
1	F	1258	0	1225	51	0
2	G	42	0	41	1	0
2	H	42	0	41	1	0
2	I	42	0	41	2	0
2	J	42	0	41	3	0
2	K	42	0	41	2	0
2	L	42	0	41	0	0
3	A	32	0	0	1	0
3	B	22	0	0	1	0
3	C	31	0	0	1	0
3	D	20	0	0	2	0
3	E	28	0	0	1	0
3	F	29	0	0	6	0
3	G	3	0	0	0	0
3	H	1	0	0	0	0
3	I	2	0	0	0	0
3	J	2	0	0	1	0
3	K	3	0	0	0	0
3	L	2	0	0	0	0
All	All	7975	0	7596	278	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 278 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1028:LYS:HD3	1:B:1090:LEU:HD21	1.63	0.80
1:F:1067:PHE:HB3	3:F:7016:HOH:O	1.84	0.78
1:D:1095:PRO:HG3	1:D:1117:ALA:HA	1.67	0.77
1:D:1028:LYS:O	1:D:1030:PRO:HD3	1.87	0.74
1:D:1028:LYS:HD3	1:D:1090:LEU:HD21	1.69	0.74

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	162/164 (99%)	143 (88%)	15 (9%)	4 (2%)	7	4
1	B	162/164 (99%)	134 (83%)	21 (13%)	7 (4%)	3	1
1	C	162/164 (99%)	141 (87%)	20 (12%)	1 (1%)	30	32
1	D	162/164 (99%)	134 (83%)	27 (17%)	1 (1%)	30	32
1	E	162/164 (99%)	143 (88%)	18 (11%)	1 (1%)	30	32
1	F	162/164 (99%)	138 (85%)	22 (14%)	2 (1%)	16	14
2	G	4/6 (67%)	3 (75%)	1 (25%)	0	100	100
2	H	4/6 (67%)	3 (75%)	1 (25%)	0	100	100
2	I	4/6 (67%)	3 (75%)	1 (25%)	0	100	100
2	J	4/6 (67%)	3 (75%)	1 (25%)	0	100	100
2	K	4/6 (67%)	3 (75%)	1 (25%)	0	100	100
2	L	4/6 (67%)	3 (75%)	1 (25%)	0	100	100
All	All	996/1020 (98%)	851 (85%)	129 (13%)	16 (2%)	12	9

5 of 16 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1043	GLU
1	B	1086	GLU
1	B	1105	PRO
1	B	1081	GLU
1	C	1151	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	132/132 (100%)	125 (95%)	7 (5%)	28	34
1	B	132/132 (100%)	118 (89%)	14 (11%)	8	8
1	C	132/132 (100%)	125 (95%)	7 (5%)	28	34
1	D	132/132 (100%)	124 (94%)	8 (6%)	23	27
1	E	132/132 (100%)	121 (92%)	11 (8%)	14	14
1	F	132/132 (100%)	123 (93%)	9 (7%)	20	22
2	G	4/4 (100%)	3 (75%)	1 (25%)	1	0
2	H	4/4 (100%)	3 (75%)	1 (25%)	1	0
2	I	4/4 (100%)	3 (75%)	1 (25%)	1	0
2	J	4/4 (100%)	2 (50%)	2 (50%)	0	0
2	K	4/4 (100%)	4 (100%)	0	100	100
2	L	4/4 (100%)	3 (75%)	1 (25%)	1	0
All	All	816/816 (100%)	754 (92%)	62 (8%)	16	17

5 of 62 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	1013	ASP
1	D	1129	PHE
2	H	101	HIS
1	D	1061	MET
1	E	1039	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	1126	HIS
1	F	1137	ASN
1	C	1149	ASN
1	B	1003	ASN
1	B	1149	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	164/164 (100%)	0.99	13 (7%) 15 23	5, 13, 20, 23	0
1	B	164/164 (100%)	2.28	88 (53%) 0 0	11, 20, 28, 34	0
1	C	164/164 (100%)	1.26	30 (18%) 2 3	5, 16, 25, 35	0
1	D	164/164 (100%)	1.40	34 (20%) 1 2	7, 18, 25, 33	0
1	E	164/164 (100%)	1.04	24 (14%) 3 6	3, 12, 21, 27	0
1	F	164/164 (100%)	2.14	83 (50%) 0 0	10, 18, 26, 30	0
2	G	4/6 (66%)	1.24	0 100 100	17, 19, 21, 29	0
2	H	6/6 (100%)	1.40	1 (16%) 2 4	11, 16, 22, 24	0
2	I	6/6 (100%)	1.20	0 100 100	7, 10, 11, 26	0
2	J	6/6 (100%)	1.61	1 (16%) 2 4	9, 10, 12, 18	0
2	K	4/6 (66%)	2.25	1 (25%) 1 2	12, 13, 20, 23	0
2	L	6/6 (100%)	1.72	2 (33%) 0 0	16, 19, 24, 24	0
All	All	1016/1020 (99%)	1.52	277 (27%) 1 1	3, 17, 26, 35	0

The worst 5 of 277 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1104	GLY	12.2
1	F	1107	THR	7.3
1	F	1080	GLY	6.1
1	F	1127	VAL	5.9
1	D	1080	GLY	5.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.