



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 22, 2016 – 09:43 PM EDT

PDB ID : 5AXH
Title : Crystal structure of thermophilic dextranase from Thermoanaerobacter pseudethanolicus, D312G mutant in complex with isomaltohexaose
Authors : Suzuki, N.; Kishine, N.; Fujimoto, Z.; Sakurai, M.; Momma, M.; Ko, J.A.; Nam, S.H.; Kimura, A.; Kim, Y.M.
Deposited on : 2015-07-29
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027107
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0122
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027107

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLC	A	701	-	-	-	X
3	PO4	B	708	-	-	-	X
4	GOL	A	708	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 10326 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dextranase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	586	Total	C	N	O	S	0	1	0
			4786	3084	792	899	11			
1	B	587	Total	C	N	O	S	0	4	0
			4802	3091	796	904	11			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP B0KBZ7
A	2	GLY	-	expression tag	UNP B0KBZ7
A	312	GLY	ASP	engineered mutation	UNP B0KBZ7
A	611	LEU	-	expression tag	UNP B0KBZ7
A	612	GLU	-	expression tag	UNP B0KBZ7
A	613	HIS	-	expression tag	UNP B0KBZ7
A	614	HIS	-	expression tag	UNP B0KBZ7
A	615	HIS	-	expression tag	UNP B0KBZ7
A	616	HIS	-	expression tag	UNP B0KBZ7
A	617	HIS	-	expression tag	UNP B0KBZ7
A	618	HIS	-	expression tag	UNP B0KBZ7
B	1	MET	-	expression tag	UNP B0KBZ7
B	2	GLY	-	expression tag	UNP B0KBZ7
B	312	GLY	ASP	engineered mutation	UNP B0KBZ7
B	611	LEU	-	expression tag	UNP B0KBZ7
B	612	GLU	-	expression tag	UNP B0KBZ7
B	613	HIS	-	expression tag	UNP B0KBZ7
B	614	HIS	-	expression tag	UNP B0KBZ7
B	615	HIS	-	expression tag	UNP B0KBZ7
B	616	HIS	-	expression tag	UNP B0KBZ7
B	617	HIS	-	expression tag	UNP B0KBZ7
B	618	HIS	-	expression tag	UNP B0KBZ7

- Molecule 2 is ALPHA-D-GLUCOSE (three-letter code: GLC) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			11	6	5		
2	B	1	Total	C	O	0	0
			11	6	5		
2	B	1	Total	C	O	0	0
			11	6	5		
2	B	1	Total	C	O	0	0
			11	6	5		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		
3	A	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		
3	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is BETA-D-GLUCOSE (three-letter code: BGC) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			12	6	6		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	303	Total	O	0	0
			303	303		
6	B	291	Total	O	0	0
			291	291		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Dextranase



• Molecule 1: Dextranase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	78.33Å 99.18Å 169.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.92 – 2.20 38.92 – 2.20	Depositor EDS
% Data completeness (in resolution range)	98.3 (38.92-2.20) 98.4 (38.92-2.20)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.69 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0124	Depositor
R, R_{free}	0.160 , 0.205 0.168 , 0.210	Depositor DCC
R_{free} test set	3350 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	30.7	Xtriage
Anisotropy	0.031	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 34.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 66563 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10326	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, GOL, PO4, BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/4916	0.68	0/6660
1	B	0.52	0/4947	0.67	0/6702
All	All	0.52	0/9863	0.68	0/13362

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4786	0	4651	9	0
1	B	4802	0	4664	6	0
2	A	45	0	39	0	0
2	B	55	0	46	0	0
3	A	10	0	0	0	0
3	B	10	0	0	0	0
4	A	12	0	16	1	0
5	B	12	0	11	0	0
6	A	303	0	0	1	0
6	B	291	0	0	0	0
All	All	10326	0	9427	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:183:PHE:CE1	1:A:232:VAL:HG13	2.41	0.55
1:B:352:ASN:HD22	1:B:374:GLU:CD	2.10	0.54
1:A:222:ILE:HG12	1:A:232:VAL:HG11	1.91	0.51
1:B:375:ILE:HD11	1:B:386:LEU:HD23	1.93	0.50
1:B:352:ASN:ND2	1:B:374:GLU:OE2	2.43	0.49
1:B:183:PHE:CE1	1:B:232:VAL:HG13	2.47	0.49
1:A:493:ILE:HD11	1:A:513:ALA:HB2	1.96	0.47
1:A:163:LYS:NZ	1:A:223:GLU:OE1	2.48	0.46
1:A:374:GLU:CD	1:A:376:TRP:HE1	2.19	0.46
1:B:522:ILE:HD11	1:B:555:TYR:CG	2.52	0.45
1:A:361:ASN:HB3	6:A:997:HOH:O	2.18	0.43
1:B:522:ILE:HD11	1:B:555:TYR:CD1	2.54	0.42
1:A:420:PHE:CE2	1:A:447:MET:HB3	2.54	0.42
1:A:382:HIS:HA	1:A:533:SER:O	2.19	0.42
1:A:470:ASN:OD1	4:A:708:GOL:H11	2.22	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	585/618 (95%)	565 (97%)	19 (3%)	1 (0%)	52	59
1	B	589/618 (95%)	571 (97%)	18 (3%)	0	100	100
All	All	1174/1236 (95%)	1136 (97%)	37 (3%)	1 (0%)	56	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	262	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	510/536 (95%)	501 (98%)	9 (2%)	66	79
1	B	514/536 (96%)	508 (99%)	6 (1%)	78	88
All	All	1024/1072 (96%)	1009 (98%)	15 (2%)	72	84

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	ASP
1	A	39	THR
1	A	91	LYS
1	A	146	TRP
1	A	152	TYR
1	A	162	SER
1	A	239	TYR
1	A	558	GLU
1	A	559	ASN
1	B	30	LYS
1	B	35	ILE
1	B	146	TRP
1	B	152	TYR
1	B	239	TYR
1	B	559	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	130	ASN
1	A	515	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	A	701	2	11,11,12	0.55	0	15,15,17	1.75	4 (26%)
2	GLC	A	702	2	11,11,12	0.48	0	15,15,17	0.86	0
2	GLC	A	703	2	11,11,12	0.50	0	15,15,17	0.91	0
2	GLC	A	704	2	12,12,12	0.52	0	17,17,17	1.89	5 (29%)
3	PO4	A	705	-	4,4,4	0.66	0	6,6,6	0.22	0
3	PO4	A	706	-	4,4,4	0.65	0	6,6,6	0.25	0
4	GOL	A	707	-	5,5,5	0.35	0	5,5,5	0.40	0
4	GOL	A	708	-	5,5,5	0.52	0	5,5,5	1.56	2 (40%)
2	GLC	B	701	2	11,11,12	0.45	0	15,15,17	1.80	3 (20%)
2	GLC	B	702	2	11,11,12	0.56	0	15,15,17	0.88	1 (6%)
2	GLC	B	703	2	11,11,12	0.37	0	15,15,17	0.97	1 (6%)
2	GLC	B	704	2	11,11,12	0.44	0	15,15,17	1.03	1 (6%)
2	GLC	B	705	2,5	11,11,12	0.38	0	15,15,17	1.03	0
5	BGC	B	706	2	12,12,12	0.46	0	17,17,17	1.10	1 (5%)
3	PO4	B	707	-	4,4,4	0.52	0	6,6,6	0.25	0
3	PO4	B	708	-	4,4,4	0.53	0	6,6,6	0.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	701	2	-	0/2/19/22	0/1/1/1
2	GLC	A	702	2	-	0/2/19/22	0/1/1/1
2	GLC	A	703	2	-	0/2/19/22	0/1/1/1
2	GLC	A	704	2	-	0/2/22/22	0/1/1/1
3	PO4	A	705	-	-	0/0/0/0	0/0/0/0
3	PO4	A	706	-	-	0/0/0/0	0/0/0/0
4	GOL	A	707	-	-	0/4/4/4	0/0/0/0
4	GOL	A	708	-	-	0/4/4/4	0/0/0/0
2	GLC	B	701	2	-	0/2/19/22	0/1/1/1
2	GLC	B	702	2	-	0/2/19/22	0/1/1/1
2	GLC	B	703	2	-	0/2/19/22	0/1/1/1
2	GLC	B	704	2	-	0/2/19/22	0/1/1/1
2	GLC	B	705	2,5	-	0/2/19/22	0/1/1/1
5	BGC	B	706	2	-	0/2/22/22	0/1/1/1
3	PO4	B	707	-	-	0/0/0/0	0/0/0/0
3	PO4	B	708	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	704	GLC	O5-C1-C2	-3.03	106.06	110.89
2	B	703	GLC	O5-C1-C2	-3.00	106.10	110.89
2	A	704	GLC	O3-C3-C2	-2.80	104.06	110.36
2	B	702	GLC	O5-C1-C2	-2.48	106.93	110.89
2	A	701	GLC	O4-C4-C3	-2.09	105.66	110.36
2	A	704	GLC	O5-C5-C4	2.03	113.55	109.67
2	A	701	GLC	O5-C1-C2	2.06	114.19	110.89
4	A	708	GOL	O1-C1-C2	2.16	120.94	109.97
4	A	708	GOL	O3-C3-C2	2.52	122.73	109.97
2	A	701	GLC	C3-C4-C5	2.54	114.75	110.23
5	B	706	BGC	C1-O5-C5	2.68	118.66	113.54
2	A	704	GLC	C1-C2-C3	2.69	115.08	110.68
2	B	701	GLC	C3-C4-C5	3.14	115.83	110.23
2	B	701	GLC	O5-C5-C4	3.26	115.53	110.13
2	A	704	GLC	C1-O5-C5	3.62	120.46	113.54
2	B	701	GLC	C1-O5-C5	3.78	117.70	112.14
2	A	704	GLC	O5-C1-C2	4.32	117.56	110.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	701	GLC	C1-O5-C5	5.02	119.52	112.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	708	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	586/618 (94%)	-0.17	28 (4%) 34 34	20, 31, 60, 105	0
1	B	587/618 (94%)	-0.21	16 (2%) 58 57	20, 31, 57, 100	0
All	All	1173/1236 (94%)	-0.19	44 (3%) 44 43	20, 31, 59, 105	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	33	ASN	8.4
1	A	33	ASN	7.9
1	A	32	ASP	7.3
1	B	32	ASP	5.6
1	A	558	GLU	4.6
1	B	558	GLU	4.6
1	A	131	LYS	4.5
1	A	272	HIS	4.5
1	A	31	ASP	4.3
1	A	35	ILE	4.1
1	A	128	LYS	4.1
1	B	35	ILE	4.0
1	B	560	ASP	4.0
1	A	71	GLN	4.0
1	B	31	ASP	3.9
1	B	561	ASN	3.7
1	A	34	ASN	3.7
1	A	611	LEU	3.7
1	A	129	GLY	3.5
1	A	26	ILE	3.2
1	A	277	ASP	3.2
1	B	34	ASN	3.1
1	A	30	LYS	3.0
1	A	264	GLU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	611	LEU	2.7
1	A	377	PRO	2.7
1	A	130	ASN	2.7
1	A	375	ILE	2.6
1	A	27	PHE	2.6
1	A	37	ALA	2.6
1	B	131	LYS	2.6
1	A	36	VAL	2.6
1	A	29	PHE	2.5
1	B	559	ASN	2.5
1	A	38	LEU	2.5
1	B	29	PHE	2.3
1	A	77	THR	2.3
1	A	72	GLU	2.2
1	B	416	ARG	2.2
1	A	262	HIS	2.2
1	B	343[A]	ASN	2.1
1	B	103	LYS	2.1
1	A	70	LEU	2.1
1	B	130	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	PO4	B	708	5/5	0.94	0.29	4.42	74,75,78,80	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	GOL	A	708	6/6	0.91	0.22	3.39	39,43,45,47	0
2	GLC	A	701	11/12	0.86	0.18	2.58	51,61,63,65	0
2	GLC	B	703	11/12	0.96	0.13	1.73	27,29,31,31	0
3	PO4	B	707	5/5	0.94	0.18	1.16	55,57,61,62	0
5	BGC	B	706	12/12	0.86	0.17	1.08	56,76,78,81	0
4	GOL	A	707	6/6	0.88	0.14	0.75	54,58,58,59	0
3	PO4	A	705	5/5	0.91	0.12	0.23	79,82,84,85	0
2	GLC	B	701	11/12	0.92	0.12	0.06	50,52,57,59	0
2	GLC	B	705	11/12	0.92	0.13	-0.43	40,49,52,55	0
2	GLC	B	704	11/12	0.96	0.12	-0.48	31,32,35,38	0
2	GLC	A	704	12/12	0.97	0.08	-0.70	30,33,37,39	0
2	GLC	A	702	11/12	0.97	0.08	-0.89	34,37,43,46	0
2	GLC	A	703	11/12	0.97	0.07	-0.95	24,30,33,36	0
2	GLC	B	702	11/12	0.96	0.08	-1.55	34,36,39,43	0
3	PO4	A	706	5/5	0.96	0.12	-	67,69,72,73	0

6.5 Other polymers [i](#)

There are no such residues in this entry.