



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:34 AM GMT

PDB ID : 3BBT
Title : crystal structure of the ErbB4 kinase in complex with lapatinib
Authors : Qiu, C.
Deposited on : 2007-11-11
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	328	<div> <div>6%</div> <div> <div></div> <div>69%</div> <div>11%</div> <div>•</div> <div>16%</div> </div> </div>
1	D	328	<div> <div>4%</div> <div> <div></div> <div>65%</div> <div>14%</div> <div>21%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	FMM	B	91	-	-	-	X
2	FMM	D	91	-	-	-	X

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4431 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Receptor tyrosine-protein kinase erbB-4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	274	Total	C	N	O	S	0	0	0
			2201	1420	380	382	19			
1	D	259	Total	C	N	O	S	0	0	0
			2091	1352	361	360	18			

- Molecule 2 is N-{3-CHLORO-4-[(3-FLUOROBENZYL)OXY]PHENYL}-6-[5-([2-(METHYLSULFONYL)ETHYL]AMINO)METHYL]-2-FURYL]-4-QUINAZOLINAMINE (three-letter code: FMM) (formula: C₂₉H₂₆ClFN₄O₄S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	B	1	Total	C	Cl	F	N	O	0	0
			27	21	1	1	3	1		
2	D	1	Total	C	Cl	F	N	O	0	0
			28	22	1	1	3	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	44	Total 44	O 44	0	0
3	D	40	Total 40	O 40	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	102.68Å 102.68Å 185.12Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.64 – 2.80 29.64 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (29.64-2.80) 99.9 (29.64-2.80)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 2.80Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.251 , 0.289 0.245 , 0.281	Depositor DCC
R_{free} test set	1381 reflections (5.37%)	DCC
Wilson B-factor (Å ²)	65.0	Xtriage
Anisotropy	0.358	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 38.8	EDS
Estimated twinning fraction	0.334 for h,-h-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.40$, $\langle L^2 \rangle = 0.23$	Xtriage
Outliers	0 of 27083 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4431	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FMM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.35	0/2251	0.56	3/3043 (0.1%)
1	D	0.34	0/2139	0.51	1/2891 (0.0%)
All	All	0.34	0/4390	0.53	4/5934 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	4

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	965	ASP	CB-CG-OD2	5.65	123.39	118.30
1	D	818	ASP	CB-CG-OD2	5.51	123.26	118.30
1	B	818	ASP	CB-CG-OD2	5.21	122.99	118.30
1	B	966	ASP	CB-CG-OD2	5.00	122.80	118.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	720	LYS	Peptide
1	B	721	ILE	Peptide
1	B	733	GLY	Peptide
1	B	972	SER	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2201	0	2264	38	0
1	D	2091	0	2153	31	0
2	B	27	0	14	0	0
2	D	28	0	14	4	0
3	B	44	0	0	0	0
3	D	40	0	0	0	0
All	All	4431	0	4445	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:720:LYS:O	1:B:721:ILE:HG12	1.39	1.21
1:B:720:LYS:C	1:B:721:ILE:HG12	1.55	1.09
1:B:965:ASP:CG	1:B:966:ASP:H	1.81	0.84
1:B:720:LYS:C	1:B:721:ILE:CG1	2.42	0.82
1:B:721:ILE:N	1:B:722:PRO:HD2	1.95	0.81
1:D:816:HIS:O	1:D:817:ARG:HB2	1.80	0.78
1:B:733:GLY:O	1:B:843:LEU:HD21	1.88	0.74
1:B:696:VAL:O	1:B:697:LYS:HG3	1.90	0.72
1:B:727:ILE:HG12	1:B:768:GLN:HG2	1.73	0.71
1:B:816:HIS:O	1:B:817:ARG:HB2	1.89	0.71
1:B:817:ARG:NH1	1:B:840:ALA:HB1	2.04	0.70
1:B:720:LYS:O	1:B:721:ILE:CG1	2.30	0.68
1:D:886:TRP:HD1	1:D:928:MET:CE	2.07	0.68
1:D:727:ILE:HG12	1:D:768:GLN:HG2	1.79	0.65
1:B:965:ASP:OD1	1:B:966:ASP:N	2.30	0.65
1:B:965:ASP:CG	1:B:966:ASP:N	2.50	0.65
1:B:963:GLN:NE2	1:B:969:LYS:H	1.96	0.63
1:D:713:VAL:HG22	1:D:720:LYS:HG2	1.82	0.62
1:B:817:ARG:NH1	1:B:840:ALA:CB	2.63	0.61
1:B:696:VAL:O	1:B:697:LYS:CG	2.50	0.59
1:D:885:ILE:O	1:D:889:MET:HG2	2.05	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:904:ILE:N	1:D:905:PRO:HD2	2.21	0.55
1:B:772:GLN:HE22	1:B:827:LYS:HE2	1.72	0.55
1:D:886:TRP:HD1	1:D:928:MET:HE1	1.71	0.55
1:B:951:ARG:HA	1:B:954:ARG:HH11	1.73	0.54
1:D:886:TRP:HD1	1:D:928:MET:HE3	1.73	0.53
1:B:696:VAL:C	1:B:697:LYS:HG3	2.28	0.53
1:D:815:VAL:HG12	1:D:817:ARG:HG3	1.91	0.52
1:B:726:LYS:HB3	1:B:769:LEU:HB2	1.91	0.51
1:B:723:VAL:HG22	1:B:771:THR:O	2.11	0.51
1:D:718:THR:HG23	1:D:719:VAL:HG13	1.93	0.50
1:D:769:LEU:HD23	2:D:91:FMM:H282	1.94	0.49
1:B:735:LYS:O	1:B:737:ASN:OD1	2.30	0.49
1:D:689:LYS:HB2	1:D:692:GLU:HB2	1.93	0.48
1:B:963:GLN:HE22	1:B:969:LYS:HB2	1.78	0.48
1:D:816:HIS:ND1	1:D:816:HIS:C	2.67	0.48
1:B:733:GLY:O	1:B:843:LEU:CD2	2.60	0.48
1:B:864:LEU:HD21	1:B:909:GLU:HG2	1.95	0.48
1:B:736:ALA:C	1:B:737:ASN:OD1	2.52	0.48
1:D:792:SER:HB3	1:D:960:LEU:HB2	1.95	0.47
1:D:752:HIS:CD2	1:D:754:HIS:H	2.33	0.47
1:D:754:HIS:CE1	1:D:801:GLN:HG2	2.49	0.47
1:D:884:THR:O	1:D:887:GLU:HB2	2.15	0.47
1:D:906:ASP:N	1:D:906:ASP:OD1	2.46	0.47
1:B:750:MET:HE1	1:B:809:LEU:HD23	1.97	0.47
1:B:735:LYS:O	1:B:737:ASN:N	2.48	0.46
1:D:776:HIS:HB2	1:D:826:VAL:HB	1.96	0.46
1:D:886:TRP:CD1	1:D:928:MET:HE1	2.51	0.45
1:D:886:TRP:CD1	1:D:928:MET:CE	2.96	0.44
1:D:837:PHE:CZ	2:D:91:FMM:H32	2.53	0.44
1:B:696:VAL:HG23	1:B:709:LYS:O	2.17	0.44
1:B:720:LYS:HB2	1:D:720:LYS:H	1.82	0.43
1:D:774:MET:HA	1:D:775:PRO:HD3	1.83	0.43
1:B:701:SER:HB3	1:B:706:THR:HG23	1.99	0.43
1:D:777:GLY:HA2	2:D:91:FMM:C16	2.49	0.42
1:D:946:ALA:O	1:D:950:SER:HB2	2.18	0.42
1:D:721:ILE:HA	1:D:722:PRO:HD2	1.78	0.42
1:B:809:LEU:HD22	1:B:814:LEU:HD22	2.01	0.42
1:D:774:MET:H	2:D:91:FMM:H19	1.84	0.42
1:D:894:LYS:HA	1:D:895:PRO:HD3	1.91	0.41
1:B:817:ARG:HH11	1:B:840:ALA:CB	2.34	0.41
1:B:816:HIS:ND1	1:B:816:HIS:C	2.73	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:963:GLN:HE22	1:B:969:LYS:H	1.68	0.41
1:B:750:MET:CE	1:B:809:LEU:HD23	2.50	0.41
1:D:886:TRP:CD1	1:D:928:MET:HE3	2.54	0.41
1:D:762:CYS:HB3	1:D:768:GLN:HB2	2.03	0.40
1:B:688:LEU:HD12	1:B:770:VAL:HG21	2.03	0.40
1:B:734:PRO:O	1:B:767:ILE:HD13	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	266/328 (81%)	256 (96%)	6 (2%)	4 (2%)	13	40
1	D	251/328 (76%)	244 (97%)	6 (2%)	1 (0%)	39	74
All	All	517/656 (79%)	500 (97%)	12 (2%)	5 (1%)	19	52

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	722	PRO
1	B	736	ALA
1	B	734	PRO
1	B	765	PRO
1	D	685	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	243/288 (84%)	231 (95%)	12 (5%)	31	65
1	D	231/288 (80%)	227 (98%)	4 (2%)	68	92
All	All	474/576 (82%)	458 (97%)	16 (3%)	44	78

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	696	VAL
1	B	713	VAL
1	B	721	ILE
1	B	742	ASP
1	B	771	THR
1	B	773	LEU
1	B	788	ASP
1	B	913	ARG
1	B	924	VAL
1	B	958	ARG
1	B	963	GLN
1	B	972	SER
1	D	723	VAL
1	D	728	LEU
1	D	835	THR
1	D	901	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	957	GLN
1	B	963	GLN
1	D	752	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FMM	B	91	-	30,30,44	1.20	2 (6%)	41,41,62	2.63	11 (26%)
2	FMM	D	91	-	31,31,44	1.18	2 (6%)	43,43,62	2.72	11 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMM	B	91	-	-	0/9/9/21	0/4/4/5
2	FMM	D	91	-	-	0/9/9/21	0/4/4/5

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	91	FMM	C21-C39	-3.92	1.40	1.44
2	B	91	FMM	C21-C39	-3.83	1.40	1.44
2	D	91	FMM	C36-CL3	2.61	1.80	1.73
2	B	91	FMM	C36-CL3	2.70	1.80	1.73

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	91	FMM	N18-C19-N20	-10.39	120.94	128.89
2	B	91	FMM	N18-C19-N20	-10.26	121.03	128.89
2	D	91	FMM	C40-C39-C21	-6.25	121.56	124.89
2	B	91	FMM	C39-C21-N20	-3.92	118.67	121.46
2	D	91	FMM	C39-C21-N20	-3.70	118.82	121.46
2	B	91	FMM	C39-C17-N18	-3.01	119.68	122.88
2	D	91	FMM	C23-N22-C21	-2.81	122.21	128.40
2	D	91	FMM	C32-C33-C35	-2.74	119.82	123.35
2	D	91	FMM	C39-C17-N18	-2.72	119.99	122.88
2	B	91	FMM	C32-C33-C35	-2.43	120.22	123.35
2	B	91	FMM	C23-N22-C21	-2.11	123.75	128.40
2	D	91	FMM	C28-O27-C26	2.07	121.93	117.82
2	D	91	FMM	N22-C21-N20	2.09	120.83	118.85
2	B	91	FMM	O27-C26-C36	2.17	119.21	116.36
2	B	91	FMM	N22-C21-N20	2.39	121.12	118.85
2	B	91	FMM	C28-O27-C26	2.53	122.83	117.82
2	D	91	FMM	C21-C39-C17	4.64	118.41	115.77
2	B	91	FMM	C21-C39-C17	5.32	118.80	115.77
2	D	91	FMM	C19-N18-C17	5.69	120.45	115.19
2	B	91	FMM	C19-N18-C17	5.76	120.51	115.19
2	B	91	FMM	C19-N20-C21	6.70	121.31	116.48
2	D	91	FMM	C19-N20-C21	6.83	121.40	116.48

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	91	FMM	4	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	B	274/328 (83%)	0.76	19 (6%)	20 11	16, 23, 39, 42	0
1	D	259/328 (78%)	0.72	13 (5%)	32 21	16, 23, 39, 41	0
All	All	533/656 (81%)	0.74	32 (6%)	25 15	16, 23, 39, 42	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	843	LEU	6.6
1	B	704	PHE	6.1
1	B	717	GLU	5.1
1	D	704	PHE	5.1
1	D	715	GLU	4.2
1	D	734	PRO	4.1
1	D	735	LYS	3.7
1	B	842	LEU	3.6
1	B	714	PRO	3.4
1	B	722	PRO	2.9
1	B	898	GLY	2.7
1	B	738	VAL	2.7
1	B	868	HIS	2.6
1	D	697	LYS	2.5
1	D	864	LEU	2.5
1	B	902	ARG	2.5
1	D	963	GLN	2.4
1	B	864	LEU	2.4
1	B	764	SER	2.4
1	B	715	GLU	2.4
1	B	728	LEU	2.4
1	B	684	GLN	2.4
1	B	711	ILE	2.4
1	D	707	VAL	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	902	ARG	2.3
1	B	712	TRP	2.2
1	B	729	ASN	2.2
1	D	769	LEU	2.2
1	B	908	LEU	2.1
1	D	714	PRO	2.1
1	D	746	ILE	2.0
1	D	870	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	FMM	B	91	27/40	0.54	0.62	7.46	122,123,123,123	0
2	FMM	D	91	28/40	0.70	0.47	3.47	99,100,100,100	0

6.5 Other polymers [i](#)

There are no such residues in this entry.