



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:39 AM GMT

PDB ID : 3BDM  
Title : yeast 20S proteasome:glidobactin A-complex  
Authors : Groll, M.; Dudler, R.; Kaiser, M.  
Deposited on : 2007-11-15  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	<div> <div>2%</div> <div> <div></div> <div>66%</div> <div>32%</div> <div>.</div> </div> </div>
1	O	250	<div> <div>7%</div> <div> <div></div> <div>66%</div> <div>31%</div> <div>.</div> </div> </div>
2	B	258	<div> <div>5%</div> <div> <div></div> <div>53%</div> <div>37%</div> <div>5%</div> <div>5%</div> </div> </div>
2	P	258	<div> <div>6%</div> <div> <div></div> <div>53%</div> <div>37%</div> <div>5%</div> <div>5%</div> </div> </div>
3	C	254	<div> <div>6%</div> <div> <div></div> <div>50%</div> <div>40%</div> <div>5%</div> <div>5%</div> </div> </div>
3	Q	254	<div> <div>14%</div> <div> <div></div> <div>50%</div> <div>41%</div> <div>5%</div> <div>5%</div> </div> </div>
4	D	260	<div> <div>3%</div> <div> <div></div> <div>59%</div> <div>31%</div> <div>.</div> <div>7%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	R	260	
5	E	234	
5	S	234	
6	F	287	
6	T	287	
7	G	252	
7	U	252	
8	H	232	
8	V	232	
9	I	205	
9	W	205	
10	J	198	
10	X	198	
11	K	212	
11	Y	212	
12	L	241	
12	Z	241	
13	O	266	
13	M	266	
14	1	196	
14	N	196	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
15	GDT	V	500	-	-	-	X

## 2 Entry composition

There are 16 unique types of molecules in this entry. The entry contains 51022 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteasome component Y7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	250	Total	C	N	O	S	0	0	0
			1915	1219	315	377	4			
1	O	250	Total	C	N	O	S	0	0	0
			1915	1219	315	377	4			

- Molecule 2 is a protein called Proteasome component Y13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	244	Total	C	N	O	S	0	0	0
			1904	1201	321	379	3			
2	P	244	Total	C	N	O	S	0	0	0
			1904	1201	321	379	3			

- Molecule 3 is a protein called Proteasome component PRE6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	241	Total	C	N	O	S	0	0	0
			1890	1181	331	374	4			
3	Q	241	Total	C	N	O	S	0	0	0
			1890	1181	331	374	4			

- Molecule 4 is a protein called Proteasome component PUP2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	242	Total	C	N	O	S	0	0	0
			1861	1162	314	378	7			
4	R	242	Total	C	N	O	S	0	0	0
			1861	1162	314	378	7			

- Molecule 5 is a protein called Proteasome component PRE5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	233	Total	C	N	O	S	0	0	0
			1795	1129	312	350	4			
5	S	233	Total	C	N	O	S	0	0	0
			1795	1129	312	350	4			

- Molecule 6 is a protein called Proteasome component C1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	244	Total	C	N	O	S	0	0	0
			1896	1205	330	357	4			
6	T	244	Total	C	N	O	S	0	0	0
			1896	1205	330	357	4			

- Molecule 7 is a protein called Proteasome component C7-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	243	Total	C	N	O	S	0	0	0
			1921	1221	322	370	8			
7	U	243	Total	C	N	O	S	0	0	0
			1921	1221	322	370	8			

- Molecule 8 is a protein called Proteasome component PUP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	222	Total	C	N	O	S	0	0	0
			1684	1061	293	323	7			
8	V	222	Total	C	N	O	S	0	0	0
			1684	1061	293	323	7			

- Molecule 9 is a protein called Proteasome component PUP3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	204	Total	C	N	O	S	0	0	0
			1581	1010	258	305	8			
9	W	204	Total	C	N	O	S	0	0	0
			1581	1010	258	305	8			

- Molecule 10 is a protein called Proteasome component C11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	198	Total	C	N	O	S	0	0	0
			1585	1005	269	305	6			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	X	198	Total	C	N	O	S	0	0	0
			1585	1005	269	305	6			

- Molecule 11 is a protein called Proteasome component PRE2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			
11	Y	212	Total	C	N	O	S	0	0	0
			1644	1045	280	312	7			

- Molecule 12 is a protein called Proteasome component C5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			
12	Z	222	Total	C	N	O	S	0	0	0
			1757	1115	303	335	4			


- Molecule 13 is a protein called Proteasome component PRE4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	M	233	Total	C	N	O	S	0	0	0
			1824	1154	312	351	7			
13	0	233	Total	C	N	O	S	0	0	0
			1824	1154	312	351	7			

- Molecule 14 is a protein called Proteasome component PRE3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	N	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			
14	1	196	Total	C	N	O	S	0	0	0
			1512	955	250	300	7			

- Molecule 15 is (2E,4E)-N-[(2S,3R)-3-HYDROXY-1-[(3Z,5S,8S,10S)-10-HYDROXY-5-METHYL-2,7-DIOXO-1,6-DIAZACYCLODEDEC-3-EN-8-YL]AMINO]-1-OXOBUTAN-2-YL]DODECA-2,4-DIENAMIDE (three-letter code: GDT) (formula: C<sub>27</sub>H<sub>44</sub>N<sub>4</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
15	H	1	Total	C	N	O	0	0
			37	27	4	6		
15	K	1	Total	C	N	O	0	0
			37	27	4	6		
15	V	1	Total	C	N	O	0	0
			37	27	4	6		
15	Y	1	Total	C	N	O	0	0
			37	27	4	6		

- Molecule 16 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	A	57	Total	O	0	0
			57	57		
16	B	38	Total	O	0	0
			38	38		
16	C	42	Total	O	0	0
			42	42		
16	D	40	Total	O	0	0
			40	40		
16	E	24	Total	O	0	0
			24	24		
16	F	46	Total	O	0	0
			46	46		
16	G	61	Total	O	0	0
			61	61		
16	H	48	Total	O	0	0
			48	48		


Continued on next page...

*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	I	66	Total 66	O 66	0	0
16	J	52	Total 52	O 52	0	0
16	K	46	Total 46	O 46	0	0
16	L	60	Total 60	O 60	0	0
16	M	69	Total 69	O 69	0	0
16	N	56	Total 56	O 56	0	0
16	O	33	Total 33	O 33	0	0
16	P	32	Total 32	O 32	0	0
16	Q	26	Total 26	O 26	0	0
16	R	34	Total 34	O 34	0	0
16	S	20	Total 20	O 20	0	0
16	T	39	Total 39	O 39	0	0
16	U	58	Total 58	O 58	0	0
16	V	47	Total 47	O 47	0	0
16	W	60	Total 60	O 60	0	0
16	X	42	Total 42	O 42	0	0
16	Y	49	Total 49	O 49	0	0
16	Z	53	Total 53	O 53	0	0
16	0	74	Total 74	O 74	0	0
16	1	64	Total 64	O 64	0	0









• Molecule 2: Proteasome component Y13




• Molecule 3: Proteasome component PRE6




• Molecule 3: Proteasome component PRE6







- Molecule 4: Proteasome component PUP2



- Molecule 4: Proteasome component PUP2




- Molecule 5: Proteasome component PRE5




- Molecule 5: Proteasome component PRE5






• Molecule 7: Proteasome component C7-alpha




• Molecule 8: Proteasome component PUP1




• Molecule 8: Proteasome component PUP1




• Molecule 9: Proteasome component PUP3




• Molecule 9: Proteasome component PUP3




• Molecule 10: Proteasome component C11




• Molecule 10: Proteasome component C11




- Molecule 11: Proteasome component PRE2




- Molecule 11: Proteasome component PRE2




- Molecule 12: Proteasome component C5




- Molecule 12: Proteasome component C5







- Molecule 13: Proteasome component PRE4



- Molecule 13: Proteasome component PRE4




- Molecule 14: Proteasome component PRE3



- Molecule 14: Proteasome component PRE3







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	132.77Å 302.79Å 143.21Å 90.00° 112.24° 90.00°	Depositor
Resolution (Å)	15.00 – 2.70 49.11 – 2.70	Depositor EDS
% Data completeness (in resolution range)	97.9 (15.00-2.70) 97.9 (49.11-2.70)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.89 (at 2.69Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.234 , 0.262 0.247 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	46.0	Xtriage
Anisotropy	0.658	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 62.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	0 of 278774 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	51022	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.48% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GDT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/1952	0.64	0/2642
1	O	0.38	0/1952	0.64	0/2642
2	B	0.38	0/1934	0.63	0/2618
2	P	0.38	0/1934	0.63	0/2618
3	C	0.37	0/1919	0.63	0/2598
3	Q	0.36	0/1919	0.63	0/2598
4	D	0.38	0/1886	0.64	0/2541
4	R	0.37	0/1886	0.63	0/2541
5	E	0.36	0/1823	0.62	0/2463
5	S	0.37	0/1823	0.62	0/2463
6	F	0.39	0/1936	0.64	0/2614
6	T	0.39	0/1936	0.64	0/2614
7	G	0.42	0/1959	0.66	0/2652
7	U	0.41	0/1959	0.66	0/2652
8	H	0.39	0/1715	0.67	0/2326
8	V	0.39	0/1715	0.66	0/2326
9	I	0.41	0/1611	0.66	0/2174
9	W	0.42	0/1611	0.67	0/2174
10	J	0.41	0/1613	0.66	0/2173
10	X	0.42	0/1613	0.68	0/2173
11	K	0.40	0/1681	0.66	0/2274
11	Y	0.40	0/1681	0.65	0/2274
12	L	0.42	0/1795	0.66	0/2420
12	Z	0.41	0/1795	0.66	0/2420
13	0	0.41	0/1855	0.69	0/2514
13	M	0.40	0/1855	0.68	0/2514
14	1	0.44	0/1541	0.67	0/2087
14	N	0.44	0/1541	0.68	0/2087
All	All	0.40	0/50440	0.65	0/68192

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying

if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
12	L	0	1
12	Z	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
12	L	145	TYR	Sidechain
12	Z	145	TYR	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1915	0	1926	79	0
1	O	1915	0	1926	81	0
2	B	1904	0	1901	110	0
2	P	1904	0	1901	109	0
3	C	1890	0	1900	126	0
3	Q	1890	0	1900	121	0
4	D	1861	0	1836	82	0
4	R	1861	0	1836	83	0
5	E	1795	0	1797	70	0
5	S	1795	0	1797	76	0
6	F	1896	0	1886	83	0
6	T	1896	0	1886	82	0
7	G	1921	0	1910	85	0
7	U	1921	0	1910	95	0
8	H	1684	0	1687	60	0
8	V	1684	0	1687	66	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	I	1581	0	1574	57	0
9	W	1581	0	1574	66	0
10	J	1585	0	1590	83	0
10	X	1585	0	1590	78	0
11	K	1644	0	1594	76	0
11	Y	1644	0	1594	80	0
12	L	1757	0	1711	61	0
12	Z	1757	0	1711	70	0
13	O	1824	0	1832	63	0
13	M	1824	0	1832	66	0
14	1	1512	0	1481	64	0
14	N	1512	0	1481	68	0
15	H	37	0	43	2	0
15	K	37	0	43	2	0
15	V	37	0	43	3	0
15	Y	37	0	43	4	0
16	O	74	0	0	4	0
16	1	64	0	0	3	0
16	A	57	0	0	5	0
16	B	38	0	0	3	0
16	C	42	0	0	8	0
16	D	40	0	0	4	0
16	E	24	0	0	1	0
16	F	46	0	0	5	0
16	G	61	0	0	5	0
16	H	48	0	0	2	0
16	I	66	0	0	3	0
16	J	52	0	0	4	0
16	K	46	0	0	2	0
16	L	60	0	0	2	0
16	M	69	0	0	4	0
16	N	56	0	0	3	0
16	O	33	0	0	3	0
16	P	32	0	0	4	0
16	Q	26	0	0	3	0
16	R	34	0	0	3	0
16	S	20	0	0	1	0
16	T	39	0	0	6	0
16	U	58	0	0	11	0
16	V	47	0	0	6	0
16	W	60	0	0	4	0
16	X	42	0	0	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
16	Y	49	0	0	2	0
16	Z	53	0	0	8	0
All	All	51022	0	49422	2017	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 2017 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
11:Y:10(B):LYS:HD2	11:Y:10(B):LYS:H	1.14	1.10
11:K:10(B):LYS:H	11:K:10(B):LYS:HD2	1.15	1.09
14:N:136:GLY:HA2	14:1:161:GLN:HE21	1.12	1.07
14:N:161:GLN:HE21	14:1:136:GLY:HA2	1.15	1.06
2:P:71:ASN:ND2	2:P:72:ASP:H	1.56	1.04

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/250 (99%)	228 (92%)	15 (6%)	5 (2%)	9	24
1	O	248/250 (99%)	228 (92%)	16 (6%)	4 (2%)	12	30
2	B	242/258 (94%)	220 (91%)	19 (8%)	3 (1%)	16	39
2	P	242/258 (94%)	219 (90%)	19 (8%)	4 (2%)	11	29
3	C	239/254 (94%)	215 (90%)	18 (8%)	6 (2%)	7	18
3	Q	239/254 (94%)	217 (91%)	17 (7%)	5 (2%)	9	23
4	D	240/260 (92%)	215 (90%)	20 (8%)	5 (2%)	9	23
4	R	240/260 (92%)	216 (90%)	18 (8%)	6 (2%)	7	18

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
5	E	231/234 (99%)	210 (91%)	15 (6%)	6 (3%)	7	16
5	S	231/234 (99%)	210 (91%)	16 (7%)	5 (2%)	8	22
6	F	242/287 (84%)	228 (94%)	12 (5%)	2 (1%)	24	51
6	T	242/287 (84%)	226 (93%)	14 (6%)	2 (1%)	24	51
7	G	241/252 (96%)	225 (93%)	13 (5%)	3 (1%)	16	39
7	U	241/252 (96%)	225 (93%)	13 (5%)	3 (1%)	16	39
8	H	220/232 (95%)	203 (92%)	13 (6%)	4 (2%)	11	27
8	V	220/232 (95%)	204 (93%)	12 (6%)	4 (2%)	11	27
9	I	202/205 (98%)	192 (95%)	9 (4%)	1 (0%)	34	63
9	W	202/205 (98%)	190 (94%)	10 (5%)	2 (1%)	19	45
10	J	196/198 (99%)	184 (94%)	10 (5%)	2 (1%)	19	45
10	X	196/198 (99%)	184 (94%)	10 (5%)	2 (1%)	19	45
11	K	210/212 (99%)	192 (91%)	17 (8%)	1 (0%)	34	63
11	Y	210/212 (99%)	193 (92%)	16 (8%)	1 (0%)	34	63
12	L	220/241 (91%)	208 (94%)	11 (5%)	1 (0%)	34	63
12	Z	220/241 (91%)	208 (94%)	11 (5%)	1 (0%)	34	63
13	O	231/266 (87%)	212 (92%)	18 (8%)	1 (0%)	39	69
13	M	231/266 (87%)	213 (92%)	17 (7%)	1 (0%)	39	69
14	I	194/196 (99%)	187 (96%)	7 (4%)	0	100	100
14	N	194/196 (99%)	188 (97%)	6 (3%)	0	100	100
All	All	6312/6690 (94%)	5840 (92%)	392 (6%)	80 (1%)	15	37

5 of 80 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	5	THR
1	A	56	SER
3	C	58	LEU
4	D	12(G)	GLU
6	F	64	ASN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	209/209 (100%)	201 (96%)	8 (4%)	40	71
1	O	209/209 (100%)	200 (96%)	9 (4%)	35	66
2	B	203/216 (94%)	187 (92%)	16 (8%)	15	34
2	P	203/216 (94%)	186 (92%)	17 (8%)	14	30
3	C	213/226 (94%)	200 (94%)	13 (6%)	23	49
3	Q	213/226 (94%)	200 (94%)	13 (6%)	23	49
4	D	198/215 (92%)	188 (95%)	10 (5%)	29	59
4	R	198/215 (92%)	187 (94%)	11 (6%)	26	54
5	E	192/193 (100%)	174 (91%)	18 (9%)	11	25
5	S	192/193 (100%)	173 (90%)	19 (10%)	10	22
6	F	201/238 (84%)	180 (90%)	21 (10%)	9	20
6	T	201/238 (84%)	181 (90%)	20 (10%)	9	22
7	G	207/210 (99%)	192 (93%)	15 (7%)	18	41
7	U	207/210 (99%)	192 (93%)	15 (7%)	18	41
8	H	181/190 (95%)	174 (96%)	7 (4%)	39	70
8	V	181/190 (95%)	174 (96%)	7 (4%)	39	70
9	I	172/173 (99%)	166 (96%)	6 (4%)	43	74
9	W	172/173 (99%)	166 (96%)	6 (4%)	43	74
10	J	175/175 (100%)	165 (94%)	10 (6%)	25	53
10	X	175/175 (100%)	164 (94%)	11 (6%)	22	48
11	K	169/169 (100%)	158 (94%)	11 (6%)	21	46
11	Y	169/169 (100%)	158 (94%)	11 (6%)	21	46
12	L	185/201 (92%)	174 (94%)	11 (6%)	24	51
12	Z	185/201 (92%)	174 (94%)	11 (6%)	24	51
13	0	199/224 (89%)	187 (94%)	12 (6%)	24	50
13	M	199/224 (89%)	188 (94%)	11 (6%)	27	55
14	1	162/162 (100%)	151 (93%)	11 (7%)	20	43
14	N	162/162 (100%)	151 (93%)	11 (7%)	20	43
All	All	5332/5602 (95%)	4991 (94%)	341 (6%)	22	47



5 of 341 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
13	M	149	GLN
2	P	206	THR
12	Z	138	LEU
14	N	36	ARG
1	O	124	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 202 such sidechains are listed below:

Mol	Chain	Res	Type
13	M	89	GLN
3	Q	97	GLN
12	Z	84	GLN
13	M	157	ASN
1	O	191	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
15	GDT	H	500	8	36,37,37	1.97	3 (8%)	41,46,46	1.61	7 (17%)
15	GDT	K	500	11	36,37,37	1.92	2 (5%)	41,46,46	1.64	7 (17%)
15	GDT	V	500	8	36,37,37	2.07	3 (8%)	41,46,46	1.60	7 (17%)
15	GDT	Y	500	11	36,37,37	1.75	3 (8%)	41,46,46	1.70	7 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
15	GDT	H	500	8	-	2/50/50/50	0/0/1/1
15	GDT	K	500	11	-	2/50/50/50	0/0/1/1
15	GDT	V	500	8	-	2/50/50/50	0/0/1/1
15	GDT	Y	500	11	-	2/50/50/50	0/0/1/1

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
15	K	500	GDT	C22-N23	-2.34	1.40	1.46
15	Y	500	GDT	C18-C19	2.05	1.56	1.52
15	Y	500	GDT	C26-C24	2.09	1.52	1.48
15	H	500	GDT	C31-N30	2.19	1.39	1.34
15	V	500	GDT	C31-N30	2.42	1.39	1.34

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
15	Y	500	GDT	O25-C24-C26	-4.57	114.21	123.01
15	K	500	GDT	O25-C24-C26	-4.43	114.47	123.01
15	V	500	GDT	O25-C24-C26	-3.98	115.34	123.01
15	Y	500	GDT	O25-C24-N23	-3.91	117.49	122.53
15	K	500	GDT	O25-C24-N23	-3.65	117.83	122.53

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
15	K	500	GDT	O25-C24-N23-C22
15	K	500	GDT	C26-C24-N23-C22

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
15	H	500	GDT	O25-C24-N23-C22
15	Y	500	GDT	C26-C24-N23-C22
15	H	500	GDT	C26-C24-N23-C22

There are no ring outliers.

4 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
15	H	500	GDT	2	0
15	K	500	GDT	2	0
15	V	500	GDT	3	0
15	Y	500	GDT	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	250/250 (100%)	0.10	6 (2%) 62 62	34, 52, 82, 108	0
1	O	250/250 (100%)	0.27	17 (6%) 20 19	37, 60, 85, 105	0
2	B	244/258 (94%)	0.26	12 (4%) 33 32	34, 58, 92, 117	0
2	P	244/258 (94%)	0.30	16 (6%) 22 20	38, 58, 92, 117	0
3	C	241/254 (94%)	0.38	15 (6%) 24 23	35, 62, 106, 123	0
3	Q	241/254 (94%)	0.69	35 (14%) 3 2	41, 66, 110, 124	0
4	D	242/260 (93%)	0.27	9 (3%) 45 45	39, 58, 90, 123	0
4	R	242/260 (93%)	0.37	17 (7%) 19 17	42, 62, 94, 125	0
5	E	233/234 (99%)	0.27	13 (5%) 28 26	37, 58, 84, 108	0
5	S	233/234 (99%)	0.31	17 (7%) 18 16	32, 58, 88, 107	0
6	F	244/287 (85%)	0.13	10 (4%) 41 41	32, 53, 88, 102	0
6	T	244/287 (85%)	0.14	11 (4%) 37 36	29, 52, 87, 105	0
7	G	243/252 (96%)	0.01	9 (3%) 45 45	29, 48, 78, 113	0
7	U	243/252 (96%)	0.04	6 (2%) 61 61	33, 52, 78, 112	0
8	H	222/232 (95%)	-0.13	3 (1%) 78 77	27, 43, 64, 106	0
8	V	222/232 (95%)	-0.16	3 (1%) 78 77	28, 46, 64, 109	0
9	I	204/205 (99%)	-0.02	3 (1%) 76 76	31, 50, 66, 79	0
9	W	204/205 (99%)	0.04	2 (0%) 84 85	31, 48, 66, 78	0
10	J	198/198 (100%)	-0.00	5 (2%) 61 61	33, 49, 70, 119	0
10	X	198/198 (100%)	0.02	7 (3%) 48 48	36, 49, 69, 124	0
11	K	212/212 (100%)	0.15	6 (2%) 56 57	30, 51, 78, 85	0
11	Y	212/212 (100%)	0.14	8 (3%) 44 44	33, 54, 76, 85	0
12	L	222/241 (92%)	-0.00	6 (2%) 58 58	32, 50, 74, 90	0
12	Z	222/241 (92%)	0.09	4 (1%) 71 72	37, 50, 74, 90	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
13	0	233/266 (87%)	-0.16	1 (0%) 93 94	29, 44, 61, 69	0
13	M	233/266 (87%)	-0.15	2 (0%) 85 86	27, 45, 60, 69	0
14	1	196/196 (100%)	-0.12	3 (1%) 76 76	32, 42, 62, 84	0
14	N	196/196 (100%)	-0.16	2 (1%) 84 85	28, 41, 63, 83	0
All	All	6368/6690 (95%)	0.12	248 (3%) 43 43	27, 52, 85, 125	0

The worst 5 of 248 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	R	12(E)	SER	10.1
4	R	12(F)	GLY	9.9
4	D	12(D)	ALA	9.5
10	X	193	GLN	8.3
3	Q	236	ILE	8.3

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
15	GDT	V	500	37/37	0.88	0.25	3.47	35,40,62,63	0
15	GDT	H	500	37/37	0.89	0.23	1.89	31,35,53,53	0
15	GDT	K	500	37/37	0.94	0.19	1.07	31,41,46,47	0
15	GDT	Y	500	37/37	0.93	0.19	0.43	39,44,55,56	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.