



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 06:34 PM GMT

PDB ID : 1BEV
Title : BOVINE ENTEROVIRUS VG-5-27
Authors : Smyth, M.; Tate, J.; Lyons, C.; Hoey, E.; Martin, S.; Stuart, D.
Deposited on : 1996-04-03
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	1	281	 71% 20% 5% 5%
2	2	248	 79% 17% • •
3	3	242	 69% 24% 6% •
4	4	68	 40% 15% • 41%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6315 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BOVINE ENTEROVIRUS COAT PROTEINS VP1 TO VP4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	1	268	Total	C	N	O	S	0	0	0
			2099	1323	361	404	11			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
1	15	ALA	VAL	CONFLICT	UNP P12915
1	24	SER	THR	CONFLICT	UNP P12915
1	94	HIS	ASN	CONFLICT	UNP P12915
1	237	TYR	CYS	CONFLICT	UNP P12915

- Molecule 2 is a protein called BOVINE ENTEROVIRUS COAT PROTEINS VP1 TO VP4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	2	244	Total	C	N	O	S	0	0	0
			1890	1201	325	357	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
2	62	ARG	ALA	CONFLICT	UNP P12915
2	101	LEU	ILE	CONFLICT	UNP P12915

- Molecule 3 is a protein called BOVINE ENTEROVIRUS COAT PROTEINS VP1 TO VP4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	3	242	Total	C	N	O	S	0	0	0
			1871	1195	306	358	12			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
3	32	PRO	LEU	CONFLICT	UNP P12915
3	154	ILE	VAL	CONFLICT	UNP P12915


- Molecule 4 is a protein called BOVINE ENTEROVIRUS COAT PROTEINS VP1 TO VP4.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	4	40	Total	C	N	O	0	0	0
			299	185	52	62			

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
4	16	GLN	GLY	CONFLICT	UNP P12915

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	2	1	Total	O	S	0	0
			5	4	1		
5	1	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is MYRISTIC ACID (three-letter code: MYR) (formula: C₁₄H₂₈O₂).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	1	1	Total	C	O	0
			16	14	2	

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	1	49	Total	O	0	0
			49	49		
7	2	51	Total	O	0	0
			51	51		
7	3	30	Total	O	0	0
			30	30		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: BOVINE ENTEROVIRUS COAT PROTEINS VP1 TO VP4



• Molecule 2: BOVINE ENTEROVIRUS COAT PROTEINS VP1 TO VP4



• Molecule 3: BOVINE ENTEROVIRUS COAT PROTEINS VP1 TO VP4



• Molecule 4: BOVINE ENTEROVIRUS COAT PROTEINS VP1 TO VP4

Chain 4: 40% 15% • 41%

GLY	ALA	GLN	LEU	SER	ARG	ASN	THR	ALA	GLY	SER	HIS	THR	THR	GLN	THR	TYR	ALA	THR	GLY	GLY	S23	T24	I25	N28	N29	I30	A37	Q41	R42	K43	Q44	D45	F46	T47	Q48	D59	V60	I61	K62	GLU	THR	ALA	VAL	PRO	LEU	LYS
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	388.20 Å 392.60 Å 360.70 Å 90.00° 112.70° 90.00°	Depositor
Resolution (Å)	25.00 – 3.00	Depositor
% Data completeness (in resolution range)	42.5 (25.00-3.00)	Depositor
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.225 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	6315	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MYR, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	1	0.48	0/2158	0.78	1/2939 (0.0%)
2	2	0.46	1/1944 (0.1%)	0.76	1/2664 (0.0%)
3	3	0.48	0/1919	0.79	2/2618 (0.1%)
4	4	0.50	0/305	0.68	0/416
All	All	0.48	1/6326 (0.0%)	0.77	4/8637 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	2	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	2	61	ASP	CB-CG	5.02	1.62	1.51

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	3	173	SER	N-CA-C	6.50	128.54	111.00
1	1	98	ASP	CB-CA-C	-6.05	98.30	110.40
2	2	181	ASN	N-CA-CB	-5.58	100.55	110.60
3	3	63	GLY	N-CA-C	5.15	125.98	113.10

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	2	216	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1	2099	0	1997	57	0
2	2	1890	0	1816	35	0
3	3	1871	0	1833	71	0
4	4	299	0	261	5	0
5	1	5	0	0	0	0
5	2	5	0	0	0	0
6	1	16	0	27	2	0
7	1	49	0	0	3	0
7	2	51	0	0	6	0
7	3	30	0	0	3	0
All	All	6315	0	5934	145	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 145 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:3:63:GLY:O	3:3:64:VAL:HG12	1.32	1.27
2:2:220:THR:O	2:2:221:THR:HG22	1.44	1.18
3:3:173:SER:O	3:3:174:HIS:HB2	1.52	1.07
2:2:182:SER:OG	7:2:956:HOH:O	1.81	0.97
1:1:52:GLU:HG3	1:1:57:THR:HG21	1.48	0.92

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	1	266/281 (95%)	252 (95%)	11 (4%)	3 (1%)	17	58
2	2	242/248 (98%)	220 (91%)	19 (8%)	3 (1%)	16	56
3	3	240/242 (99%)	225 (94%)	10 (4%)	5 (2%)	9	40
4	4	38/68 (56%)	32 (84%)	2 (5%)	4 (10%)	1	3
All	All	786/839 (94%)	729 (93%)	42 (5%)	15 (2%)	10	43

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	2	6	ALA
3	3	63	GLY
3	3	90	GLY
4	4	24	THR
4	4	30	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	1	228/239 (95%)	200 (88%)	28 (12%)	6	25
2	2	198/205 (97%)	189 (96%)	9 (4%)	34	74
3	3	206/206 (100%)	184 (89%)	22 (11%)	8	31
4	4	30/55 (54%)	22 (73%)	8 (27%)	0	3
All	All	662/705 (94%)	595 (90%)	67 (10%)	9	34

5 of 67 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	2	62	ARG
3	3	12	GLN
4	4	41	GLN
2	2	86	LEU
2	2	102	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
2	2	130	HIS
2	2	173	GLN
4	4	28	ASN
2	2	131	GLN
2	2	153	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	MYR	1	900	-	12,15,15	0.61	0	12,15,15	0.74	0
5	SO4	1	951	-	4,4,4	0.45	0	6,6,6	0.35	0
5	SO4	2	950	-	4,4,4	0.31	0	6,6,6	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	MYR	1	900	-	-	0/11/13/13	0/0/0/0
5	SO4	1	951	-	-	0/0/0/0	0/0/0/0
5	SO4	2	950	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	1	900	MYR	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.