



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:35 AM GMT

PDB ID : 3BF5  
Title : Crystal structure of putative ribokinase (10640157) from *Thermoplasma acidophilum* at 1.91 Å resolution  
Authors : Joint Center for Structural Genomics (JCSG)  
Deposited on : 2007-11-20  
Resolution : 1.91 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	5832 (1.94-1.90)
Clashscore	102246	6540 (1.94-1.90)
Ramachandran outliers	100387	6464 (1.94-1.90)
Sidechain outliers	100360	6465 (1.94-1.90)
RSRZ outliers	91569	5846 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	<div> <div>16%</div> <div>86%</div> <div>7%</div> <div>7%</div> </div>
1	B	306	<div> <div>29%</div> <div>82%</div> <div>9%</div> <div>9%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PGO	A	288	X	-	-	X
3	PGO	A	289	X	-	-	-
3	PGO	A	290	X	-	-	-
3	PGO	B	289	X	-	-	X
3	PGO	B	290	X	-	-	-
3	PGO	B	291	X	-	-	X
3	PGO	B	292	X	-	-	-

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4673 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribokinase related protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	284	Total	C	N	O	S	Se	0	10	0
			2266	1459	369	421	1	16			
1	B	279	Total	C	N	O	S	Se	0	4	0
			2153	1386	350	403	1	13			

There are 44 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	MSE	-	LEADER SEQUENCE	UNP Q9HJT3
A	-17	GLY	-	LEADER SEQUENCE	UNP Q9HJT3
A	-16	SER	-	LEADER SEQUENCE	UNP Q9HJT3
A	-15	ASP	-	LEADER SEQUENCE	UNP Q9HJT3
A	-14	LYS	-	LEADER SEQUENCE	UNP Q9HJT3
A	-13	ILE	-	LEADER SEQUENCE	UNP Q9HJT3
A	-12	HIS	-	LEADER SEQUENCE	UNP Q9HJT3
A	-11	HIS	-	LEADER SEQUENCE	UNP Q9HJT3
A	-10	HIS	-	LEADER SEQUENCE	UNP Q9HJT3
A	-9	HIS	-	LEADER SEQUENCE	UNP Q9HJT3
A	-8	HIS	-	LEADER SEQUENCE	UNP Q9HJT3
A	-7	HIS	-	LEADER SEQUENCE	UNP Q9HJT3
A	-6	GLU	-	LEADER SEQUENCE	UNP Q9HJT3
A	-5	ASN	-	LEADER SEQUENCE	UNP Q9HJT3
A	-4	LEU	-	LEADER SEQUENCE	UNP Q9HJT3
A	-3	TYR	-	LEADER SEQUENCE	UNP Q9HJT3
A	-2	PHE	-	LEADER SEQUENCE	UNP Q9HJT3
A	-1	GLN	-	LEADER SEQUENCE	UNP Q9HJT3
A	0	GLY	-	LEADER SEQUENCE	UNP Q9HJT3
A	112	ALA	GLU	ENGINEERED	UNP Q9HJT3
A	113	ALA	LYS	ENGINEERED	UNP Q9HJT3
A	115	ALA	LYS	ENGINEERED	UNP Q9HJT3
B	-18	MSE	-	LEADER SEQUENCE	UNP Q9HJT3
B	-17	GLY	-	LEADER SEQUENCE	UNP Q9HJT3
B	-16	SER	-	LEADER SEQUENCE	UNP Q9HJT3

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-15	ASP	-	LEADER SEQUENCE	UNP Q9HJT3
B	-14	LYS	-	LEADER SEQUENCE	UNP Q9HJT3
B	-13	ILE	-	LEADER SEQUENCE	UNP Q9HJT3
B	-12	HIS	-	LEADER SEQUENCE	UNP Q9HJT3
B	-11	HIS	-	LEADER SEQUENCE	UNP Q9HJT3
B	-10	HIS	-	LEADER SEQUENCE	UNP Q9HJT3
B	-9	HIS	-	LEADER SEQUENCE	UNP Q9HJT3
B	-8	HIS	-	LEADER SEQUENCE	UNP Q9HJT3
B	-7	HIS	-	LEADER SEQUENCE	UNP Q9HJT3
B	-6	GLU	-	LEADER SEQUENCE	UNP Q9HJT3
B	-5	ASN	-	LEADER SEQUENCE	UNP Q9HJT3
B	-4	LEU	-	LEADER SEQUENCE	UNP Q9HJT3
B	-3	TYR	-	LEADER SEQUENCE	UNP Q9HJT3
B	-2	PHE	-	LEADER SEQUENCE	UNP Q9HJT3
B	-1	GLN	-	LEADER SEQUENCE	UNP Q9HJT3
B	0	GLY	-	LEADER SEQUENCE	UNP Q9HJT3
B	112	ALA	GLU	ENGINEERED	UNP Q9HJT3
B	113	ALA	LYS	ENGINEERED	UNP Q9HJT3
B	115	ALA	LYS	ENGINEERED	UNP Q9HJT3

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is S-1,2-PROPANEDIOL (three-letter code: PGO) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			5	3	2		
3	B	1	Total	C	O	0	0
			5	3	2		
3	B	1	Total	C	O	0	0
			5	3	2		
3	A	1	Total	C	O	0	0
			5	3	2		
3	B	1	Total	C	O	0	0
			5	3	2		
3	A	1	Total	C	O	0	0
			5	3	2		
3	B	1	Total	C	O	0	0
			5	3	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	144	Total	O	0	0
			144	144		
4	B	70	Total	O	0	0
			70	70		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ribokinase related protein



- Molecule 1: Ribokinase related protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.45Å 59.18Å 177.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.18 – 1.91 29.19 – 1.91	Depositor EDS
% Data completeness (in resolution range)	99.1 (29.18-1.91) 99.2 (29.19-1.91)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.86 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.202 , 0.245 0.206 , 0.247	Depositor DCC
$R_{free}$ test set	2364 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	30.6	Xtriage
Anisotropy	0.406	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 53.9	EDS
Estimated twinning fraction	0.048 for k,h,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 46708 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4673	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGO, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.65	0/2337	0.86	4/3129 (0.1%)
1	B	0.62	0/2204	0.75	0/2959
All	All	0.64	0/4541	0.81	4/6088 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	2	ARG	NE-CZ-NH1	11.00	125.80	120.30
1	A	2	ARG	NE-CZ-NH2	-6.30	117.15	120.30
1	A	241	ARG	NE-CZ-NH1	5.81	123.21	120.30
1	A	240	ARG	NE-CZ-NH1	5.54	123.07	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2266	0	2199	9	0
1	B	2153	0	2040	24	0
2	B	5	0	0	0	0
3	A	15	0	24	0	0
3	B	20	0	32	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	144	0	0	0	0
4	B	70	0	0	0	0
All	All	4673	0	4295	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:179:VAL:HG12	1:B:183[A]:MSE:HE2	1.13	1.10
1:B:179:VAL:HG12	1:B:183[A]:MSE:CE	2.00	0.90
1:A:40:ALA:HB1	1:A:56:LEU:HD11	1.68	0.75
1:B:152:ILE:HG13	1:B:183[A]:MSE:HE1	1.70	0.74
1:B:118:LEU:HB2	1:B:136:MSE:HE3	1.73	0.70
1:B:160:LEU:HD11	1:B:183[A]:MSE:CE	2.30	0.62
1:B:206:MSE:SE	1:B:243[A]:ILE:HG21	2.50	0.61
1:B:160:LEU:HD11	1:B:183[A]:MSE:HE3	1.82	0.61
1:B:1:MSE:HE3	1:B:123:GLU:HB2	1.84	0.60
1:B:7:PHE:CZ	1:B:136:MSE:HE1	2.39	0.57
1:A:153:HIS:HA	1:A:183[A]:MSE:HE1	1.86	0.57
1:A:1:MSE:HE3	1:A:123:GLU:HB2	1.87	0.55
1:B:243[B]:ILE:HD12	3:B:292:PGO:H33	1.88	0.55
1:B:1:MSE:HE3	1:B:123:GLU:CB	2.36	0.55
1:B:40:ALA:HB1	1:B:56:LEU:HD11	1.90	0.54
1:B:118:LEU:HD22	1:B:136:MSE:CE	2.38	0.53
1:B:160:LEU:CD1	1:B:183[A]:MSE:HE3	2.39	0.53
1:B:62:MSE:HE1	3:B:290:PGO:H12	1.90	0.52
1:B:175:HIS:O	1:B:179:VAL:HG23	2.10	0.51
1:A:206:MSE:SE	1:A:243[B]:ILE:HG21	2.61	0.50
1:B:97:THR:HG22	1:B:99:GLY:H	1.76	0.50
1:B:118:LEU:HD22	1:B:136:MSE:HE3	1.94	0.49
1:A:157:LYS:HG3	1:A:184:THR:HA	1.95	0.47
1:B:206:MSE:SE	3:B:292:PGO:C3	3.14	0.46
1:B:232:GLY:HA2	1:B:253:ILE:HD12	2.00	0.44
1:B:183[B]:MSE:HE2	1:B:183[B]:MSE:HA	1.99	0.43
1:A:282[B]:MSE:HE3	1:A:283:PHE:O	2.19	0.43
1:B:127:PHE:CZ	1:B:136:MSE:HE2	2.54	0.43
1:A:2:ARG:HG3	1:A:2:ARG:HH11	1.85	0.42
1:A:22:PRO:HG2	1:B:103:VAL:HG13	2.03	0.41
1:A:128:SER:HA	1:A:147:ASP:O	2.21	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:44:ALA:HB1	1:B:77:ILE:HD13	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	292/306 (95%)	282 (97%)	10 (3%)	0	100	100
1	B	279/306 (91%)	268 (96%)	11 (4%)	0	100	100
All	All	571/612 (93%)	550 (96%)	21 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/249 (96%)	234 (98%)	5 (2%)	61	54
1	B	217/249 (87%)	213 (98%)	4 (2%)	66	60
All	All	456/498 (92%)	447 (98%)	9 (2%)	63	56

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	ARG
1	A	53	PRO
1	A	68	TYR
1	A	143	LYS
1	A	163	PHE
1	B	68	TYR
1	B	87	ASP
1	B	150	GLN
1	B	163	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	264	ASN
1	A	278	ASN
1	B	28	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	PGO	A	288	-	4,4,4	0.51	0	2,4,4	0.76	0
3	PGO	A	289	-	4,4,4	0.57	0	2,4,4	0.66	0
3	PGO	A	290	-	4,4,4	0.39	0	2,4,4	0.26	0
2	PO4	B	288	-	4,4,4	0.65	0	6,6,6	0.27	0
3	PGO	B	289	-	4,4,4	0.72	0	2,4,4	0.51	0
3	PGO	B	290	-	4,4,4	0.58	0	2,4,4	0.11	0
3	PGO	B	291	-	4,4,4	0.76	0	2,4,4	0.73	0
3	PGO	B	292	-	4,4,4	0.52	0	2,4,4	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PGO	A	288	-	1/1/1/1	0/2/2/2	0/0/0/0
3	PGO	A	289	-	1/1/1/1	0/2/2/2	0/0/0/0
3	PGO	A	290	-	1/1/1/1	0/2/2/2	0/0/0/0
2	PO4	B	288	-	-	0/0/0/0	0/0/0/0
3	PGO	B	289	-	1/1/1/1	0/2/2/2	0/0/0/0
3	PGO	B	290	-	1/1/1/1	0/2/2/2	0/0/0/0
3	PGO	B	291	-	1/1/1/1	0/2/2/2	0/0/0/0
3	PGO	B	292	-	1/1/1/1	0/2/2/2	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

All (7) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	288	PGO	C2
3	A	290	PGO	C2
3	B	291	PGO	C2
3	A	289	PGO	C2
3	B	292	PGO	C2
3	B	289	PGO	C2
3	B	290	PGO	C2

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	290	PGO	1	0
3	B	292	PGO	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	271/306 (88%)	1.20	48 (17%) <b>2</b> <b>2</b>	32, 40, 58, 72	0
1	B	266/306 (86%)	1.79	90 (33%) <b>0</b> <b>0</b>	32, 40, 54, 63	0
All	All	537/612 (87%)	1.49	138 (25%) <b>1</b> <b>1</b>	32, 40, 57, 72	0

All (138) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	217	PRO	8.9
1	B	223	VAL	7.9
1	A	0	GLY	7.6
1	B	216	ILE	6.8
1	B	119	ALA	6.4
1	B	213	PHE	6.4
1	A	283	PHE	6.2
1	B	268	ASN	6.1
1	B	218	SER	6.0
1	B	219	SER	5.9
1	B	263	GLU	5.4
1	B	74	SER	5.2
1	A	100	LYS	5.1
1	B	221	ASP	5.0
1	A	145	ILE	4.8
1	A	209	LYS	4.8
1	A	171[A]	ILE	4.8
1	B	252	ILE	4.7
1	A	223	VAL	4.5
1	B	281	ARG	4.4
1	B	267	LEU	4.4
1	B	212	ASP	4.4
1	B	220	GLY	4.2
1	B	202	SER	4.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	253	ILE	4.1
1	A	119	ALA	4.0
1	B	262	ILE	4.0
1	B	279	TYR	3.9
1	B	50	PHE	3.7
1	A	264	ASN	3.7
1	A	40	ALA	3.7
1	A	47	ALA	3.7
1	A	153	HIS	3.6
1	B	256	HIS	3.6
1	B	222	THR	3.6
1	B	161	LYS	3.5
1	B	155	TYR	3.4
1	B	16	ILE	3.4
1	B	280	ARG	3.4
1	B	118	LEU	3.4
1	B	158	ASP	3.4
1	B	211	TYR	3.3
1	B	277	GLU	3.3
1	A	231	ALA	3.3
1	B	231	ALA	3.3
1	A	82	VAL	3.2
1	B	153	HIS	3.2
1	B	139	SER	3.2
1	B	43	PHE	3.1
1	A	6	TYR	3.1
1	A	101	LYS	3.1
1	A	194	ILE	3.0
1	B	121	GLU	3.0
1	A	243[A]	ILE	3.0
1	B	46	VAL	3.0
1	B	214	PRO	2.9
1	B	160	LEU	2.9
1	A	43	PHE	2.9
1	A	39	THR	2.8
1	A	56	LEU	2.8
1	A	229	PHE	2.8
1	A	281	ARG	2.8
1	A	117	GLN	2.8
1	B	200	ARG	2.8
1	B	97	THR	2.8
1	B	145	ILE	2.7

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	48	GLN	2.7
1	A	272	LEU	2.7
1	B	199	GLU	2.7
1	A	59	ALA	2.7
1	A	126	HIS	2.7
1	B	233	LEU	2.7
1	B	276	THR	2.6
1	B	87	ASP	2.6
1	B	85	PHE	2.6
1	B	82	VAL	2.6
1	B	254	ALA	2.6
1	A	99	GLY	2.6
1	B	156	SER	2.6
1	B	194	ILE	2.6
1	B	243[A]	ILE	2.6
1	B	257	VAL	2.6
1	A	132	ASN	2.6
1	B	201	GLY	2.6
1	B	258	ILE	2.6
1	B	272	LEU	2.6
1	B	57	TYR	2.6
1	B	236	ALA	2.5
1	B	52	ILE	2.5
1	B	270	GLU	2.5
1	B	54	PHE	2.5
1	B	234	TYR	2.5
1	B	47	ALA	2.5
1	A	144	ILE	2.4
1	A	266	SER	2.4
1	A	149[A]	SER	2.4
1	B	274	ARG	2.4
1	A	234	TYR	2.4
1	A	265	PHE	2.4
1	B	140	ILE	2.3
1	A	10	LEU	2.3
1	A	41	GLY	2.3
1	A	77	ILE	2.3
1	B	144	ILE	2.3
1	A	135	ASP	2.3
1	A	22	PRO	2.3
1	B	260	ASP	2.3
1	B	261	GLY	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	4	LEU	2.2
1	B	150	GLN	2.2
1	B	15	LEU	2.2
1	B	141	ARG	2.2
1	B	239	ASN	2.2
1	B	72	ILE	2.2
1	B	229	PHE	2.2
1	B	182	GLU	2.2
1	A	188	SER	2.2
1	B	2	ARG	2.2
1	A	131	PRO	2.2
1	B	195	VAL	2.2
1	B	215	ALA	2.1
1	B	166	ILE	2.1
1	A	233	LEU	2.1
1	A	246	GLY	2.1
1	B	171	ILE	2.1
1	B	10	LEU	2.1
1	A	158	ASP	2.1
1	B	188	SER	2.1
1	B	198	GLY	2.1
1	B	126	HIS	2.1
1	B	53	PRO	2.1
1	B	178	ARG	2.0
1	A	237	LEU	2.0
1	A	267	LEU	2.0
1	B	133	TYR	2.0
1	B	116	PRO	2.0
1	A	12	ILE	2.0
1	A	19	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	PGO	B	291	5/5	0.74	0.61	14.43	65,68,69,70	0
3	PGO	A	288	5/5	0.82	0.29	5.44	43,46,48,49	0
3	PGO	B	289	5/5	0.61	0.32	4.60	58,58,59,60	0
3	PGO	B	290	5/5	0.53	0.30	1.56	56,56,59,62	0
3	PGO	A	290	5/5	0.89	0.18	0.62	56,61,67,69	0
3	PGO	A	289	5/5	0.70	0.22	0.50	68,71,74,76	0
2	PO4	B	288	5/5	0.95	0.16	-0.40	43,52,56,57	0
3	PGO	B	292	5/5	0.58	0.41	-	62,68,73,79	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.