



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:37 AM GMT

PDB ID : 3BHT
Title : Structure of phosphorylated Thr160 CDK2/cyclin A in complex with the inhibitor meriolin 3
Authors : Echalier, A.; Bettayeb, K.; Ferandin, Y.; Lozach, O.; Clement, M.; Valette, A.; Liger, F.; Marquet, B.; Morris, J.C.; Endicott, J.A.; Joseph, B.; Meijer, L.
Deposited on : 2007-11-29
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	300	<div> <div>10%</div> <div>84%</div> <div>13%</div> <div>...</div> </div>
1	C	300	<div> <div>12%</div> <div>77%</div> <div>11%</div> <div>10%</div> </div>
2	B	262	<div> <div>6%</div> <div>90%</div> <div>9%</div> </div>
2	D	262	<div> <div>12%</div> <div>88%</div> <div>10%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MG	D	2	-	-	-	X
5	SGM	B	4	-	-	-	X
5	SGM	D	3	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9983 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cell division protein kinase 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	298	Total	C	N	O	P	S	0	4	0
			2416	1568	412	426	1	9			
1	C	271	Total	C	N	O	P	S	0	2	0
			2185	1417	373	387	1	7			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	EXPRESSION TAG	UNP P24941
A	0	SER	-	EXPRESSION TAG	UNP P24941
C	-1	GLY	-	EXPRESSION TAG	UNP P24941
C	0	SER	-	EXPRESSION TAG	UNP P24941


- Molecule 2 is a protein called Cyclin-A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	262	Total	C	N	O	S	0	3	0
			2130	1379	346	395	10			
2	D	262	Total	C	N	O	S	0	1	0
			2118	1371	344	393	10			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	D	1	Total	Mg	0	0
			1	1		

- Molecule 4 is 4-(4-METHOXY-1H-PYRROLO[2,3-B]PYRIDIN-3-YL)PYRIMIDIN-2-AMINE (three-letter code: MFR) (formula: C₁₂H₁₁N₅O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			18	12	5	1		
4	C	1	Total	C	N	O	0	0
			18	12	5	1		

- Molecule 5 is MONOTHIOGLYCEROL (three-letter code: SGM) (formula: $C_3H_8O_2S$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	O	S	0	0
			6	3	2	1		
5	B	1	Total	C	O	S	0	0
			6	3	2	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	D	1	Total	C	O	S	0	0
			6	3	2	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	345	Total	O	0	0
			345	345		
6	B	292	Total	O	0	0
			292	292		
6	C	227	Total	O	0	0
			227	227		
6	D	214	Total	O	0	0
			214	214		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cell division protein kinase 2




- Molecule 1: Cell division protein kinase 2




- Molecule 2: Cyclin-A2



- Molecule 2: Cyclin-A2





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	74.13Å 133.77Å 147.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.00 19.87 – 2.00	Depositor EDS
% Data completeness (in resolution range)	95.5 (20.00-2.00) 95.5 (19.87-2.00)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.71 (at 2.01Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.181 , 0.220 0.181 , 0.215	Depositor DCC
R_{free} test set	4777 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	25.7	Xtriage
Anisotropy	0.229	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 39.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 95265 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9983	wwPDB-VP
Average B, all atoms (Å ²)	9.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, MG, SGM, MFR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	0/2477	0.74	2/3357 (0.1%)
1	C	0.54	0/2229	0.62	0/3016
2	B	0.55	0/2190	0.58	0/2978
2	D	0.51	0/2171	0.67	2/2952 (0.1%)
All	All	0.59	0/9067	0.66	4/12303 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	174	GLU	N-CA-C	10.17	138.45	111.00
1	A	217	ARG	NE-CZ-NH2	-8.40	116.10	120.30
2	D	175	VAL	N-CA-C	-6.80	92.63	111.00
1	A	163	VAL	N-CA-C	-6.39	93.75	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	14	THR	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2416	0	2471	52	1
1	C	2185	0	2250	41	1
2	B	2130	0	2146	19	0
2	D	2118	0	2136	34	0
3	B	1	0	0	0	0
3	D	1	0	0	0	0
4	A	18	0	11	0	0
4	C	18	0	11	0	0
5	B	12	0	15	3	0
5	D	6	0	8	3	0
6	A	345	0	0	9	0
6	B	292	0	0	6	0
6	C	227	0	0	5	0
6	D	214	0	0	8	0
All	All	9983	0	9048	128	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (128) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:177[A]:CYS:SG	1:A:233:MET:HE3	1.45	1.54
2:B:193:CYS:SG	5:B:4:SGM:S1	2.31	1.27
2:D:193:CYS:SG	5:D:3:SGM:S1	2.38	1.22
2:B:171:SER:OG	6:B:595:HOH:O	1.68	1.12
1:C:272:ASN:OD1	2:D:171:SER:HB3	1.47	1.11
1:A:177[A]:CYS:SG	1:A:233:MET:CE	2.38	1.11
1:C:15:TYR:HB2	1:C:16:GLY:CA	1.82	1.08
1:C:15:TYR:HB2	1:C:16:GLY:HA3	1.32	1.06
1:A:15:TYR:CD2	1:A:16:GLY:HA3	1.90	1.05
1:A:272:ASN:OD1	2:B:171:SER:HB2	1.55	1.05
2:D:432:VAL:HG23	6:D:580:HOH:O	1.59	0.99
1:A:2:GLU:HG3	2:D:293:ARG:NH2	1.82	0.94
1:A:2:GLU:CG	2:D:293:ARG:NH2	2.31	0.93

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:15:TYR:CB	1:C:16:GLY:HA3	1.91	0.93
1:C:36:ARG:HH11	1:C:36:ARG:HG3	1.33	0.91
1:A:2:GLU:HG2	1:C:73:GLU:OE2	1.70	0.91
1:A:161:HIS:HD2	6:A:353:HOH:O	1.55	0.90
2:D:171:SER:HB2	6:D:697:HOH:O	1.71	0.89
2:B:327:CYS:HG	5:B:5:SGM:HS1	1.14	0.89
1:C:71:HIS:HD2	2:D:296:HIS:HE1	1.14	0.89
1:A:160:TPO:HG23	1:A:162:GLU:HG2	1.59	0.84
1:A:15:TYR:CG	1:A:16:GLY:HA3	2.14	0.83
1:C:71:HIS:HD2	2:D:296:HIS:CE1	1.97	0.83
1:A:268:HIS:CD2	6:A:589:HOH:O	2.33	0.81
1:A:2:GLU:HG2	2:D:293:ARG:NH2	1.99	0.78
1:A:15:TYR:CD2	1:A:16:GLY:CA	2.65	0.77
1:A:177[A]:CYS:HG	1:A:233:MET:HE3	1.50	0.77
1:C:71:HIS:CD2	2:D:296:HIS:HE1	2.01	0.76
2:B:183:HIS:HB2	2:B:317:GLN:HE22	1.51	0.74
2:D:344:ALA:HB1	2:D:348:LEU:HD22	1.68	0.74
1:A:60:HIS:HD2	1:A:62:ASN:H	1.33	0.74
1:C:36:ARG:CG	1:C:36:ARG:HH11	2.00	0.73
1:A:95:ALA:O	1:A:96:LEU:HB2	1.87	0.73
1:A:161:HIS:O	1:A:161:HIS:CG	2.42	0.71
2:B:327:CYS:SG	5:B:5:SGM:S1	2.82	0.71
1:A:60:HIS:CD2	1:A:62:ASN:H	2.09	0.70
1:A:160:TPO:CG2	1:A:162:GLU:HG2	2.21	0.70
1:A:59[A]:ASN:ND2	6:A:333:HOH:O	2.26	0.69
2:D:305:ASP:HB3	5:D:3:SGM:H12	1.76	0.67
2:B:344:ALA:HB1	2:B:348:LEU:HD22	1.77	0.66
1:A:268:HIS:NE2	6:A:589:HOH:O	2.28	0.66
1:A:161:HIS:O	1:A:162:GLU:C	2.36	0.64
1:C:15:TYR:CB	1:C:16:GLY:CA	2.60	0.64
1:A:272:ASN:CG	2:B:171:SER:HB2	2.17	0.63
1:A:15:TYR:HD2	1:A:16:GLY:CA	2.12	0.62
2:B:210:MET:HE1	2:B:250:ARG:CB	2.29	0.62
1:A:15:TYR:CB	1:A:16:GLY:HA3	2.29	0.62
1:C:268:HIS:HD2	1:C:270:ASP:H	1.46	0.62
1:C:15:TYR:HB2	1:C:16:GLY:HA2	1.79	0.62
1:A:88:LYS:HE2	6:A:593:HOH:O	2.00	0.62
2:D:285:THR:HG22	2:D:286:TYR:CD1	2.35	0.62
1:C:52:ILE:HD11	1:C:78:LEU:HD21	1.82	0.60
1:C:71:HIS:CD2	2:D:296:HIS:CE1	2.84	0.60
1:A:2:GLU:CG	2:D:293:ARG:HH22	2.14	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:161:HIS:CD2	1:A:161:HIS:O	2.54	0.59
2:B:194:LYS:HD3	2:B:351:LEU:HD23	1.83	0.59
1:C:272:ASN:OD1	2:D:171:SER:CB	2.38	0.59
1:A:162:GLU:HG3	1:A:163:VAL:HG12	1.83	0.59
1:C:41:THR:HG22	1:C:42:GLU:N	2.16	0.59
1:C:60:HIS:CD2	1:C:62:ASN:H	2.21	0.58
1:C:59:ASN:ND2	6:C:458:HOH:O	2.37	0.58
1:C:278:LYS:HE3	2:D:177:ASP:O	2.04	0.58
2:D:287:THR:HB	6:D:710:HOH:O	2.04	0.57
2:B:210:MET:HE1	2:B:250:ARG:HB2	1.85	0.57
2:B:395:HIS:HE1	2:B:427:PRO:O	1.87	0.56
2:B:183:HIS:HD2	6:B:553:HOH:O	1.87	0.56
1:C:60:HIS:HD2	1:C:62:ASN:H	1.51	0.56
1:A:0:SER:O	1:A:1:MET:CB	2.54	0.55
1:A:0:SER:O	1:A:1:MET:HB2	2.07	0.54
1:A:161:HIS:O	1:A:162:GLU:O	2.27	0.53
1:C:218:THR:HG23	1:C:251:VAL:HG21	1.91	0.52
1:A:2:GLU:HG2	2:D:293:ARG:HH21	1.75	0.52
1:C:161:HIS:O	1:C:162:GLU:C	2.46	0.52
1:A:119:HIS:HD2	6:B:543:HOH:O	1.91	0.52
1:C:15:TYR:HB3	6:C:353:HOH:O	2.09	0.52
1:C:161:HIS:O	1:C:162:GLU:O	2.28	0.51
1:A:2:GLU:CG	2:D:293:ARG:HH21	2.22	0.51
1:C:161:HIS:HD2	6:C:372:HOH:O	1.94	0.50
2:B:233:HIS:HE1	6:B:447:HOH:O	1.94	0.50
1:C:216:PHE:HB3	1:C:221:THR:HG23	1.94	0.50
1:C:71:HIS:NE2	2:D:304:PHE:HE1	2.10	0.49
2:D:233:HIS:HE1	6:D:551:HOH:O	1.94	0.49
1:C:15:TYR:CG	1:C:16:GLY:HA3	2.47	0.49
1:A:260:ARG:HD3	6:A:334:HOH:O	2.11	0.49
2:B:225:TYR:HE2	2:B:281:ILE:HG21	1.78	0.49
2:B:179[A]:HIS:CE1	6:B:492:HOH:O	2.66	0.48
1:C:161:HIS:CE1	6:C:490:HOH:O	2.67	0.48
2:B:289:LYS:HE3	2:B:293:ARG:HE	1.78	0.48
1:C:83:LEU:HD23	1:C:136:ASN:HB3	1.95	0.47
1:A:223:ASP:H	1:A:226:VAL:HG12	1.79	0.47
1:A:162:GLU:HG3	1:A:163:VAL:CG1	2.45	0.46
1:C:71:HIS:HE1	6:D:663:HOH:O	1.99	0.46
2:D:285:THR:HG22	2:D:286:TYR:CG	2.50	0.46
1:C:119:HIS:HD2	6:D:535:HOH:O	1.98	0.46
1:A:60:HIS:HE1	6:A:441:HOH:O	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:183:HIS:HB2	2:D:317:GLN:HE22	1.81	0.45
1:A:88:LYS:HA	1:A:91:MET:CE	2.47	0.45
2:B:300:LYS:HE3	6:B:614:HOH:O	2.16	0.45
1:A:268:HIS:HD2	1:A:270:ASP:H	1.65	0.44
2:D:285:THR:HG22	2:D:286:TYR:CE1	2.52	0.44
6:C:439:HOH:O	2:D:296:HIS:HD2	2.01	0.44
1:A:98:GLY:HA2	1:A:199:ARG:NE	2.33	0.43
2:D:196:LYS:HG2	2:D:199:TYR:HB3	2.01	0.43
1:A:15:TYR:CB	1:A:16:GLY:CA	2.96	0.43
1:A:161:HIS:CD2	6:A:353:HOH:O	2.44	0.43
1:C:88:LYS:HG3	1:C:131:GLN:NE2	2.34	0.43
2:B:194:LYS:HD3	2:B:351:LEU:CD2	2.48	0.43
1:C:36:ARG:HG3	1:C:36:ARG:NH1	2.15	0.43
1:A:65:LYS:HD2	6:A:333:HOH:O	2.18	0.42
1:A:55:LEU:HD21	1:A:146:PHE:CD1	2.54	0.42
1:A:88:LYS:HA	1:A:91:MET:HE3	2.00	0.42
1:C:13:GLY:C	1:C:15:TYR:H	2.23	0.42
1:C:270:ASP:HA	1:C:271:PRO:HD3	1.92	0.42
1:A:-1:GLY:HA3	1:A:0:SER:HA	1.74	0.42
2:D:323:GLN:HA	2:D:324:PRO:HA	1.87	0.42
2:D:361:HIS:HB2	2:D:372:TRP:HB2	2.01	0.42
2:D:193:CYS:HG	5:D:3:SGM:C1	2.29	0.41
2:D:414:LYS:HG2	2:D:423:LEU:HG	2.02	0.41
1:C:73:GLU:OE2	2:D:293:ARG:NH2	2.53	0.41
1:C:41:THR:HG22	1:C:42:GLU:H	1.84	0.41
1:A:162:GLU:HG3	1:A:163:VAL:HA	2.02	0.41
1:C:133:LEU:HD11	1:C:192:ILE:HD13	2.02	0.41
2:D:403:GLN:HG3	6:D:527:HOH:O	2.21	0.41
2:D:179:HIS:HB2	6:D:716:HOH:O	2.20	0.41
1:A:119:HIS:HE1	1:A:185:ASP:OD2	2.04	0.40
1:C:253:PRO:HD2	1:C:254:PRO:HD3	2.01	0.40
1:A:175:LEU:HB2	1:A:233:MET:HE1	2.02	0.40
1:A:86:ASP:OD2	1:A:89:LYS:HE3	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:297:ARG:NH2	1:C:251:VAL:CG2[3_454]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	297/300 (99%)	283 (95%)	9 (3%)	5 (2%)	11	4
1	C	267/300 (89%)	254 (95%)	8 (3%)	5 (2%)	10	4
2	B	263/262 (100%)	261 (99%)	2 (1%)	0	100	100
2	D	261/262 (100%)	256 (98%)	5 (2%)	0	100	100
All	All	1088/1124 (97%)	1054 (97%)	24 (2%)	10 (1%)	21	13

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	15	TYR
1	A	162	GLU
1	C	162	GLU
1	C	163	VAL
1	A	164	VAL
1	C	164	VAL
1	A	96	LEU
1	C	14	THR
1	C	16	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/263 (101%)	254 (96%)	11 (4%)	36	31

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	239/263 (91%)	228 (95%)	11 (5%)	33	28
2	B	238/235 (101%)	232 (98%)	6 (2%)	55	55
2	D	236/235 (100%)	227 (96%)	9 (4%)	40	36
All	All	978/996 (98%)	941 (96%)	37 (4%)	40	36

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	0	SER
1	A	75	LYS
1	A	97	THR
1	A	101	LEU
1	A	122	ARG
1	A	148	LEU
1	A	150	ARG
1	A	162	GLU
1	A	163	VAL
1	A	297	ARG
1	A	298	LEU
2	B	196	LYS
2	B	199	TYR
2	B	232	LEU
2	B	292	LEU
2	B	348	LEU
2	B	384	LEU
1	C	14	THR
1	C	15	TYR
1	C	36	ARG
1	C	55	LEU
1	C	83	LEU
1	C	101	LEU
1	C	122	ARG
1	C	131	GLN
1	C	148	LEU
1	C	150	ARG
1	C	163	VAL
2	D	175	VAL
2	D	196	LYS
2	D	232	LEU
2	D	284	ASP
2	D	292	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	296	HIS
2	D	348	LEU
2	D	391	LEU
2	D	403	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (29) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	60	HIS
1	A	74	ASN
1	A	119	HIS
1	A	161	HIS
2	B	183	HIS
2	B	233	HIS
2	B	254	GLN
2	B	317	GLN
2	B	395	HIS
2	B	403	GLN
1	C	59	ASN
1	C	60	HIS
1	C	71	HIS
1	C	84	HIS
1	C	85	GLN
1	C	113	GLN
1	C	119	HIS
1	C	161	HIS
1	C	265	GLN
1	C	268	HIS
1	C	287	GLN
1	C	295	HIS
2	D	173	ASN
2	D	183	HIS
2	D	233	HIS
2	D	254	GLN
2	D	296	HIS
2	D	317	GLN
2	D	396	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	TPO	A	160	1	8,10,11	2.31	3 (37%)	7,14,16	1.98	2 (28%)
1	TPO	C	160	1	8,10,11	2.35	3 (37%)	7,14,16	1.20	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	160	1	-	0/8/11/13	0/0/0/0
1	TPO	C	160	1	-	0/8/11/13	0/0/0/0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	160	TPO	P-O3P	-4.37	1.39	1.54
1	C	160	TPO	P-O3P	-4.17	1.39	1.54
1	C	160	TPO	P-O2P	-3.73	1.41	1.54
1	A	160	TPO	P-O2P	-3.67	1.41	1.54
1	C	160	TPO	P-O1P	-3.08	1.41	1.51
1	A	160	TPO	P-O1P	-2.78	1.42	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	160	TPO	OG1-P-O1P	-3.35	98.73	107.11
1	C	160	TPO	O-C-CA	-2.06	119.99	125.44
1	A	160	TPO	O3P-P-O2P	2.40	116.53	107.38

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	160	TPO	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	MFR	A	299	-	19,20,20	1.24	2 (10%)	20,28,28	3.65	11 (55%)
5	SGM	B	4	-	5,5,5	0.48	0	5,5,5	0.58	0
5	SGM	B	5	-	5,5,5	0.47	0	5,5,5	0.41	0
4	MFR	C	299	-	19,20,20	1.27	2 (10%)	20,28,28	3.58	11 (55%)
5	SGM	D	3	-	5,5,5	0.44	0	5,5,5	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MFR	A	299	-	-	0/2/6/6	0/3/3/3
5	SGM	B	4	-	-	0/4/4/4	0/0/0/0
5	SGM	B	5	-	-	0/4/4/4	0/0/0/0
4	MFR	C	299	-	-	0/2/6/6	0/3/3/3
5	SGM	D	3	-	-	0/4/4/4	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	299	MFR	C9-C10	2.37	1.45	1.41
4	C	299	MFR	C9-C10	2.55	1.45	1.41
4	A	299	MFR	C1-C10	2.94	1.49	1.42
4	C	299	MFR	C1-C10	3.04	1.49	1.42

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	299	MFR	N7-C12-N5	-5.61	119.92	125.78
4	C	299	MFR	N7-C12-N5	-5.29	120.26	125.78
4	C	299	MFR	C8-C9-C11	-4.23	116.25	126.30
4	C	299	MFR	C14-C13-N7	-3.85	119.51	123.90
4	A	299	MFR	C14-C13-N7	-3.12	120.34	123.90
4	A	299	MFR	C8-C9-C11	-3.02	119.13	126.30
4	A	299	MFR	C2-C1-C10	-2.66	115.23	120.51
4	A	299	MFR	C14-C11-N5	-2.58	118.46	122.01
4	C	299	MFR	C2-C1-C10	-2.55	115.44	120.51
4	C	299	MFR	C14-C11-N5	-2.37	118.74	122.01
4	C	299	MFR	C15-O1-C1	3.11	122.32	117.77
4	A	299	MFR	C15-O1-C1	3.31	122.61	117.77
4	A	299	MFR	O1-C1-C10	3.45	121.34	115.89
4	A	299	MFR	C3-C2-C1	3.58	122.46	117.71
4	C	299	MFR	C3-C2-C1	3.59	122.47	117.71
4	C	299	MFR	N6-C12-N7	3.76	120.78	117.39
4	C	299	MFR	O1-C1-C10	4.33	122.73	115.89
4	A	299	MFR	N6-C12-N7	4.57	121.51	117.39
4	A	299	MFR	C13-N7-C12	4.68	120.53	116.32
4	C	299	MFR	C13-N7-C12	5.02	120.84	116.32
4	C	299	MFR	C11-N5-C12	9.41	121.44	116.34
4	A	299	MFR	C11-N5-C12	10.67	122.12	116.34

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	4	SGM	1	0
5	B	5	SGM	2	0
5	D	3	SGM	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	297/300 (99%)	0.77	31 (10%) 8 9	2, 5, 22, 43	0
1	C	270/300 (90%)	1.02	37 (13%) 4 4	2, 6, 33, 59	0
2	B	262/262 (100%)	0.50	15 (5%) 27 29	2, 5, 16, 27	0
2	D	262/262 (100%)	0.84	31 (11%) 6 6	2, 6, 21, 43	0
All	All	1091/1124 (97%)	0.78	114 (10%) 8 9	2, 6, 24, 59	0

All (114) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	249	SER	11.3
1	C	221	THR	9.1
1	A	96	LEU	8.9
1	A	14	THR	7.6
1	C	251	VAL	7.2
2	D	171	SER	7.1
2	D	432	VAL	7.0
1	C	14	THR	6.8
2	D	428	GLU	6.7
1	C	250	LYS	6.1
1	A	-1	GLY	5.9
1	C	248	PHE	5.8
1	A	41	THR	5.8
1	A	0	SER	5.7
1	C	15	TYR	5.7
2	B	171	SER	5.7
1	C	13	GLY	5.4
1	C	253	PRO	5.2
1	A	36	ARG	4.9
2	D	284	ASP	4.8
1	C	287	GLN	4.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	38	ASP	4.7
2	D	431	ASN	4.6
2	B	283[A]	ASP	4.5
1	C	72	THR	4.5
2	D	283	ASP	4.2
1	C	295	HIS	4.2
2	B	432	VAL	4.1
2	D	429	THR	4.1
1	C	296	LEU	4.1
1	C	288	ASP	4.0
2	B	284	ASP	3.9
1	A	2	GLU	3.8
1	A	1	MET	3.8
1	C	282	ALA	3.8
2	D	336	LEU	3.8
1	C	252	VAL	3.7
1	C	269	TYR	3.6
1	C	84	HIS	3.5
1	C	73	GLU	3.5
1	A	73	GLU	3.4
1	C	2	GLU	3.4
2	B	323	GLN	3.4
1	C	257	GLU	3.3
2	D	362	LEU	3.3
2	D	332	LEU	3.3
1	C	173	ILE	3.3
2	D	430	LEU	3.3
2	D	280	TYR	3.2
2	B	428	GLU	3.2
1	A	97	THR	3.2
1	C	108	LEU	3.2
1	A	247	ASP	3.1
1	C	220	GLY	3.1
2	B	431	ASN	3.1
2	B	280	TYR	3.1
1	A	55	LEU	3.0
1	C	36	ARG	3.0
2	D	359	ALA	2.9
2	D	186	LEU	2.9
1	A	170	ALA	2.9
2	D	355	ILE	2.9
1	C	284	PRO	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	363	ALA	2.9
2	D	365	TYR	2.8
2	B	285	THR	2.8
1	A	104	ILE	2.8
1	A	298	LEU	2.8
2	D	394	LEU	2.8
1	A	74	ASN	2.7
1	C	293	VAL	2.7
1	C	216	PHE	2.7
2	B	201	LYS	2.7
1	A	206[A]	ASP	2.7
1	A	232	SER	2.7
1	A	162	GLU	2.6
2	D	179	HIS	2.6
1	A	15	TYR	2.6
2	D	425	ASN	2.5
2	B	362	LEU	2.5
2	D	396	GLN	2.5
1	A	95	ALA	2.5
1	A	32	LEU	2.5
2	D	357	ALA	2.5
1	C	0[A]	SER	2.5
2	B	296	HIS	2.5
1	A	287	GLN	2.5
1	A	175	LEU	2.4
2	D	400	ARG	2.4
1	C	273	LYS	2.4
2	D	356	ALA	2.4
1	A	103	LEU	2.4
1	C	5	GLN	2.4
2	D	333	ALA	2.3
1	A	297	ARG	2.3
2	D	256	VAL	2.3
1	C	41	THR	2.3
2	D	423	LEU	2.3
1	C	206	ASP	2.2
2	B	293	ARG	2.2
1	A	64	VAL	2.2
1	C	184	VAL	2.2
2	D	385	GLU	2.1
2	D	323	GLN	2.1
1	A	233	MET	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	71	HIS	2.1
1	C	258	ASP	2.1
1	A	231	THR	2.1
2	B	429	THR	2.1
1	A	161	HIS	2.1
2	D	360	PHE	2.1
2	D	378	GLN	2.0
2	B	385[A]	GLU	2.0
1	C	209	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	TPO	A	160	11/12	0.98	0.11	-	2,4,6,7	0
1	TPO	C	160	11/12	0.98	0.10	-	2,4,7,7	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	SGM	B	4	6/6	0.88	0.26	3.53	30,31,33,34	0
5	SGM	D	3	6/6	0.75	0.28	2.86	41,43,43,44	0
3	MG	D	2	1/1	0.98	0.23	2.58	9,9,9,9	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	MG	B	1	1/1	0.95	0.16	-0.00	15,15,15,15	0
4	MFR	C	299	18/18	0.95	0.10	-1.61	2,2,4,5	0
4	MFR	A	299	18/18	0.96	0.10	-1.67	2,2,2,2	0
5	SGM	B	5	6/6	0.42	0.41	-	61,63,63,64	0

6.5 Other polymers [i](#)

There are no such residues in this entry.