



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:06 PM GMT

PDB ID : 4BJZ
Title : Crystal structure of 3-hydroxybenzoate 6-hydroxylase uncovers lipid- assisted flavoprotein strategy for regioselective aromatic hydroxylation: Native data
Authors : Orru, R.; Montersino, S.; Barendregt, A.; Westphal, A.H.; van Duijn, E.; Mattevi, A.; van Berkel, W.J.H.
Deposited on : 2013-04-21
Resolution : 1.51 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2658 (1.54-1.50)
Clashscore	102246	2887 (1.54-1.50)
Ramachandran outliers	100387	2818 (1.54-1.50)
Sidechain outliers	100360	2816 (1.54-1.50)
RSRZ outliers	91569	2660 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	424	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	P3A	A	600[A]	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	P3A	A	600[B]	-	-	-	X
4	CL	A	1398	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3636 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROBABLE SALICYLATE MONOOXYGENASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	395	3083	1936	548	588	11	0	1	0

There are 25 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	400	LYS	-	EXPRESSION TAG	UNP Q0SFK6
A	401	LEU	-	EXPRESSION TAG	UNP Q0SFK6
A	402	GLY	-	EXPRESSION TAG	UNP Q0SFK6
A	403	PRO	-	EXPRESSION TAG	UNP Q0SFK6
A	404	GLU	-	EXPRESSION TAG	UNP Q0SFK6
A	405	GLN	-	EXPRESSION TAG	UNP Q0SFK6
A	406	LYS	-	EXPRESSION TAG	UNP Q0SFK6
A	407	LEU	-	EXPRESSION TAG	UNP Q0SFK6
A	408	ILE	-	EXPRESSION TAG	UNP Q0SFK6
A	409	SER	-	EXPRESSION TAG	UNP Q0SFK6
A	410	GLU	-	EXPRESSION TAG	UNP Q0SFK6
A	411	GLU	-	EXPRESSION TAG	UNP Q0SFK6
A	412	ASP	-	EXPRESSION TAG	UNP Q0SFK6
A	413	LEU	-	EXPRESSION TAG	UNP Q0SFK6
A	414	ASN	-	EXPRESSION TAG	UNP Q0SFK6
A	415	SER	-	EXPRESSION TAG	UNP Q0SFK6
A	416	ALA	-	EXPRESSION TAG	UNP Q0SFK6
A	417	VAL	-	EXPRESSION TAG	UNP Q0SFK6
A	418	ASP	-	EXPRESSION TAG	UNP Q0SFK6
A	419	HIS	-	EXPRESSION TAG	UNP Q0SFK6
A	420	HIS	-	EXPRESSION TAG	UNP Q0SFK6
A	421	HIS	-	EXPRESSION TAG	UNP Q0SFK6
A	422	HIS	-	EXPRESSION TAG	UNP Q0SFK6
A	423	HIS	-	EXPRESSION TAG	UNP Q0SFK6
A	424	HIS	-	EXPRESSION TAG	UNP Q0SFK6


- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:

C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is PHOSPHATIDYLGLYCEROL-PHOSPHOGLYCEROL (three-letter code: P3A) (formula: C₄₁H₇₈O₁₅P₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	1
			50	36	12	2		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total 2	Cl 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	448	Total 448	O 448	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PROBABLE SALICYLATE MONOOXYGENASE



4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	106.79 Å 106.79 Å 142.72 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.51 33.70 – 1.51	Depositor EDS
% Data completeness (in resolution range)	98.8 (30.00-1.51) 98.8 (33.70-1.51)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.65 (at 1.51 Å)	Xtriage
Refinement program	REFMAC 5.8.0033	Depositor
R, R_{free}	0.168 , 0.201 0.181 , 0.213	Depositor DCC
R_{free} test set	3215 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	14.9	Xtriage
Anisotropy	0.008	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 45.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 63866 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3636	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: P3A, FAD, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.96	0/3158	1.05	12/4287 (0.3%)

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	377	ARG	NE-CZ-NH2	-12.97	113.81	120.30
1	A	122	ARG	NE-CZ-NH2	-11.44	114.58	120.30
1	A	122	ARG	NE-CZ-NH1	10.81	125.71	120.30
1	A	372	ARG	NE-CZ-NH2	-8.88	115.86	120.30
1	A	377	ARG	NE-CZ-NH1	7.27	123.93	120.30
1	A	92	ASP	CB-CG-OD1	6.21	123.89	118.30
1	A	313	ASP	CB-CG-OD1	6.14	123.82	118.30
1	A	288	ARG	NE-CZ-NH1	5.86	123.23	120.30
1	A	365	ASP	CB-CG-OD2	-5.63	113.23	118.30
1	A	388	ASP	CB-CG-OD2	-5.62	113.24	118.30
1	A	319	ASP	CB-CG-OD2	-5.38	113.46	118.30
1	A	365	ASP	CB-CG-OD1	5.22	123.00	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3083	0	2962	51	0
2	A	53	0	31	0	0
3	A	50	0	6	0	0
4	A	2	0	0	2	0
5	A	448	0	0	42	1
All	All	3636	0	2999	52	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:202:GLU:HB3	5:A:2261:HOH:O	1.43	1.18
1:A:224:MET:HG2	5:A:2253:HOH:O	1.54	1.05
1:A:122:ARG:HG2	5:A:2163:HOH:O	1.68	0.92
1:A:271:ARG:HG3	5:A:2313:HOH:O	1.72	0.88
1:A:86:GLU:OE1	5:A:2116:HOH:O	1.99	0.80
1:A:177:GLU:HG2	5:A:2237:HOH:O	1.86	0.75
1:A:220:ARG:HB3	5:A:2273:HOH:O	1.86	0.73
1:A:96:GLU:CG	5:A:2130:HOH:O	2.37	0.72
1:A:242:ASN:HB2	5:A:2289:HOH:O	1.89	0.72
1:A:86:GLU:CB	5:A:2116:HOH:O	2.40	0.69
1:A:96:GLU:HG2	5:A:2133:HOH:O	1.93	0.67
1:A:172:LYS:HE2	5:A:2188:HOH:O	1.94	0.65
1:A:96:GLU:HG2	5:A:2130:HOH:O	1.94	0.64
1:A:202:GLU:HG2	5:A:2263:HOH:O	1.97	0.63
1:A:86:GLU:HB3	5:A:2116:HOH:O	1.98	0.61
4:A:1398:CL:CL	4:A:1399:CL:CL	2.92	0.61
1:A:338:LYS:HE2	5:A:2368:HOH:O	2.02	0.60
1:A:271:ARG:HD3	5:A:2314:HOH:O	2.02	0.60
1:A:290:ILE:HG23	5:A:2345:HOH:O	2.01	0.60
1:A:220:ARG:NH1	1:A:224:MET:HE2	2.17	0.59
1:A:3:ASN:CB	5:A:2001:HOH:O	2.51	0.58
1:A:242:ASN:CB	5:A:2289:HOH:O	2.47	0.57
1:A:224:MET:HG3	5:A:2273:HOH:O	2.05	0.57
1:A:96:GLU:CG	5:A:2133:HOH:O	2.49	0.57
1:A:323:GLU:HG3	5:A:2359:HOH:O	2.07	0.54
1:A:224:MET:N	5:A:2273:HOH:O	2.40	0.54
1:A:299:PRO:HD2	5:A:2347:HOH:O	2.07	0.54
1:A:292:LEU:HD12	5:A:2345:HOH:O	2.08	0.52
1:A:72:PHE:N	5:A:2096:HOH:O	2.42	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:66:ASP:HB2	5:A:2075:HOH:O	2.10	0.51
1:A:224:MET:CG	5:A:2253:HOH:O	2.29	0.51
1:A:397:ARG:C	5:A:2444:HOH:O	2.50	0.50
1:A:301:GLN:HA	4:A:1398:CL:CL	2.49	0.48
1:A:216:GLN:HG2	1:A:227:GLN:HG2	1.95	0.48
1:A:258:GLU:OE2	1:A:262:ARG:HD3	2.14	0.47
1:A:177:GLU:CG	5:A:2237:HOH:O	2.57	0.47
1:A:397:ARG:NH1	5:A:2370:HOH:O	2.21	0.46
1:A:220:ARG:CZ	1:A:224:MET:HE2	2.45	0.46
1:A:71:ALA:C	5:A:2096:HOH:O	2.53	0.46
1:A:3:ASN:HD21	1:A:5:GLN:NE2	2.14	0.45
1:A:292:LEU:CD1	5:A:2345:HOH:O	2.64	0.44
1:A:258:GLU:O	1:A:262:ARG:HG3	2.17	0.44
1:A:169:LEU:HD22	5:A:2220:HOH:O	2.17	0.44
1:A:143:LYS:HE3	5:A:2200:HOH:O	2.17	0.43
1:A:160:LEU:HB3	5:A:2345:HOH:O	2.18	0.43
1:A:3:ASN:CA	5:A:2001:HOH:O	2.66	0.43
1:A:96:GLU:HG3	5:A:2130:HOH:O	2.12	0.42
1:A:292:LEU:HG	5:A:2345:HOH:O	2.20	0.41
1:A:385:LYS:HE3	1:A:386:TYR:CE1	2.55	0.41
1:A:77:ILE:HG13	1:A:93:LEU:HD11	2.03	0.41
1:A:202:GLU:CG	5:A:2263:HOH:O	2.64	0.40
1:A:143:LYS:HG3	5:A:2200:HOH:O	2.20	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:2298:HOH:O	5:A:2368:HOH:O[3_545]	1.75	0.45

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	394/424 (93%)	386 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	312/337 (93%)	304 (97%)	8 (3%)	54	19

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ASN
1	A	43	GLU
1	A	53	HIS
1	A	143	LYS
1	A	198	ASP
1	A	271	ARG
1	A	341	ASN
1	A	347	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	FAD	A	500	-	48,58,58	1.55	8 (16%)	54,89,89	2.54	15 (27%)
3	P3A	A	600[A]	-	43,43,57	1.09	4 (9%)	47,48,67	1.08	3 (6%)
3	P3A	A	600[B]	-	43,43,57	1.03	4 (9%)	47,48,67	1.07	3 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	500	-	-	0/30/50/50	0/6/6/6
3	P3A	A	600[A]	-	-	0/45/45/65	0/0/0/0
3	P3A	A	600[B]	-	-	0/45/45/65	0/0/0/0

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	FAD	C1'-N10	-2.56	1.45	1.48
2	A	500	FAD	C10-N1	-2.54	1.31	1.35
3	A	600[A]	P3A	O37-C38	-2.46	1.40	1.46
3	A	600[B]	P3A	O37-C38	-2.46	1.40	1.46
2	A	500	FAD	C6-C5X	-2.37	1.38	1.41
2	A	500	FAD	C9A-N10	2.13	1.41	1.38
2	A	500	FAD	C9A-C5X	2.36	1.47	1.42
3	A	600[B]	P3A	P16-O18	2.49	1.58	1.50
2	A	500	FAD	C8-C7	2.74	1.48	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	600[A]	P3A	O41-C42	2.85	1.41	1.33
3	A	600[B]	P3A	O41-C42	2.85	1.41	1.33
3	A	600[A]	P3A	P16-O18	3.59	1.62	1.50
3	A	600[A]	P3A	O37-C21	3.79	1.45	1.34
3	A	600[B]	P3A	O37-C21	3.79	1.45	1.34
2	A	500	FAD	C4X-C10	4.34	1.49	1.41
2	A	500	FAD	O4B-C1B	4.88	1.47	1.41

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	FAD	N3A-C2A-N1A	-7.93	122.82	128.89
2	A	500	FAD	C4-C4X-C10	-7.71	115.01	119.94
2	A	500	FAD	C4X-C10-N10	-6.92	116.44	120.52
2	A	500	FAD	C4B-O4B-C1B	-4.07	105.25	109.72
2	A	500	FAD	C1B-N9A-C4A	-2.99	122.43	126.94
2	A	500	FAD	O4B-C1B-N9A	-2.86	102.10	108.10
2	A	500	FAD	C4X-C4-N3	-2.63	119.99	123.59
3	A	600[A]	P3A	C34-C33-C32	-2.34	102.44	114.53
3	A	600[B]	P3A	C34-C33-C32	-2.34	102.44	114.53
3	A	600[B]	P3A	O18-P16-O17	-2.20	104.15	114.72
3	A	600[A]	P3A	O37-C38-C40	-2.14	100.84	108.36
3	A	600[B]	P3A	O37-C38-C40	-2.14	100.84	108.36
2	A	500	FAD	N6A-C6A-N1A	2.02	123.53	119.20
2	A	500	FAD	O4B-C4B-C3B	2.02	109.21	105.15
2	A	500	FAD	C2A-N1A-C6A	2.08	122.49	118.77
2	A	500	FAD	C1'-N10-C9A	2.25	121.39	118.86
3	A	600[A]	P3A	O15-P16-O17	2.36	119.52	110.85
2	A	500	FAD	C2B-C1B-N9A	2.42	117.99	114.29
2	A	500	FAD	C4X-N5-C5X	3.55	120.85	116.76
2	A	500	FAD	C4-C4X-N5	3.78	123.31	118.72
2	A	500	FAD	C4-N3-C2	7.90	122.07	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	395/424 (93%)	0.14	13 (3%) 50 52	8, 17, 36, 59	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	198	ASP	5.0
1	A	197	LEU	3.5
1	A	5	GLN	2.8
1	A	326	SER	2.6
1	A	192	TYR	2.4
1	A	242	ASN	2.3
1	A	262	ARG	2.3
1	A	69	SER	2.3
1	A	327	THR	2.3
1	A	220	ARG	2.1
1	A	265	ASP	2.1
1	A	271	ARG	2.1
1	A	241	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	P3A	A	600[A]	44/58	0.79	0.17	3.68	20,24,30,33	6
3	P3A	A	600[B]	44/58	0.79	0.17	3.68	20,24,30,33	6
2	FAD	A	500	53/53	0.98	0.09	-0.28	9,13,18,20	0
4	CL	A	1399	1/1	0.98	0.06	-4.06	24,24,24,24	0
4	CL	A	1398	1/1	0.99	0.05	-4.58	18,18,18,18	0

6.5 Other polymers [i](#)

There are no such residues in this entry.