



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:46 AM GMT

PDB ID : 2BLA  
Title : SP21 DOUBLE MUTANT P. VIVAX DIHYDROFOLATE REDUCTASE IN COMPLEX WITH PYRIMETHAMINE  
Authors : Kongsaree, P.; Khongsuk, P.; Leartsakulpanich, U.; Chitnumsub, P.; Tarnchompoo, B.; Walkinshaw, M.D.; Yuthavong, Y.  
Deposited on : 2005-03-02  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	238	<div> <div>3%</div> <div>60%</div> <div>29%</div> <div>9%</div> </div>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2009 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called DIHYDROFOLATE REDUCTASE-THYMIDYLATE SYNTHASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	217	1781	1148	295	328	10	0	0	0

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	3	ASN	ASP	CONFLICT	UNP Q5U9H1
A	213	GLU	GLN	CONFLICT	UNP Q5U9H1

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula:  $C_{21}H_{30}N_7O_{17}P_3$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	48	21	7	17	3	0	0

- Molecule 3 is 5-(4-CHLORO-PHENYL)-6-ETHYL-PYRIMIDINE-2,4-DIAMINE (three-letter code: CP6) (formula:  $C_{12}H_{13}ClN_4$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	Cl	N	0	0
			17	12	1	4		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula:  $C_6H_{13}NO_4S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	151	Total	O	0	0
			151	151		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DIHYDROFOLATE REDUCTASE-THYMIDYLATE SYNTHASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	135.27Å 54.40Å 46.09Å 90.00° 108.13° 90.00°	Depositor
Resolution (Å)	6.00 – 2.50 32.14 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.1 (6.00-2.50) 99.2 (32.14-2.50)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.24 (at 2.51Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.194 , 0.282 0.204 , 0.192	Depositor DCC
$R_{free}$ test set	479 reflections (4.68%)	DCC
Wilson B-factor (Å <sup>2</sup> )	41.3	Xtriage
Anisotropy	0.416	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 66.0	EDS
Estimated twinning fraction	0.033 for -h-2*k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 11056 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2009	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.93% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, MES, CP6

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.52	0/1819	0.64	0/2450

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1781	0	1802	53	0
2	A	48	0	26	4	0
3	A	17	0	13	1	0
4	A	12	0	13	0	0
5	A	151	0	0	10	0
All	All	2009	0	1854	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:175:GLY:HA3	5:A:2106:HOH:O	1.65	0.96
1:A:174:GLY:HA3	2:A:1239:NDP:H5N	1.54	0.90
1:A:224:ASN:HD21	1:A:227:GLY:H	1.34	0.75
1:A:78:ARG:O	1:A:82:MET:HG2	1.90	0.72
1:A:34:ARG:HD3	5:A:2136:HOH:O	1.91	0.70
1:A:215:ARG:HG2	1:A:238:VAL:HB	1.73	0.69
1:A:22:THR:HG22	1:A:24:GLU:H	1.57	0.68
1:A:210:ASP:OD1	1:A:212:SER:OG	2.12	0.68
1:A:76:ARG:O	1:A:80:LEU:HG	1.94	0.67
1:A:218:SER:HB2	1:A:234:VAL:HB	1.80	0.64
1:A:174:GLY:CA	2:A:1239:NDP:H5N	2.26	0.64
1:A:186:ASN:HB2	5:A:2118:HOH:O	1.99	0.63
1:A:115:ARG:HD2	1:A:140:LEU:HD13	1.79	0.63
1:A:111:VAL:HG21	1:A:128:LEU:HD12	1.81	0.62
1:A:133:ASN:HD22	1:A:133:ASN:N	1.97	0.60
1:A:47:TRP:O	1:A:48:LYS:HB2	2.01	0.60
1:A:109:ASN:OD1	1:A:168:TYR:HB3	2.04	0.58
1:A:56:TYR:O	1:A:60:VAL:HG23	2.06	0.55
1:A:117:ASN:ND2	3:A:1240:CP6:H11	2.22	0.54
1:A:189:LYS:HG2	5:A:2004:HOH:O	2.05	0.54
1:A:144:ASP:HB2	5:A:2087:HOH:O	2.07	0.54
1:A:215:ARG:HB2	1:A:215:ARG:HH11	1.73	0.54
1:A:48:LYS:HA	1:A:48:LYS:HE3	1.91	0.53
1:A:124:GLN:H	1:A:124:GLN:HE21	1.56	0.53
1:A:127:PRO:HD3	1:A:147:GLU:HB3	1.90	0.53
1:A:215:ARG:NH1	1:A:215:ARG:HB2	2.24	0.52
1:A:146:LYS:HE2	5:A:2089:HOH:O	2.09	0.52
1:A:221:GLU:HG3	5:A:2141:HOH:O	2.11	0.51
1:A:141:THR:HB	1:A:143:GLU:OE2	2.11	0.51
1:A:40:GLY:HA2	1:A:46:PRO:HD3	1.92	0.51
1:A:124:GLN:NE2	1:A:124:GLN:H	2.10	0.49
1:A:18:LYS:HG2	1:A:35:THR:HG22	1.95	0.49
1:A:39:LEU:O	2:A:1239:NDP:H2N	2.13	0.48
1:A:2:GLU:N	5:A:2001:HOH:O	2.48	0.47
1:A:133:ASN:C	1:A:149:VAL:HG23	2.35	0.47
1:A:135:VAL:HG23	1:A:149:VAL:HG21	1.96	0.46
1:A:155:ILE:O	1:A:158:LEU:HB3	2.16	0.46
1:A:162:LEU:HD22	1:A:167:TYR:CZ	2.50	0.46
1:A:103:ASN:HB3	1:A:105:ASP:OD2	2.15	0.46
1:A:104:ALA:N	5:A:2076:HOH:O	2.15	0.45
1:A:133:ASN:O	1:A:149:VAL:HG23	2.17	0.45
1:A:78:ARG:HG3	1:A:81:ARG:HH21	1.83	0.44

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:113:MET:HA	1:A:174:GLY:O	2.18	0.44
1:A:8:PHE:HB3	1:A:167:TYR:OH	2.17	0.44
1:A:50:ASN:O	1:A:54:MET:HG3	2.18	0.43
1:A:222:VAL:HG12	1:A:223:TYR:N	2.32	0.43
1:A:186:ASN:HA	5:A:2120:HOH:O	2.19	0.43
1:A:136:LEU:HD23	1:A:152:ILE:HG13	2.01	0.43
1:A:41:ASN:O	1:A:44:THR:HG22	2.19	0.42
1:A:129:PRO:HD2	1:A:131:ARG:HH21	1.84	0.42
1:A:193:PHE:CE1	1:A:195:ARG:HG2	2.54	0.41
1:A:159:LEU:O	1:A:163:LYS:HG3	2.21	0.41
1:A:117:ASN:HD21	2:A:1239:NDP:C6N	2.34	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/238 (90%)	202 (95%)	9 (4%)	2 (1%)	21	37

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	48	LYS
1	A	81	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/215 (94%)	192 (95%)	10 (5%)	30	53

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	ARG
1	A	45	LEU
1	A	48	LYS
1	A	124	GLN
1	A	133	ASN
1	A	197	ASN
1	A	208	GLU
1	A	215	ARG
1	A	218	SER
1	A	224	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	108	GLN
1	A	124	GLN
1	A	186	ASN
1	A	224	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NDP	A	1239	-	42,52,52	1.81	5 (11%)	55,80,80	1.91	15 (27%)
3	CP6	A	1240	-	18,18,18	3.53	11 (61%)	21,25,25	1.76	7 (33%)
4	MES	A	1241	-	11,12,12	1.67	3 (27%)	14,16,16	0.76	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	1239	-	-	0/30/77/77	0/5/5/5
3	CP6	A	1240	-	-	0/6/6/6	0/2/2/2
4	MES	A	1241	-	-	0/6/14/14	0/1/1/1

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1239	NDP	C4N-C5N	-5.22	1.37	1.49
3	A	1240	CP6	C10-CL1	-4.58	1.64	1.74
2	A	1239	NDP	C2D-C1D	2.31	1.60	1.53
4	A	1241	MES	C5-N4	2.47	1.53	1.46
4	A	1241	MES	C7-N4	2.61	1.53	1.47
4	A	1241	MES	C3-N4	2.90	1.54	1.46
3	A	1240	CP6	C2-N14	3.06	1.40	1.34
3	A	1240	CP6	C9-C8	3.26	1.44	1.38
2	A	1239	NDP	C4A-N3A	3.30	1.40	1.35
3	A	1240	CP6	C12-C11	3.34	1.44	1.38
3	A	1240	CP6	C3-N1	3.47	1.40	1.35
3	A	1240	CP6	C2-N6	3.49	1.41	1.35
2	A	1239	NDP	C6N-C5N	3.55	1.40	1.33
3	A	1240	CP6	C2-N1	3.98	1.42	1.35

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1240	CP6	C11-C10	4.63	1.47	1.38
3	A	1240	CP6	C8-C7	4.89	1.49	1.39
2	A	1239	NDP	C2N-C3N	5.44	1.48	1.34
3	A	1240	CP6	C4-C5	5.50	1.53	1.41
3	A	1240	CP6	C4-C3	6.68	1.52	1.43

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1239	NDP	N3A-C2A-N1A	-5.35	124.80	128.89
2	A	1239	NDP	C3N-C2N-N1N	-3.92	117.53	123.14
2	A	1239	NDP	C4B-O4B-C1B	-3.88	105.46	109.72
3	A	1240	CP6	C4-C3-N13	-3.71	117.94	120.86
2	A	1239	NDP	C3B-C2B-C1B	-3.34	96.27	102.73
2	A	1239	NDP	C1D-N1N-C2N	-3.14	115.44	120.91
2	A	1239	NDP	O7N-C7N-N7N	-2.65	116.18	122.76
2	A	1239	NDP	C3D-C2D-C1D	-2.53	96.32	101.40
3	A	1240	CP6	C4-C3-N1	-2.14	121.23	122.55
3	A	1240	CP6	C2-N1-C3	2.07	119.40	117.04
3	A	1240	CP6	C12-C7-C4	2.11	124.31	120.78
2	A	1239	NDP	C2B-C3B-C4B	2.12	106.86	101.85
2	A	1239	NDP	C2D-C3D-C4D	2.32	107.37	102.61
3	A	1240	CP6	C7-C4-C3	2.41	123.37	120.74
2	A	1239	NDP	O3B-C3B-C2B	2.57	118.58	111.16
2	A	1239	NDP	PN-O3-PA	2.64	140.14	132.73
3	A	1240	CP6	N13-C3-N1	2.68	120.83	116.95
2	A	1239	NDP	P2B-O2B-C2B	2.73	128.12	121.56
2	A	1239	NDP	O3B-C3B-C4B	2.86	119.64	111.05
3	A	1240	CP6	C9-C10-CL1	2.97	124.23	119.35
2	A	1239	NDP	O4B-C1B-N9A	3.10	114.59	108.10
2	A	1239	NDP	C5N-C4N-C3N	3.81	123.02	112.52

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1239	NDP	4	0
3	A	1240	CP6	1	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	217/238 (91%)	-0.10	7 (3%) 51 56	28, 46, 66, 84	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	12	ALA	3.0
1	A	103	ASN	2.8
1	A	238	VAL	2.8
1	A	11	TYR	2.4
1	A	13	ILE	2.3
1	A	172	ILE	2.1
1	A	21	PRO	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	MES	A	1241	12/12	0.93	0.17	1.14	47,59,79,83	0
3	CP6	A	1240	17/17	0.90	0.18	0.36	15,23,44,52	0
2	NDP	A	1239	48/48	0.97	0.12	-0.57	40,55,88,92	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.