



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:08 PM GMT

PDB ID : 4BLP
Title : P4 PROTEIN FROM BACTERIOPHAGE PHI13
Authors : El Omari, K.; Meier, C.; Kainov, D.; Sutton, G.; Grimes, J.M.; Poranen, M.M.; Bamford, D.H.; Tuma, R.; Stuart, D.I.; Mancini, E.J.
Deposited on : 2013-05-04
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	358	<div> <div>4%</div> <div>70%</div> <div>5%</div> <div>24%</div> </div>
1	B	358	<div> <div>4%</div> <div>70%</div> <div>5%</div> <div>25%</div> </div>
1	C	358	<div> <div>4%</div> <div>69%</div> <div>7%</div> <div>25%</div> </div>
1	D	358	<div> <div>3%</div> <div>72%</div> <div>.</div> <div>24%</div> </div>
1	E	358	<div> <div>4%</div> <div>69%</div> <div>6%</div> <div>24%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	358	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	B	1304	-	-	-	X
3	GOL	C	1304	-	-	-	X
3	GOL	D	1307	-	-	-	X

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 13687 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PACKAGING ENZYME P4.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	271	Total	C	N	O	S	0	5	0
			1991	1263	342	380	6			
1	B	270	Total	C	N	O	S	0	12	0
			2019	1282	347	384	6			
1	C	270	Total	C	N	O	S	0	7	0
			1994	1266	343	379	6			
1	D	273	Total	C	N	O	S	0	2	0
			1991	1259	343	383	6			
1	E	271	Total	C	N	O	S	0	8	0
			2007	1275	344	382	6			
1	F	272	Total	C	N	O	S	0	7	0
			2008	1274	343	385	6			

- Molecule 2 is CITRATE ANION (three-letter code: FLC) (formula: $C_6H_5O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			13	6	7		
2	F	1	Total	C	O	0	0
			13	6	7		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	293	Total	O	0	0
			293	293		
4	B	263	Total	O	0	0
			263	263		
4	C	250	Total	O	0	0
			250	250		
4	D	293	Total	O	0	0
			293	293		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	277	Total 277	O 277	0	0
4	F	251	Total 251	O 251	0	0

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

[illegible]

Chain B:

Chain C:

69% 7% 25%


4%

MET THR ASP GLU LYS LYS VAL PRO SER LYS PRO ALA ARG PRO ALA ARG LYS PRO ASP LYS PRO ARG ASP LYS PRO ALA VAL PRO LYS V34 A35 A36 V39 K51 K52 S53 T60 V70 V77 V100 V110 V127 R128 P129 S134 H137


R142 S146 T155 G156 A157 V170 E179 F180 F181 D182 M183 L203 F220 G221 I222 I245 L249 M260 R285 R294 R299 I300 F301 S302 G303

PRO SER GLY PRO SER ALA ARG ARG ARG LEU LEU GLU SER ASN ASN ASN ILE ASP PRO ASP ASP ASP GLU GLU ASN ASP ASN ARG ARG ARG GLY ALA ARG ILE SER SER LEU


- Molecule 1: PACKAGING ENZYME P4



- Molecule 1: PACKAGING ENZYME P4



- Molecule 1: PACKAGING ENZYME P4



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	97.22Å 116.24Å 149.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.85 – 1.70 19.85 – 1.60	Depositor EDS
% Data completeness (in resolution range)	91.1 (19.85-1.70) 99.6 (19.85-1.60)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 1.60Å)	Xtriage
Refinement program	BUSTER 2.11.2	Depositor
R, R_{free}	0.163 , 0.188 0.161 , 0.182	Depositor DCC
R_{free} test set	9317 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	21.0	Xtriage
Anisotropy	0.458	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 55.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 221346 reflections (0.000%)	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	13687	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	0/2037	0.61	0/2785
1	B	0.60	0/2086	0.61	0/2853
1	C	0.59	0/2046	0.62	0/2799
1	D	0.61	0/2029	0.62	0/2777
1	E	0.62	0/2062	0.63	0/2821
1	F	0.60	0/2061	0.63	0/2821
All	All	0.60	0/12321	0.62	0/16856

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1991	0	2051	13	0
1	B	2019	0	2097	14	0
1	C	1994	0	2062	13	0
1	D	1991	0	2033	8	0
1	E	2007	0	2077	15	0
1	F	2008	0	2065	17	0
2	A	13	0	5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	13	0	5	0	0
3	B	6	0	8	0	0
3	C	6	0	8	0	0
3	D	6	0	8	0	0
3	E	6	0	8	0	0
4	A	293	0	0	1	0
4	B	263	0	0	2	0
4	C	250	0	0	1	0
4	D	293	0	0	2	0
4	E	277	0	0	2	0
4	F	251	0	0	1	0
All	All	13687	0	12427	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:70[B]:VAL:HG11	1:E:193:LEU:HD21	1.68	0.75
1:E:183:MET:SD	1:F:132:SER:HA	2.38	0.63
1:C:77[B]:VAL:HG22	1:C:110:VAL:HG22	1.80	0.62
1:A:128:ARG:HH21	1:F:107:ARG:HH12	1.47	0.62
1:D:55:ARG:NH1	4:D:2036:HOH:O	2.32	0.61
1:A:294:ARG:HD3	1:F:179[B]:GLU:HG2	1.81	0.61
1:E:137:HIS:HE1	1:E:146:SER:OG	1.85	0.59
1:B:179[B]:GLU:HG2	1:C:294:ARG:HD3	1.83	0.59
1:C:137:HIS:HE1	1:C:146:SER:OG	1.86	0.59
1:E:70[B]:VAL:HG23	1:E:238:ALA:HB1	1.85	0.58
1:B:170[B]:VAL:HG11	1:B:203:LEU:HD13	1.86	0.57
1:A:137:HIS:HE1	1:A:146:SER:OG	1.86	0.57
1:E:77[B]:VAL:HG22	1:E:110:VAL:HG22	1.85	0.57
1:C:170[B]:VAL:HG11	1:C:203:LEU:HD13	1.86	0.57
1:B:137:HIS:HE1	1:B:146:SER:OG	1.89	0.56
1:A:170[B]:VAL:HG11	1:A:203:LEU:HD13	1.88	0.56
1:D:137:HIS:HE1	1:D:146:SER:OG	1.88	0.55
1:E:170[B]:VAL:HG11	1:E:203:LEU:HD13	1.88	0.55
1:A:77[B]:VAL:HG22	1:A:110:VAL:HG22	1.88	0.55
1:F:137:HIS:HE1	1:F:146:SER:OG	1.89	0.55
1:B:294:ARG:HG2	1:B:299[B]:ARG:HD3	1.88	0.55
1:B:70[B]:VAL:HG12	1:B:222:ILE:HD11	1.87	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:128:ARG:NH2	1:F:107:ARG:HH12	2.08	0.52
1:F:70:VAL:HG12	1:F:222:ILE:HD11	1.90	0.52
1:B:300:ILE:HD11	4:B:2171:HOH:O	2.11	0.51
1:A:34:VAL:HG22	1:A:105:LEU:HD22	1.93	0.50
1:A:128:ARG:HH21	1:F:107:ARG:NH1	2.08	0.50
1:F:141:VAL:HG11	1:F:300:ILE:HD11	1.94	0.50
1:C:53:SER:HA	1:C:60:THR:O	2.12	0.50
1:D:180:THR:O	1:D:183:MET:HG3	2.12	0.50
4:A:2256:HOH:O	1:F:223:THR:HG21	2.12	0.49
1:B:103:ARG:NH2	4:B:2115:HOH:O	2.41	0.49
1:C:220:PHE:HE1	1:C:260:MET:CE	2.26	0.49
1:C:155:THR:HG22	4:D:2161:HOH:O	2.12	0.48
1:C:179:GLU:HG3	4:C:2186:HOH:O	2.13	0.48
1:B:53:SER:HA	1:B:60:THR:O	2.13	0.48
1:C:70:VAL:HG12	1:C:222:ILE:HD11	1.96	0.47
1:E:53:SER:HA	1:E:60:THR:O	2.14	0.47
1:A:53:SER:HA	1:A:60:THR:O	2.14	0.47
1:F:53:SER:HA	1:F:60:THR:O	2.15	0.47
1:B:151:VAL:HG21	1:B:163:LEU:HD21	1.97	0.47
1:E:183:MET:HG2	1:F:131:ALA:O	2.14	0.47
1:A:276:SER:OG	1:F:216:ARG:NH2	2.48	0.46
1:D:53:SER:HA	1:D:60:THR:O	2.16	0.46
1:B:70[B]:VAL:HG12	1:B:222:ILE:CD1	2.46	0.45
1:F:151:VAL:HG21	1:F:163:LEU:HD21	1.99	0.45
1:E:127:VAL:HG21	1:E:245[A]:ILE:HG12	1.99	0.45
1:F:229:GLY:HA2	1:F:265:LYS:HB3	1.97	0.44
1:C:127:VAL:HG21	1:C:245[A]:ILE:HG12	2.00	0.44
1:B:103:ARG:NH1	1:B:103:ARG:O	2.51	0.44
1:E:183:MET:HG3	4:E:2211:HOH:O	2.18	0.44
1:D:223:THR:HG21	4:E:2242:HOH:O	2.18	0.43
1:D:176:GLU:OE2	1:D:216:ARG:HD2	2.18	0.43
1:E:179:GLU:O	1:F:131:ALA:HB3	2.19	0.43
1:A:151:VAL:HG21	1:A:163:LEU:HD21	2.01	0.43
1:B:189:ALA:HB1	1:C:129:PRO:HD2	2.01	0.42
1:A:179[B]:GLU:HG2	1:B:294:ARG:HD3	1.99	0.42
1:A:221:GLY:HA3	1:B:124:LEU:HD22	2.01	0.42
1:D:188:ILE:HD12	1:D:203:LEU:HD11	2.01	0.41
1:C:220:PHE:HE1	1:C:260:MET:HE1	1.85	0.41
1:F:142:ARG:HG2	4:F:2144:HOH:O	2.21	0.41
1:D:151:VAL:HG21	1:D:163:LEU:HD21	2.02	0.41
1:E:294:ARG:HG2	1:E:299:ARG:HD3	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:39:VAL:HG21	1:F:59:MET:HE3	2.02	0.41
1:E:197[A]:LEU:HD21	1:E:242:LEU:HD22	2.01	0.41
1:C:77[B]:VAL:HG11	1:C:249:LEU:HD11	2.03	0.41
1:E:283:VAL:O	1:E:284:ASN:ND2	2.49	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/358 (76%)	269 (99%)	3 (1%)	0	100	100
1	B	280/358 (78%)	276 (99%)	4 (1%)	0	100	100
1	C	275/358 (77%)	272 (99%)	3 (1%)	0	100	100
1	D	273/358 (76%)	268 (98%)	5 (2%)	0	100	100
1	E	277/358 (77%)	274 (99%)	3 (1%)	0	100	100
1	F	277/358 (77%)	274 (99%)	3 (1%)	0	100	100
All	All	1654/2148 (77%)	1633 (99%)	21 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	217/287 (76%)	215 (99%)	2 (1%)	84	76
1	B	223/287 (78%)	222 (100%)	1 (0%)	93	90
1	C	218/287 (76%)	212 (97%)	6 (3%)	51	29
1	D	215/287 (75%)	213 (99%)	2 (1%)	84	76
1	E	220/287 (77%)	218 (99%)	2 (1%)	84	76
1	F	219/287 (76%)	217 (99%)	2 (1%)	84	76
All	All	1312/1722 (76%)	1297 (99%)	15 (1%)	80	69

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	134	SER
1	A	299	ARG
1	B	134	SER
1	C	39	VAL
1	C	51	LYS
1	C	134	SER
1	C	299[A]	ARG
1	C	299[B]	ARG
1	C	302	SER
1	D	134	SER
1	D	183	MET
1	E	134	SER
1	E	179	GLU
1	F	51	LYS
1	F	134	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	137	HIS
1	B	137	HIS
1	C	137	HIS
1	C	284	ASN
1	D	137	HIS
1	E	137	HIS
1	F	137	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FLC	A	1304	-	3,12,12	0.24	0	3,17,17	1.79	1 (33%)
3	GOL	B	1304	-	5,5,5	0.15	0	5,5,5	0.29	0
3	GOL	C	1304	-	5,5,5	0.27	0	5,5,5	0.13	0
3	GOL	D	1307	-	5,5,5	0.17	0	5,5,5	0.25	0
3	GOL	E	1305	-	5,5,5	0.13	0	5,5,5	0.21	0
2	FLC	F	1307	-	3,12,12	0.28	0	3,17,17	1.61	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FLC	A	1304	-	-	0/6/16/16	0/0/0/0
3	GOL	B	1304	-	-	0/4/4/4	0/0/0/0
3	GOL	C	1304	-	-	0/4/4/4	0/0/0/0
3	GOL	D	1307	-	-	0/4/4/4	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	E	1305	-	-	0/4/4/4	0/0/0/0
2	FLC	F	1307	-	-	0/6/16/16	0/0/0/0

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1304	FLC	CB-CG-CGC	2.22	118.50	114.96
2	F	1307	FLC	CB-CA-CAC	2.25	118.56	114.96

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	271/358 (75%)	-0.06	14 (5%)	31 33	8, 20, 43, 73	3 (1%)
1	B	270/358 (75%)	-0.07	15 (5%)	28 30	8, 16, 40, 69	4 (1%)
1	C	270/358 (75%)	-0.14	13 (4%)	34 38	7, 17, 40, 73	5 (1%)
1	D	273/358 (76%)	-0.28	10 (3%)	45 50	8, 15, 37, 65	3 (1%)
1	E	271/358 (75%)	-0.12	14 (5%)	31 33	7, 15, 37, 84	6 (2%)
1	F	272/358 (75%)	-0.13	12 (4%)	38 42	8, 17, 35, 56	5 (1%)
All	All	1627/2148 (75%)	-0.14	78 (4%)	34 38	7, 17, 40, 84	26 (1%)

All (78) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	303	GLY	6.2
1	C	301	PHE	5.6
1	B	100	VAL	5.5
1	D	228	PRO	5.4
1	B	301	PHE	5.3
1	F	306	ALA	5.3
1	E	35	ALA	5.3
1	A	32	PRO	5.3
1	B	34	VAL	5.3
1	E	93	ASP	4.9
1	B	35	ALA	4.8
1	A	183	MET	4.7
1	F	263	ASP	4.6
1	C	156	GLY	4.6
1	F	35	ALA	4.5
1	E	304	GLU	4.2
1	F	264	ASP	4.2
1	E	303	GLY	4.0
1	C	35	ALA	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	F	228	PRO	4.0
1	D	184	GLU	3.9
1	F	93	ASP	3.9
1	A	301	PHE	3.8
1	B	263	ASP	3.6
1	E	34	VAL	3.5
1	B	133	LEU	3.5
1	A	33	LYS	3.3
1	E	92	SER	3.3
1	E	301	PHE	3.2
1	A	167	THR	3.2
1	B	285	ASN	3.0
1	C	36	ALA	3.0
1	C	157	ALA	3.0
1	E	36	ALA	3.0
1	A	91	ALA	2.9
1	C	155	THR	2.9
1	B	228	PRO	2.9
1	F	262	SER	2.8
1	B	36	ALA	2.8
1	E	142	ARG	2.8
1	A	285	ASN	2.8
1	B	303	GLY	2.8
1	A	133	LEU	2.7
1	A	35	ALA	2.6
1	C	285	ASN	2.6
1	B	155	THR	2.6
1	D	134	SER	2.6
1	D	142	ARG	2.6
1	F	185	GLU	2.5
1	D	305	PRO	2.5
1	C	34	VAL	2.5
1	D	262	SER	2.5
1	F	96	SER	2.5
1	B	229	GLY	2.4
1	A	303	GLY	2.4
1	A	92	SER	2.4
1	B	103	ARG	2.4
1	E	167	THR	2.4
1	E	181	PHE	2.4
1	A	263	ASP	2.4
1	B	262	SER	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	132	SER	2.3
1	C	142	ARG	2.3
1	D	306	ALA	2.3
1	A	185	GLU	2.3
1	E	133	LEU	2.3
1	E	183	MET	2.2
1	C	181	PHE	2.2
1	C	183	MET	2.2
1	D	223	THR	2.2
1	F	285	ASN	2.2
1	E	155	THR	2.2
1	C	100	VAL	2.2
1	D	155	THR	2.1
1	A	302	SER	2.1
1	F	142	ARG	2.1
1	F	223	THR	2.0
1	D	263	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GOL	C	1304	6/6	0.93	0.23	5.41	16,28,32,35	0
3	GOL	D	1307	6/6	0.88	0.17	3.21	15,23,28,29	0
3	GOL	B	1304	6/6	0.85	0.15	2.06	16,29,31,32	0
2	FLC	A	1304	13/13	0.82	0.21	1.97	30,49,54,57	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	FLC	F	1307	13/13	0.90	0.18	1.72	24,34,37,39	0
3	GOL	E	1305	6/6	0.94	0.12	1.59	13,27,33,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.