



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 06:37 PM GMT

PDB ID : 1BNS  
Title : STRUCTURAL STUDIES OF BARNASE MUTANTS  
Authors : Chen, Y.W.  
Deposited on : 1994-04-11  
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1269 (2.04-2.04)
Ramachandran outliers	100387	1258 (2.04-2.04)
Sidechain outliers	100360	1258 (2.04-2.04)
RSRZ outliers	91569	1194 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	110	<div> <div></div> <div>74%21%...</div> </div>
1	B	110	<div> <div>11%</div> <div>51%36%7%..</div> </div>
1	C	110	<div> <div></div> <div>64%27%6%.</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2777 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BARNASE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	108	Total	C	N	O	0	0	0
			851	541	146	164			
1	B	108	Total	C	N	O	0	1	0
			850	540	147	163			
1	C	107	Total	C	N	O	0	0	0
			852	540	149	163			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	26	ALA	THR	CONFLICT	UNP P00648
B	26	ALA	THR	CONFLICT	UNP P00648
C	26	ALA	THR	CONFLICT	UNP P00648

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	65	Total	O	0	0
			65	65		
2	B	73	Total	O	0	0
			73	73		
2	C	86	Total	O	0	0
			86	86		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: BARNASE

Chain A: 




#### • Molecule 1: BARNASE

Chain B: 



#### • Molecule 1: BARNASE

Chain C: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.37Å 59.37Å 82.17Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	6.00 – 2.05 13.85 – 2.05	Depositor EDS
% Data completeness (in resolution range)	(Not available) (6.00-2.05) 94.1 (13.85-2.05)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	9.87 (at 2.04Å)	Xtriage
Refinement program	PROLSQ	Depositor
R, $R_{free}$	0.169 , (Not available) 0.164 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	19.8	Xtriage
Anisotropy	0.110	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 108.6	EDS
Estimated twinning fraction	0.004 for -h,-k,l 0.045 for h,-h-k,-l 0.027 for -k,-h,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 19072 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	2777	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.44% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.97	1/872 (0.1%)	2.09	24/1181 (2.0%)
1	B	1.04	0/876	2.20	35/1188 (2.9%)
1	C	1.05	1/873 (0.1%)	2.33	36/1180 (3.1%)
All	All	1.02	2/2621 (0.1%)	2.21	95/3549 (2.7%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	110	ARG	C-OXT	-11.03	1.02	1.23
1	A	110	ARG	C-OXT	-10.29	1.03	1.23

All (95) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	59	ARG	CD-NE-CZ	19.57	151.00	123.60
1	B	110	ARG	NE-CZ-NH2	16.07	128.34	120.30
1	A	72	ARG	NE-CZ-NH2	15.51	128.05	120.30
1	A	86	ASP	CB-CG-OD1	13.39	130.35	118.30
1	C	12	ASP	CB-CG-OD2	-13.17	106.45	118.30
1	C	32	ALA	CB-CA-C	12.59	128.99	110.10
1	C	13	TYR	CB-CG-CD1	-12.03	113.78	121.00
1	A	110	ARG	NE-CZ-NH2	11.98	126.29	120.30
1	C	110	ARG	NE-CZ-NH2	11.70	126.15	120.30
1	B	93	ASP	CA-CB-CG	11.28	138.21	113.40
1	C	13	TYR	CB-CG-CD2	11.19	127.71	121.00
1	C	90	TYR	CB-CG-CD2	10.43	127.26	121.00
1	B	8	ASP	CB-CG-OD2	-10.36	108.98	118.30
1	C	87	ARG	NE-CZ-NH2	10.28	125.44	120.30
1	B	12	ASP	CB-CG-OD2	10.03	127.33	118.30
1	C	59	ARG	NE-CZ-NH1	9.99	125.30	120.30
1	C	101	ASP	CB-CG-OD2	9.90	127.21	118.30
1	C	72	ARG	NE-CZ-NH1	9.48	125.04	120.30
1	B	83	ARG	NE-CZ-NH2	9.34	124.97	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	24	TYR	CB-CG-CD2	-9.06	115.56	121.00
1	A	101	ASP	CB-CG-OD2	9.01	126.41	118.30
1	B	69	ARG	NE-CZ-NH2	-8.88	115.86	120.30
1	A	22	ASP	CB-CA-C	-8.71	92.98	110.40
1	A	24	TYR	CB-CG-CD1	8.25	125.95	121.00
1	A	86	ASP	CB-CG-OD2	-7.76	111.32	118.30
1	C	110	ARG	NE-CZ-NH1	-7.72	116.44	120.30
1	C	12	ASP	CB-CG-OD1	7.39	124.95	118.30
1	C	87	ARG	NH1-CZ-NH2	-7.37	111.30	119.40
1	B	44	ASP	CB-CG-OD2	-7.32	111.71	118.30
1	A	8	ASP	CB-CG-OD2	7.12	124.71	118.30
1	A	12	ASP	CB-CG-OD2	-7.00	112.00	118.30
1	B	93	ASP	CB-CG-OD1	6.87	124.48	118.30
1	B	22	ASP	CB-CG-OD2	6.84	124.45	118.30
1	B	54	ASP	CB-CG-OD1	6.79	124.41	118.30
1	C	54	ASP	CB-CG-OD1	-6.75	112.23	118.30
1	C	75	ASP	CB-CG-OD1	6.68	124.31	118.30
1	B	75	ASP	CB-CG-OD1	6.67	124.31	118.30
1	B	19	LYS	CA-C-N	-6.60	102.68	117.20
1	C	69	ARG	NE-CZ-NH2	-6.59	117.00	120.30
1	C	110	ARG	N-CA-CB	6.53	122.35	110.60
1	B	24	TYR	CB-CG-CD2	-6.49	117.11	121.00
1	C	85	SER	N-CA-CB	-6.39	100.92	110.50
1	A	69	ARG	NE-CZ-NH1	-6.37	117.12	120.30
1	B	9	GLY	CA-C-O	6.36	132.05	120.60
1	B	95	LEU	N-CA-CB	6.35	123.10	110.40
1	C	78	TYR	CB-CG-CD1	-6.34	117.20	121.00
1	B	97	TYR	CB-CG-CD1	-6.32	117.21	121.00
1	C	110	ARG	CD-NE-CZ	6.29	132.41	123.60
1	C	97	TYR	CB-CG-CD2	-6.29	117.23	121.00
1	A	83	ARG	NE-CZ-NH2	6.17	123.38	120.30
1	A	89	LEU	CA-CB-CG	-6.09	101.30	115.30
1	A	37	ALA	CB-CA-C	6.05	119.18	110.10
1	A	38	SER	N-CA-CB	-6.03	101.46	110.50
1	B	69	ARG	CD-NE-CZ	5.97	131.96	123.60
1	C	70	THR	CA-CB-OG1	5.95	121.49	109.00
1	C	87	ARG	NE-CZ-NH1	5.89	123.25	120.30
1	C	26	ALA	N-CA-CB	-5.87	101.88	110.10
1	C	32	ALA	N-CA-CB	-5.85	101.90	110.10
1	B	110	ARG	NE-CZ-NH1	-5.83	117.38	120.30
1	B	101	ASP	CB-CG-OD1	5.80	123.52	118.30
1	B	101	ASP	N-CA-CB	-5.76	100.22	110.60

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	72	ARG	NE-CZ-NH1	-5.76	117.42	120.30
1	A	26	ALA	N-CA-CB	-5.75	102.05	110.10
1	C	90	TYR	CB-CG-CD1	-5.71	117.57	121.00
1	B	72	ARG	NE-CZ-NH2	-5.64	117.48	120.30
1	B	73	GLU	O-C-N	5.64	131.73	122.70
1	B	31	GLN	CG-CD-OE1	-5.63	110.33	121.60
1	A	101	ASP	CB-CG-OD1	-5.59	113.27	118.30
1	A	55	ILE	O-C-N	5.51	131.51	122.70
1	B	99	THR	O-C-N	5.48	131.47	122.70
1	C	66	LYS	CB-CG-CD	-5.48	97.36	111.60
1	A	69	ARG	CD-NE-CZ	5.45	131.23	123.60
1	C	83	ARG	NE-CZ-NH2	5.45	123.02	120.30
1	B	8	ASP	OD1-CG-OD2	5.44	133.64	123.30
1	A	70	THR	CA-CB-CG2	5.43	120.01	112.40
1	C	87	ARG	N-CA-CB	5.40	120.32	110.60
1	B	57	SER	O-C-N	5.39	131.33	122.70
1	B	29	GLU	OE1-CD-OE2	-5.39	116.83	123.30
1	A	78	TYR	CG-CD1-CE1	-5.33	117.03	121.30
1	B	29	GLU	CA-CB-CG	-5.33	101.68	113.40
1	B	24	TYR	CB-CG-CD1	5.32	124.19	121.00
1	C	93	ASP	CA-C-O	5.28	131.19	120.10
1	C	59	ARG	NE-CZ-NH2	-5.28	117.66	120.30
1	A	38	SER	CA-CB-OG	-5.28	96.95	111.20
1	C	31	GLN	C-N-CA	-5.16	108.79	121.70
1	C	50	SER	O-C-N	5.15	130.94	122.70
1	B	58	ASN	CA-CB-CG	5.14	124.72	113.40
1	C	7	PHE	CB-CG-CD2	5.12	124.38	120.80
1	B	5	ASN	CB-CG-OD1	-5.09	111.41	121.60
1	B	41	ASN	N-CA-CB	-5.08	101.45	110.60
1	C	71	TRP	CB-CG-CD1	-5.07	120.41	127.00
1	B	95	LEU	CA-CB-CG	5.05	126.92	115.30
1	A	6	THR	N-CA-CB	5.03	119.85	110.30
1	B	98	LYS	O-C-N	5.02	130.73	122.70
1	B	18	HIS	CA-C-O	5.00	130.60	120.10

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	851	0	811	6	0
1	B	850	0	802	22	0
1	C	852	0	820	13	0
2	A	65	0	0	0	0
2	B	73	0	0	1	0
2	C	86	0	0	3	0
All	All	2777	0	2433	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:28:SER:HA	1:C:31:GLN:HE21	1.31	0.95
1:B:64:PRO:HB2	1:B:69:ARG:NH1	2.12	0.64
1:A:26:ALA:HB3	1:A:29:GLU:HG3	1.79	0.64
1:B:64:PRO:HB2	1:B:69:ARG:HH12	1.63	0.64
1:C:28:SER:HA	1:C:31:GLN:NE2	2.09	0.63
1:C:57:SER:HB3	1:C:59:ARG:HD2	1.81	0.62
1:A:89:LEU:HD13	1:A:106:PHE:HE2	1.67	0.60
1:C:74:ALA:HB3	1:C:88:ILE:HG22	1.84	0.60
1:C:32:ALA:HB3	2:C:196:HOH:O	2.01	0.59
1:B:4:ILE:HD12	1:B:21:PRO:HB3	1.86	0.58
1:C:27:LYS:O	1:C:31:GLN:HG3	2.05	0.57
1:B:64:PRO:C	1:B:69:ARG:HH11	2.09	0.56
1:B:72:ARG:HB3	1:B:90:TYR:CZ	2.41	0.56
1:C:58:ASN:N	2:C:148:HOH:O	2.38	0.55
1:A:89:LEU:HD13	1:A:106:PHE:CE2	2.43	0.53
1:A:28:SER:HA	1:A:31:GLN:HE21	1.74	0.52
1:B:96:ILE:HB	1:B:110:ARG:HB2	1.91	0.52
1:B:96:ILE:HD12	1:B:110:ARG:HG3	1.92	0.51
1:A:21:PRO:HD2	1:A:24:TYR:CD1	2.46	0.49
1:B:7:PHE:CE2	1:B:98:LYS:HB2	2.47	0.49
1:B:89:LEU:HD13	1:B:106:PHE:HE2	1.77	0.49
1:B:93:ASP:O	1:B:94:TRP:HB2	2.13	0.49
1:B:43:ALA:HA	2:B:172:HOH:O	2.13	0.48
1:C:57:SER:CB	1:C:59:ARG:HD2	2.43	0.47
1:B:85:SER:HA	1:B:102[B]:HIS:CD2	2.49	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:89:LEU:HD13	1:B:106:PHE:CE2	2.50	0.46
1:B:21:PRO:HB2	1:B:23:ASN:OD1	2.16	0.46
1:B:15:GLN:O	1:B:18:HIS:NE2	2.49	0.45
1:A:20:LEU:HA	1:A:21:PRO:HD3	1.85	0.43
1:B:69:ARG:NH2	1:B:93:ASP:OD2	2.35	0.42
1:B:57:SER:OG	1:B:58:ASN:N	2.53	0.42
1:C:74:ALA:HB3	1:C:88:ILE:CG2	2.47	0.41
1:C:57:SER:HB2	2:C:146:HOH:O	2.20	0.41
1:B:74:ALA:HB3	1:B:88:ILE:CG2	2.50	0.41
1:B:64:PRO:HG2	1:B:71:TRP:HZ2	1.85	0.41
1:B:46:ALA:HB1	1:B:49:LYS:HG3	2.01	0.41
1:B:64:PRO:C	1:B:69:ARG:NH1	2.74	0.41
1:C:15:GLN:HE22	1:C:96:ILE:HD11	1.86	0.41
1:B:87:ARG:HB2	1:B:99:THR:HG22	2.02	0.41
1:C:55:ILE:HA	1:C:55:ILE:HD13	1.91	0.41
1:C:53:GLY:O	1:C:72:ARG:HD3	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	106/110 (96%)	100 (94%)	6 (6%)	0	100	100
1	B	107/110 (97%)	92 (86%)	13 (12%)	2 (2%)	10	2
1	C	105/110 (96%)	101 (96%)	4 (4%)	0	100	100
All	All	318/330 (96%)	293 (92%)	23 (7%)	2 (1%)	30	18

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	58	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	94	TRP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	87/91 (96%)	82 (94%)	5 (6%)	25	15
1	B	87/91 (96%)	77 (88%)	10 (12%)	7	2
1	C	88/91 (97%)	82 (93%)	6 (7%)	20	11
All	All	262/273 (96%)	241 (92%)	21 (8%)	15	7

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	67	SER
1	A	70	THR
1	A	89	LEU
1	A	98	LYS
1	A	103	TYR
1	B	19	LYS
1	B	55	ILE
1	B	66	LYS
1	B	67	SER
1	B	72	ARG
1	B	88	ILE
1	B	92	SER
1	B	93	ASP
1	B	98	LYS
1	B	103	TYR
1	C	8	ASP
1	C	15	GLN
1	C	27	LYS
1	C	57	SER
1	C	98	LYS
1	C	102	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	31	GLN
1	B	31	GLN
1	C	15	GLN
1	C	18	HIS
1	C	31	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	108/110 (98%)	-0.32	0 <span>100</span> <span>100</span>	11, 21, 40, 47	1 (0%)
1	B	108/110 (98%)	-0.01	12 (11%) <span>7</span> <span>7</span>	7, 22, 57, 66	3 (2%)
1	C	107/110 (97%)	-0.66	0 <span>100</span> <span>100</span>	7, 16, 30, 42	1 (0%)
All	All	323/330 (97%)	-0.33	12 (3%) <span>45</span> <span>51</span>	7, 19, 46, 66	5 (1%)

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	68	GLY	6.1
1	B	67	SER	4.9
1	B	71	TRP	3.3
1	B	58	ASN	3.1
1	B	16	THR	3.1
1	B	17	TYR	3.0
1	B	59	ARG	2.6
1	B	61	GLY	2.6
1	B	94	TRP	2.3
1	B	60	GLU	2.3
1	B	57	SER	2.2
1	B	105	THR	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.