



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 06:38 PM GMT

PDB ID : 1BRR  
Title : X-RAY STRUCTURE OF THE BACTERIORHODOPSIN TRIMER/LIPID COMPLEX  
Authors : Essen, L.-O.; Siegert, R.; Oesterhelt, D.  
Deposited on : 1998-07-28  
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1451 (2.90-2.90)
Clashscore	102246	1668 (2.90-2.90)
Ramachandran outliers	100387	1630 (2.90-2.90)
Sidechain outliers	100360	1632 (2.90-2.90)
RSRZ outliers	91569	1456 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	247	<div> <div>13%</div> <div>61%</div> <div>29%</div> <div>7%</div> </div>
1	B	247	<div> <div>9%</div> <div>62%</div> <div>28%</div> <div>6%</div> </div>
1	C	247	<div> <div>8%</div> <div>62%</div> <div>27%</div> <div>6%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	RET	A	999	-	-	-	X
5	ARC	A	1000	-	-	-	X
5	ARC	A	1001	-	-	-	X
5	ARC	A	1002	-	-	-	X
5	ARC	B	1000	-	-	-	X
5	ARC	B	1001	-	-	-	X
5	ARC	B	1002	-	-	-	X
5	ARC	B	1003	-	-	-	X
5	ARC	C	1000	-	-	-	X
5	ARC	C	1001	-	-	-	X
5	ARC	C	1002	-	-	-	X
6	OCT	B	1004	-	-	-	X

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 5672 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (BACTERIORHODOPSIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	230	Total	C	N	O	S	0	0	0
			1753	1177	265	302	9			
1	B	231	Total	C	N	O	S	0	0	0
			1770	1189	269	303	9			
1	C	232	Total	C	N	O	S	0	0	0
			1773	1188	271	305	9			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	PCA	GLN	?	UNP P0945

- Molecule 2 is a polymer of unknown type called SUGAR (3-MER).


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	3	Total	C	O	S	0	0
			38	18	19	1		
2	C	3	Total	C	O	S	0	0
			38	18	19	1		

- Molecule 3 is SUGAR (BETA-D-GLUCOSE) (three-letter code: BGC) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			12	6	6		

- Molecule 4 is RETINAL (three-letter code: RET) (formula:  $C_{20}H_{28}O$ ).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	C	0	0
			20	20		
4	B	1	Total	C	0	0
			20	20		
4	C	1	Total	C	0	0
			20	20		

- Molecule 5 is 3,7,11,15-TETRAMETHYL-HEXADECAN-1-OL (three-letter code: ARC) (formula: C<sub>20</sub>H<sub>42</sub>O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			21	20	1		
5	C	1	Total	C	O	0	0
			21	20	1		
5	A	1	Total	C	O	0	0
			21	20	1		
5	B	1	Total	C	O	0	0
			21	20	1		
5	B	1	Total	C	O	0	0
			21	20	1		
5	B	1	Total	C	O	0	0
			21	20	1		
5	B	1	Total	C	O	0	0
			21	20	1		
5	A	1	Total	C	O	0	0
			21	20	1		
5	A	1	Total	C	O	0	0
			21	20	1		
5	C	1	Total	C	O	0	0
			21	20	1		

- Molecule 6 is N-OCTANE (three-letter code: OCT) (formula: C<sub>8</sub>H<sub>18</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total C 8 8	0	0

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total C O 4 3 1	0	0
7	A	1	Total C 3 3	0	0
7	C	1	Total C 3 3	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: PROTEIN (BACTERIORHODOPSIN)




#### • Molecule 1: PROTEIN (BACTERIORHODOPSIN)



#### • Molecule 1: PROTEIN (BACTERIORHODOPSIN)







## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	120.52Å 105.96Å 80.19Å 90.00° 94.94° 90.00°	Depositor
Resolution (Å)	10.00 – 2.90 24.83 – 2.91	Depositor EDS
% Data completeness (in resolution range)	83.2 (10.00-2.90) 83.5 (24.83-2.91)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.67 (at 2.89Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, $R_{free}$	0.256 , 0.299 0.253 , 0.297	Depositor DCC
$R_{free}$ test set	530 reflections (3.01%)	DCC
Wilson B-factor (Å <sup>2</sup> )	65.6	Xtriage
Anisotropy	0.190	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 72.7	EDS
Estimated twinning fraction	0.000 for -1/2*h+1/2*k+1,1/2*h-1/2*k+1,1/2*h+1/2*k 0.000 for -1/2*h-1/2*k+1,-1/2*h-1/2*k-1,1/2*h-1/2*k	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 18504 reflections (0.005%)	Xtriage
$F_o, F_c$ correlation	0.87	EDS
Total number of atoms	5672	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.00% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, BGC, GLC, RET, ARC, SGA, PCA, OCT, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/1800	0.56	0/2462
1	B	0.43	0/1818	0.59	0/2485
1	C	0.45	0/1816	0.64	0/2484
All	All	0.42	0/5434	0.60	0/7431

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1753	0	1790	76	0
1	B	1770	0	1813	78	0
1	C	1773	0	1817	69	0
2	B	38	0	29	5	0
2	C	38	0	28	2	0
3	B	12	0	11	1	0
4	A	20	0	27	6	0
4	B	20	0	27	6	0
4	C	20	0	27	6	0
5	A	63	0	124	28	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	84	0	165	28	0
5	C	63	0	125	27	0
6	B	8	0	16	1	0
7	A	3	0	0	0	0
7	B	4	0	3	0	0
7	C	3	0	0	0	0
All	All	5672	0	6002	270	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 270 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:43:TYR:HB3	5:B:1000:ARC:H61	1.44	0.99
1:C:7:ARG:HH11	1:C:7:ARG:HG3	1.30	0.93
1:A:113:GLY:HA2	5:A:1002:ARC:H161	1.53	0.90
1:C:89:THR:HG21	1:C:216:LYS:HE3	1.54	0.89
2:B:250:SGA:H2	2:B:250:SGA:O3S	1.73	0.89

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	228/247 (92%)	201 (88%)	19 (8%)	8 (4%)	4	18
1	B	229/247 (93%)	201 (88%)	18 (8%)	10 (4%)	3	12
1	C	230/247 (93%)	201 (87%)	20 (9%)	9 (4%)	4	15
All	All	687/741 (93%)	603 (88%)	57 (8%)	27 (4%)	4	15

5 of 27 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	4	ILE
1	B	7	ARG
1	B	229	ILE
1	C	2	ALA
1	C	7	ARG

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	179/193 (93%)	167 (93%)	12 (7%)	20	50
1	B	181/193 (94%)	166 (92%)	15 (8%)	14	38
1	C	181/193 (94%)	164 (91%)	17 (9%)	11	32
All	All	541/579 (93%)	497 (92%)	44 (8%)	15	39

5 of 44 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	97	LEU
1	B	202	ASN
1	C	203	ILE
1	B	99	LEU
1	B	151	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	105	GLN
1	C	202	ASN
1	C	3	GLN
1	A	202	ASN
1	B	202	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	PCA	C	1	1	3,4,9	0.62	0	0,4,12	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	C	1	1	-	0/0/2/13	0/0/0/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	1	PCA	1	0

## 5.5 Carbohydrates ⓘ

6 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GLC	B	248	2	12,12,12	0.37	0	17,17,17	0.59	0
2	MAN	B	249	2	11,11,12	0.63	0	14,15,17	0.43	0
2	SGA	B	250	2	15,15,16	0.53	0	18,22,24	1.16	1 (5%)
2	GLC	C	248	2	12,12,12	0.31	0	17,17,17	0.65	0
2	MAN	C	249	2	11,11,12	0.50	0	14,15,17	0.83	1 (7%)
2	SGA	C	250	2	15,15,16	0.87	1 (6%)	18,22,24	1.15	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	248	2	-	0/2/22/22	0/1/1/1
2	MAN	B	249	2	-	0/2/19/22	0/1/1/1
2	SGA	B	250	2	-	0/7/24/27	0/1/1/1
2	GLC	C	248	2	-	0/2/22/22	0/1/1/1
2	MAN	C	249	2	-	0/2/19/22	0/1/1/1
2	SGA	C	250	2	-	0/7/24/27	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	250	SGA	O3-S	-2.18	1.50	1.57

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	250	SGA	C3-O3-S	-2.87	113.31	118.77
2	C	249	MAN	C1-O5-C5	2.06	114.86	112.25
2	C	250	SGA	O3-S-O2S	2.07	113.58	106.86
2	B	250	SGA	C1-C2-C3	3.58	113.63	109.24

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	248	GLC	4	0
2	B	249	MAN	1	0
2	B	250	SGA	1	0
2	C	248	GLC	1	0
2	C	250	SGA	1	0

## 5.6 Ligand geometry

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	ARC	A	1000	-	20,20,20	0.26	0	22,23,23	0.34	0
5	ARC	A	1001	-	20,20,20	0.22	0	22,23,23	0.38	0
5	ARC	A	1002	-	20,20,20	0.38	0	22,23,23	0.36	0
7	GOL	A	1003	-	2,2,5	0.48	0	1,1,5	0.62	0
4	RET	A	999	-	19,20,21	1.65	3 (15%)	27,27,28	1.64	4 (14%)
5	ARC	B	1000	-	20,20,20	0.22	0	22,23,23	0.33	0
5	ARC	B	1001	-	20,20,20	0.31	0	22,23,23	0.35	0
5	ARC	B	1002	-	20,20,20	0.35	0	22,23,23	0.35	0
5	ARC	B	1003	-	20,20,20	0.19	0	22,23,23	0.40	0
6	OCT	B	1004	-	7,7,7	0.24	0	6,6,6	0.52	0
7	GOL	B	1005	-	3,3,5	0.42	0	2,2,5	0.29	0
3	BGC	B	251	-	12,12,12	0.27	0	17,17,17	0.44	0
4	RET	B	999	-	19,20,21	1.65	4 (21%)	27,27,28	1.57	5 (18%)
5	ARC	C	1000	-	20,20,20	0.24	0	22,23,23	0.31	0
5	ARC	C	1001	-	20,20,20	0.28	0	22,23,23	0.28	0
5	ARC	C	1002	-	20,20,20	0.20	0	22,23,23	0.43	0
7	GOL	C	1003	-	2,2,5	0.56	0	1,1,5	0.35	0
4	RET	C	999	-	19,20,21	1.53	2 (10%)	27,27,28	1.84	7 (25%)



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ARC	A	1000	-	-	0/21/21/21	0/0/0/0
5	ARC	A	1001	-	-	0/21/21/21	0/0/0/0
5	ARC	A	1002	-	-	0/21/21/21	0/0/0/0
7	GOL	A	1003	-	-	0/0/0/4	0/0/0/0
4	RET	A	999	-	-	0/13/30/31	0/1/1/1
5	ARC	B	1000	-	-	0/21/21/21	0/0/0/0
5	ARC	B	1001	-	-	0/21/21/21	0/0/0/0
5	ARC	B	1002	-	-	0/21/21/21	0/0/0/0
5	ARC	B	1003	-	-	0/21/21/21	0/0/0/0
6	OCT	B	1004	-	-	0/5/5/5	0/0/0/0
7	GOL	B	1005	-	-	0/1/1/4	0/0/0/0
3	BGC	B	251	-	-	0/2/22/22	0/1/1/1
4	RET	B	999	-	-	0/13/30/31	0/1/1/1
5	ARC	C	1000	-	-	0/21/21/21	0/0/0/0
5	ARC	C	1001	-	-	0/21/21/21	0/0/0/0
5	ARC	C	1002	-	-	0/21/21/21	0/0/0/0
7	GOL	C	1003	-	-	0/0/0/4	0/0/0/0
4	RET	C	999	-	-	0/13/30/31	0/1/1/1

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	999	RET	C20-C13	-2.96	1.44	1.50
4	C	999	RET	C20-C13	-2.93	1.44	1.50
4	B	999	RET	C20-C13	-2.75	1.45	1.50
4	B	999	RET	C12-C13	-2.19	1.41	1.45
4	A	999	RET	C5-C6	3.20	1.39	1.34

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	999	RET	C18-C5-C6	-5.79	118.92	124.61
4	A	999	RET	C18-C5-C6	-5.07	119.63	124.61
4	B	999	RET	C18-C5-C6	-4.60	120.09	124.61
4	C	999	RET	C10-C11-C12	-3.08	113.75	123.13
4	A	999	RET	C8-C7-C6	-2.52	119.75	127.32

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

15 monomers are involved in 97 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1000	ARC	11	0
5	A	1001	ARC	11	0
5	A	1002	ARC	6	0
4	A	999	RET	6	0
5	B	1000	ARC	6	0
5	B	1001	ARC	5	0
5	B	1002	ARC	13	0
5	B	1003	ARC	6	0
6	B	1004	OCT	1	0
3	B	251	BGC	1	0
4	B	999	RET	6	0
5	C	1000	ARC	3	0
5	C	1001	ARC	9	0
5	C	1002	ARC	15	0
4	C	999	RET	6	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	230/247 (93%)	0.74	33 (14%) 4 2	33, 72, 94, 100	0
1	B	231/247 (93%)	0.42	21 (9%) 11 7	27, 49, 79, 100	0
1	C	231/247 (93%)	0.30	19 (8%) 14 9	28, 46, 74, 93	0
All	All	692/741 (93%)	0.49	73 (10%) 8 5	27, 54, 90, 100	0

The worst 5 of 73 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2	ALA	11.7
1	A	3	GLN	8.5
1	C	4	ILE	7.2
1	B	3	GLN	7.1
1	C	2	ALA	6.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	PCA	C	1	5/9	0.77	0.71	-	91,92,92,92	0

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	MAN	C	249	11/12	0.77	0.23	0.20	71,73,77,81	0
2	SGA	C	250	15/16	0.86	0.31	-0.17	80,82,83,83	0
2	MAN	B	249	11/12	0.76	0.18	-0.45	69,72,80,86	0
2	SGA	B	250	15/16	0.76	0.41	-	90,91,96,97	0
2	GLC	B	248	12/12	0.91	0.12	-	68,69,69,69	0
2	GLC	C	248	12/12	0.93	0.13	-	59,62,66,69	0

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	ARC	A	1001	21/21	0.68	0.55	9.28	69,82,87,87	0
5	ARC	C	1002	21/21	0.84	0.48	8.39	62,63,64,65	0
5	ARC	C	1001	21/21	0.58	0.56	7.14	85,92,97,98	0
5	ARC	C	1000	21/21	0.75	0.41	6.22	70,76,77,78	0
5	ARC	B	1001	21/21	0.84	0.35	5.29	48,57,69,71	0
5	ARC	B	1003	21/21	0.73	0.47	5.03	62,66,70,71	0
5	ARC	A	1002	21/21	0.87	0.36	4.35	56,63,70,70	0
5	ARC	B	1000	21/21	0.68	0.50	4.30	86,89,92,93	0
5	ARC	A	1000	21/21	0.68	0.51	4.12	84,86,92,92	0
5	ARC	B	1002	21/21	0.73	0.37	3.34	69,75,78,78	0
6	OCT	B	1004	8/8	0.74	0.33	3.23	55,60,67,69	0
4	RET	A	999	20/21	0.82	0.32	2.24	62,65,71,71	0
4	RET	B	999	20/21	0.92	0.21	0.65	30,34,49,49	0
4	RET	C	999	20/21	0.93	0.20	0.34	24,35,42,43	0
7	GOL	B	1005	4/6	0.76	0.25	-	71,71,72,73	0
7	GOL	C	1003	3/6	0.85	0.25	-	66,66,66,67	0
7	GOL	A	1003	3/6	0.92	0.11	-	69,69,70,70	0
3	BGC	B	251	12/12	0.80	0.51	-	71,74,75,75	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.