



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 10:07 PM GMT

PDB ID : 5BR0
Title : Crystal structure of hemagglutinin of A/Taiwan/2/2013 (H6N1)
Authors : Ni, F.; Kondrashkina, E.; Wang, Q.
Deposited on : 2015-05-29
Resolution : 2.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	326	<div> <div>3%</div> <div>90%</div> <div>10%</div> </div>
1	C	326	<div> <div>3%</div> <div>90%</div> <div>10%</div> </div>
2	B	171	<div> <div>2%</div> <div>96%</div> <div>.</div> </div>
2	D	171	<div> <div>2%</div> <div>96%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	B	301	-	-	-	X
4	FUC	B	302	X	-	-	-
4	FUC	D	302	X	-	-	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 8975 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called HEMAGGLUTININ HA1 CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	326	Total	C	N	O	S	0	0	0
			2577	1634	439	491	13			
1	C	326	Total	C	N	O	S	0	0	0
			2577	1634	439	491	13			

- Molecule 2 is a protein called HEMAGGLUTININ HA2 CHAIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	171	Total	C	N	O	S	0	0	0
			1381	860	242	272	7			
2	D	171	Total	C	N	O	S	0	0	0
			1381	860	242	272	7			

- Molecule 3 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	C	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		
3	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is ALPHA-L-FUCOSE (three-letter code: FUC) (formula: C₆H₁₂O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			10	6	4		
4	D	1	Total	C	O	0	0
			10	6	4		

- Molecule 5 is BETA-D-MANNOSE (three-letter code: BMA) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			11	6	5		
5	D	1	Total	C	O	0	0
			11	6	5		

- Molecule 6 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			11	6	5		
6	B	1	Total	C	O	0	0
			11	6	5		
6	D	1	Total	C	O	0	0
			11	6	5		
6	D	1	Total	C	O	0	0
			11	6	5		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	243	Total	O	0	0
			243	243		
7	B	166	Total	O	0	0
			166	166		
7	C	216	Total	O	0	0
			216	216		
7	D	180	Total	O	0	0
			180	180		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HEMAGGLUTININ HA1 CHAIN




- Molecule 1: HEMAGGLUTININ HA1 CHAIN



- Molecule 2: HEMAGGLUTININ HA2 CHAIN



- Molecule 2: HEMAGGLUTININ HA2 CHAIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 3	Depositor
Cell constants a, b, c, α , β , γ	114.02Å 114.02Å 164.44Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	53.87 – 2.39 63.18 – 2.39	Depositor EDS
% Data completeness (in resolution range)	100.0 (53.87-2.39) 100.0 (63.18-2.39)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.99 (at 2.40Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.182 , 0.210 0.189 , 0.214	Depositor DCC
R_{free} test set	4748 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	30.6	Xtriage
Anisotropy	0.425	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 52.5	EDS
Estimated twinning fraction	0.460 for -h,-k,l 0.038 for h,-h-k,-l 0.038 for -k,-h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 94734 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8975	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.26	0/2638	0.44	0/3589
1	C	0.27	0/2638	0.44	0/3589
2	B	0.29	0/1409	0.43	0/1897
2	D	0.28	0/1409	0.44	0/1897
All	All	0.27	0/8094	0.44	0/10972

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2577	0	2527	26	0
1	C	2577	0	2527	20	0
2	B	1381	0	1294	6	0
2	D	1381	0	1294	5	0
3	A	56	0	52	0	0
3	B	28	0	23	0	0
3	C	56	0	52	1	0
3	D	28	0	23	0	0
4	B	10	0	10	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	10	0	10	0	0
5	B	11	0	8	0	0
5	D	11	0	8	0	0
6	B	22	0	20	0	0
6	D	22	0	20	0	0
7	A	243	0	0	12	0
7	B	166	0	0	4	2
7	C	216	0	0	8	2
7	D	180	0	0	3	3
All	All	8975	0	7868	56	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:163:GLN:OE1	7:B:401:HOH:O	1.91	0.89
1:C:264:LYS:NZ	7:C:502:HOH:O	2.07	0.87
1:A:208:ASN:OD1	7:A:501:HOH:O	1.97	0.81
1:C:21:GLU:OE1	7:C:501:HOH:O	1.98	0.81
1:A:151:ILE:O	7:A:502:HOH:O	2.02	0.75
1:A:205:GLU:OE2	7:A:503:HOH:O	2.08	0.71
3:C:402:NAG:O4	7:C:503:HOH:O	2.07	0.71
1:C:279:ALA:O	7:C:504:HOH:O	2.09	0.70
1:A:21:GLU:OE2	7:A:505:HOH:O	2.12	0.68
1:A:21:GLU:OE1	7:A:504:HOH:O	2.11	0.67
1:C:37:LYS:NZ	7:C:507:HOH:O	2.26	0.66
2:D:106:ARG:HD2	7:D:410:HOH:O	1.97	0.64
2:B:48:ILE:HD11	2:B:110:LEU:HD23	1.87	0.56
2:D:48:ILE:HD11	2:D:110:LEU:HD23	1.88	0.56
1:C:149:VAL:HG23	1:C:253:TRP:HB2	1.88	0.55
1:C:207:MET:HG2	1:C:208:ASN:N	2.21	0.55
2:D:116:LYS:HE3	7:D:433:HOH:O	2.07	0.55
1:A:321:ARG:NH2	7:A:507:HOH:O	2.17	0.54
1:A:207:MET:HG2	1:A:208:ASN:N	2.23	0.53
2:B:82:LYS:NZ	7:B:410:HOH:O	2.42	0.51
1:A:239:GLU:OE2	7:A:508:HOH:O	2.19	0.51
1:A:123:ALA:HB3	7:A:517:HOH:O	2.09	0.51
1:C:2:LYS:HG2	2:D:139:GLU:HG2	1.93	0.51
1:A:149:VAL:HG23	1:A:253:TRP:HB2	1.93	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:43:LYS:HG2	1:C:48:ALA:HA	1.95	0.49
1:A:87:ASN:ND2	7:A:519:HOH:O	2.47	0.48
1:A:218:ARG:NE	7:A:518:HOH:O	2.46	0.48
2:B:121:LYS:NZ	7:B:404:HOH:O	2.35	0.48
1:A:45:MET:N	7:A:506:HOH:O	2.47	0.47
1:A:283:THR:HG22	1:A:301:TRP:HB3	1.96	0.47
1:A:65:PRO:HG3	1:A:144:PHE:O	2.15	0.47
1:C:325:GLN:O	1:C:326:ILE:HG13	2.15	0.47
1:A:137:SER:OG	1:A:138:TYR:N	2.49	0.45
1:A:284:ILE:HG23	1:A:285:THR:HG23	1.99	0.45
1:C:64:ASN:HA	1:C:65:PRO:HD3	1.86	0.45
1:C:120:SER:OG	7:C:505:HOH:O	2.21	0.45
1:C:283:THR:HG22	1:C:301:TRP:HB3	1.98	0.44
1:C:146:ARG:HD3	7:C:551:HOH:O	2.17	0.44
1:A:2:LYS:HG2	2:B:139:GLU:HG2	1.98	0.44
1:C:137:SER:OG	1:C:138:TYR:N	2.50	0.44
1:A:130:GLY:HA3	1:A:150:TRP:HB3	2.00	0.43
1:A:57:ILE:O	1:A:61:ILE:HG13	2.19	0.43
1:A:82:ARG:HA	1:A:83:PRO:HD3	1.84	0.43
1:A:61:ILE:H	1:A:61:ILE:HG13	1.64	0.43
1:A:1:ASP:HB2	7:B:458:HOH:O	2.19	0.43
1:C:41:PHE:CE2	1:C:272:LEU:HB2	2.54	0.43
1:C:57:ILE:O	1:C:61:ILE:HG13	2.20	0.42
1:C:130:GLY:HA3	1:C:150:TRP:HB3	2.01	0.42
2:B:58:LYS:HD3	2:B:58:LYS:HA	1.85	0.42
1:A:43:LYS:HG2	1:A:48:ALA:HA	2.00	0.42
1:C:123:ALA:HB3	7:C:542:HOH:O	2.19	0.42
1:C:288:LEU:HD21	1:C:297:VAL:HG21	2.01	0.42
1:C:65:PRO:HG3	1:C:144:PHE:O	2.19	0.42
1:A:130:GLY:N	7:A:526:HOH:O	2.52	0.42
1:A:41:PHE:CE2	1:A:272:LEU:HB2	2.55	0.41
2:D:57:ASN:ND2	7:D:408:HOH:O	2.47	0.41

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:B:559:HOH:O	7:C:654:HOH:O[1_554]	1.89	0.31
7:B:510:HOH:O	7:B:523:HOH:O[3_565]	1.97	0.23
7:D:498:HOH:O	7:D:514:HOH:O[2_555]	2.01	0.19

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:C:694:HOH:O	7:D:558:HOH:O[2_555]	2.02	0.18
7:D:451:HOH:O	7:D:474:HOH:O[2_555]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	324/326 (99%)	318 (98%)	6 (2%)	0	100	100
1	C	324/326 (99%)	317 (98%)	7 (2%)	0	100	100
2	B	169/171 (99%)	167 (99%)	2 (1%)	0	100	100
2	D	169/171 (99%)	166 (98%)	3 (2%)	0	100	100
All	All	986/994 (99%)	968 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/287 (100%)	286 (100%)	1 (0%)	94	98
1	C	287/287 (100%)	287 (100%)	0	100	100
2	B	146/146 (100%)	146 (100%)	0	100	100
2	D	146/146 (100%)	146 (100%)	0	100	100
All	All	866/866 (100%)	865 (100%)	1 (0%)	95	99

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	159	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

20 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	401	1	14,14,15	0.58	0	15,19,21	0.61	0
3	NAG	A	402	1	14,14,15	0.61	0	15,19,21	0.84	1 (6%)
3	NAG	A	403	1	14,14,15	0.42	0	15,19,21	0.64	1 (6%)
3	NAG	A	404	1	14,14,15	1.06	1 (7%)	15,19,21	1.13	2 (13%)
3	NAG	B	301	3,2,4	14,14,15	0.53	0	15,19,21	0.44	0
4	FUC	B	302	3	10,10,11	1.06	0	13,14,16	0.85	0
3	NAG	B	303	3,5	14,14,15	0.15	0	15,19,21	0.53	0
5	BMA	B	304	3,6	11,11,12	0.89	0	15,15,17	0.73	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	MAN	B	305	5	11,11,12	0.71	0	15,15,17	1.00	2 (13%)
6	MAN	B	306	5	11,11,12	0.75	0	15,15,17	1.40	3 (20%)
3	NAG	C	401	1	14,14,15	0.32	0	15,19,21	0.19	0
3	NAG	C	402	1	14,14,15	0.46	0	15,19,21	0.45	0
3	NAG	C	403	1	14,14,15	0.22	0	15,19,21	0.58	1 (6%)
3	NAG	C	404	1	14,14,15	0.22	0	15,19,21	0.30	0
3	NAG	D	301	3,2,4	14,14,15	0.45	0	15,19,21	0.48	0
4	FUC	D	302	3	10,10,11	0.63	0	13,14,16	0.83	0
3	NAG	D	303	3,5	14,14,15	0.20	0	15,19,21	0.34	0
5	BMA	D	304	3,6	11,11,12	0.61	0	15,15,17	0.83	0
6	MAN	D	305	5	11,11,12	0.72	0	15,15,17	0.94	2 (13%)
6	MAN	D	306	5	11,11,12	0.75	0	15,15,17	1.04	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	401	1	-	0/6/23/26	0/1/1/1
3	NAG	A	402	1	-	0/6/23/26	0/1/1/1
3	NAG	A	403	1	-	0/6/23/26	0/1/1/1
3	NAG	A	404	1	-	0/6/23/26	0/1/1/1
3	NAG	B	301	3,2,4	-	0/6/23/26	0/1/1/1
4	FUC	B	302	3	1/1/4/5	0/0/17/20	0/1/1/1
3	NAG	B	303	3,5	-	0/6/23/26	0/1/1/1
5	BMA	B	304	3,6	-	0/2/19/22	0/1/1/1
6	MAN	B	305	5	-	0/2/19/22	0/1/1/1
6	MAN	B	306	5	-	0/2/19/22	0/1/1/1
3	NAG	C	401	1	-	0/6/23/26	0/1/1/1
3	NAG	C	402	1	-	0/6/23/26	0/1/1/1
3	NAG	C	403	1	-	0/6/23/26	0/1/1/1
3	NAG	C	404	1	-	0/6/23/26	0/1/1/1
3	NAG	D	301	3,2,4	-	0/6/23/26	0/1/1/1
4	FUC	D	302	3	1/1/4/5	0/0/17/20	0/1/1/1
3	NAG	D	303	3,5	-	0/6/23/26	0/1/1/1
5	BMA	D	304	3,6	-	0/2/19/22	0/1/1/1
6	MAN	D	305	5	-	0/2/19/22	0/1/1/1
6	MAN	D	306	5	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	404	NAG	C1-C2	3.50	1.57	1.52

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	NAG	C1-O5-C5	-2.46	108.53	112.14
6	D	305	MAN	O2-C2-C3	-2.21	105.72	110.19
3	A	404	NAG	O5-C5-C4	-2.17	106.53	110.13
6	B	305	MAN	O2-C2-C3	-2.09	105.98	110.19
6	D	306	MAN	O2-C2-C3	-2.04	106.08	110.19
6	B	306	MAN	O2-C2-C3	-2.01	106.14	110.19
6	D	305	MAN	C1-O5-C5	2.03	115.12	112.14
3	C	403	NAG	C1-O5-C5	2.10	115.23	112.14
3	A	403	NAG	C1-O5-C5	2.12	115.26	112.14
6	B	305	MAN	C1-O5-C5	2.17	115.33	112.14
6	D	306	MAN	C1-O5-C5	2.25	115.44	112.14
6	B	306	MAN	O5-C1-C2	2.27	114.53	110.89
3	A	404	NAG	C1-O5-C5	3.02	116.58	112.14
6	B	306	MAN	C1-O5-C5	3.82	117.76	112.14

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	B	302	FUC	C1
4	D	302	FUC	C1

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	402	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	326/326 (100%)	-0.10	11 (3%) 49 49	20, 38, 60, 96	0
1	C	326/326 (100%)	-0.09	9 (2%) 56 55	19, 38, 60, 90	0
2	B	171/171 (100%)	-0.08	3 (1%) 71 71	16, 28, 48, 77	0
2	D	171/171 (100%)	-0.11	3 (1%) 71 71	16, 28, 49, 80	0
All	All	994/994 (100%)	-0.09	26 (2%) 59 58	16, 34, 58, 96	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	156	SER	6.1
1	C	156	SER	4.6
1	A	326	ILE	4.4
2	B	168	LEU	3.5
1	C	155	ASP	3.1
1	C	190	ASP	3.0
1	C	262	ASN	2.9
1	A	190	ASP	2.8
1	C	188	VAL	2.8
1	A	187	THR	2.7
1	A	123	ALA	2.7
2	D	171	GLN	2.7
2	D	168	LEU	2.7
1	C	191	ASN	2.6
2	B	171	GLN	2.6
1	A	191	ASN	2.6
1	C	123	ALA	2.6
1	A	188	VAL	2.4
1	A	155	ASP	2.3
1	A	158	THR	2.3
1	A	262	ASN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	195	SER	2.2
2	D	167	LYS	2.2
1	C	187	THR	2.1
2	B	167	LYS	2.0
1	C	153	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NAG	B	301	14/15	0.91	0.18	2.14	51,62,69,72	0
3	NAG	D	301	14/15	0.90	0.15	1.34	53,62,70,72	0
4	FUC	B	302	10/11	0.73	0.33	-	76,91,102,107	0
3	NAG	C	401	14/15	0.74	0.41	-	44,48,51,53	0
5	BMA	B	304	11/12	0.90	0.24	-	88,100,105,106	0
3	NAG	A	404	14/15	0.75	0.52	-	57,58,58,58	0
6	MAN	B	306	11/12	0.56	0.54	-	105,117,129,130	0
3	NAG	D	303	14/15	0.94	0.21	-	63,77,80,80	0
3	NAG	B	303	14/15	0.94	0.19	-	64,73,77,80	0
4	FUC	D	302	10/11	0.80	0.22	-	78,93,100,101	0
5	BMA	D	304	11/12	0.87	0.24	-	83,103,107,111	0
3	NAG	C	403	14/15	0.89	0.26	-	51,53,53,53	0
3	NAG	C	404	14/15	0.71	0.54	-	56,56,57,57	0
3	NAG	A	403	14/15	0.82	0.25	-	50,53,53,53	0
6	MAN	D	305	11/12	0.84	0.44	-	91,96,109,111	0
3	NAG	A	402	14/15	0.72	0.47	-	44,47,47,47	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NAG	A	401	14/15	0.63	0.36	-	46,49,52,53	0
6	MAN	D	306	11/12	0.51	0.64	-	108,126,130,134	0
6	MAN	B	305	11/12	0.74	0.41	-	87,99,106,109	0
3	NAG	C	402	14/15	0.71	0.41	-	46,49,51,51	0

6.5 Other polymers [i](#)

There are no such residues in this entry.