



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:11 PM GMT

PDB ID : 4BSA
Title : Crystal Structure of the Haemagglutinin (with Asn-133 Glycosylation) from an H7N9 Influenza Virus Isolated from Humans
Authors : Xiong, X.; Haire, L.F.; Martin, S.R.; Wharton, S.A.; Daniels, R.S.; Bennett, M.S.; McCauley, J.W.; Collins, P.J.; Walker, P.A.; Skehel, J.J.; Gamblin, S.J.
Deposited on : 2013-06-10
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	321	<div> <div>16%</div> <div>87%</div> <div>11%</div> <div>..</div> </div>
2	B	177	<div> <div>18%</div> <div>89%</div> <div>7%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SO4	A	1321	-	-	-	X
4	SO4	A	1326	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4216 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HEMAGGLUTININ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	317	2416	1500	437	464	15	0	0	1


There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	10	LEU	VAL	SEE REMARK 999	UNP M4YV75
A	125	THR	ALA	SEE REMARK 999	UNP M4YV75
A	217	LEU	ILE	SEE REMARK 999	UNP M4YV75

- Molecule 2 is a protein called HEMAGGLUTININ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	170	1379	851	241	280	7	0	0	0

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0

- Molecule 5 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	2	Total C N O 28 16 2 10	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	200	Total O 200 200	0	0
6	B	67	Total O 67 67	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: HEMAGGLUTININ



• Molecule 2: HEMAGGLUTININ



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	116.13Å 116.13Å 295.51Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	58.13 – 2.30 58.07 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.9 (58.13-2.30) 98.9 (58.07-2.30)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.19 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.250 , 0.296 0.251 , 0.297	Depositor DCC
R_{free} test set	1725 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	34.7	Xtriage
Anisotropy	0.022	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.7	EDS
Estimated twinning fraction	0.008 for -1/3*h+1/3*k+1/3*l,-k,8/3*h+4/3*k+1/3*l 0.018 for -2/3*h-1/3*k-1/3*l,-1/3*h-2/3*k+1/3*l,-4/3*h+4/3*k+1/3*l 0.008 for -h,1/3*h-1/3*k-1/3*l,-4/3*h-8/3*k+1/3*l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 34129 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4216	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/2462	0.58	0/3328
2	B	0.32	0/1403	0.53	0/1890
All	All	0.34	0/3865	0.56	0/5218

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2416	0	2374	31	0
2	B	1379	0	1280	7	0
3	A	56	0	52	0	0
4	A	50	0	0	0	0
4	B	20	0	0	0	0
5	B	28	0	25	2	0
6	A	200	0	0	13	1
6	B	67	0	0	1	1
All	All	4216	0	3731	37	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:211:ARG:HD3	6:A:2113:HOH:O	1.95	0.66
1:A:38:ILE:HD12	1:A:280:ILE:HD12	1.78	0.63
1:A:121:ARG:NH1	1:A:146:ASN:O	2.32	0.63
1:A:222:ASP:HB3	6:A:2128:HOH:O	2.00	0.61
1:A:173:ILE:HD11	6:A:2110:HOH:O	2.03	0.59
1:A:298:ARG:HG2	2:B:92:TRP:CE2	2.40	0.56
1:A:242:PHE:CD2	6:A:2136:HOH:O	2.53	0.55
1:A:101:ILE:HD13	1:A:227:MET:HE3	1.88	0.54
1:A:170:VAL:O	1:A:245:PRO:HB3	2.07	0.54
1:A:298:ARG:HG2	2:B:92:TRP:CD2	2.46	0.51
5:B:201:NAG:H83	6:B:2027:HOH:O	2.09	0.51
1:A:185:LEU:HA	6:A:2089:HOH:O	2.11	0.50
1:A:66:CYS:O	1:A:70:LEU:HD13	2.11	0.50
1:A:235:THR:HG23	6:A:2105:HOH:O	2.12	0.50
1:A:226:LEU:HD22	1:A:226:LEU:C	2.32	0.50
1:A:131:ARG:NH1	1:A:136:PHE:O	2.45	0.49
1:A:193:VAL:HB	6:A:2110:HOH:O	2.12	0.49
1:A:175:HIS:HA	6:A:2111:HOH:O	2.12	0.48
2:B:169:ASN:O	2:B:170:ARG:C	2.51	0.48
1:A:173:ILE:HB	1:A:222:ASP:HB2	1.97	0.46
1:A:224:HIS:CE1	6:A:2128:HOH:O	2.69	0.46
2:B:6:ILE:HD12	2:B:112:ASP:HA	1.97	0.46
1:A:38:ILE:CD1	1:A:280:ILE:HD12	2.45	0.45
1:A:117:TYR:CE1	1:A:143:LEU:HD11	2.53	0.44
2:B:75:LYS:HE3	5:B:201:NAG:H81	2.00	0.44
1:A:209:GLY:O	1:A:211:ARG:NH1	2.49	0.44
1:A:158:SER:N	6:A:2106:HOH:O	2.51	0.43
1:A:123:ASN:N	1:A:123:ASN:OD1	2.51	0.43
1:A:175:HIS:NE2	1:A:222:ASP:OD1	2.52	0.42
1:A:60:ILE:HG21	1:A:170:VAL:HG21	2.00	0.42
2:B:19:ASP:HB3	2:B:36:ALA:HB2	2.02	0.42
2:B:55:LEU:HD22	2:B:99:LEU:HD21	2.01	0.41
1:A:193:VAL:HA	1:A:237:SER:O	2.21	0.41
1:A:143:LEU:HD12	6:A:2136:HOH:O	2.21	0.41
1:A:176:SER:HB3	1:A:181:GLU:HB2	2.03	0.41
1:A:158:SER:HB2	6:A:2105:HOH:O	2.21	0.40
1:A:153:PRO:HG2	6:A:2103:HOH:O	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:2029:HOH:O	6:A:2031:HOH:O[6_555]	1.84	0.36
6:B:2044:HOH:O	6:B:2046:HOH:O[3_655]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	315/321 (98%)	300 (95%)	15 (5%)	0	100	100
2	B	168/177 (95%)	164 (98%)	4 (2%)	0	100	100
All	All	483/498 (97%)	464 (96%)	19 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	266/270 (98%)	256 (96%)	10 (4%)	40	54
2	B	145/152 (95%)	141 (97%)	4 (3%)	51	68
All	All	411/422 (97%)	397 (97%)	14 (3%)	44	59

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	HIS
1	A	70	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	94	ASN
1	A	120	ILE
1	A	127	SER
1	A	155	MET
1	A	186	TYR
1	A	222	ASP
1	A	226	LEU
1	A	298	ARG
2	B	34	THR
2	B	64	GLU
2	B	144	CYS
2	B	170	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	146	ASN
1	A	174	HIS
1	A	201	GLN
2	B	62	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	B	201	2,5	14,14,15	0.50	0	15,19,21	0.81	0
5	NAG	B	202	5	14,14,15	0.53	0	15,19,21	0.82	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	B	201	2,5	-	0/6/23/26	0/1/1/1
5	NAG	B	202	5	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	201	NAG	2	0

5.6 Ligand geometry

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	A	1123	1	14,14,15	0.48	0	15,19,21	1.47	1 (6%)
4	SO4	A	1317	-	4,4,4	0.39	0	6,6,6	0.07	0
4	SO4	A	1318	-	4,4,4	0.37	0	6,6,6	0.09	0
4	SO4	A	1319	-	4,4,4	0.37	0	6,6,6	0.08	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	A	1320	-	4,4,4	0.36	0	6,6,6	0.09	0
4	SO4	A	1321	-	4,4,4	0.42	0	6,6,6	0.09	0
4	SO4	A	1322	-	4,4,4	0.37	0	6,6,6	0.13	0
4	SO4	A	1323	-	4,4,4	0.37	0	6,6,6	0.08	0
4	SO4	A	1324	-	4,4,4	0.38	0	6,6,6	0.18	0
4	SO4	A	1325	-	4,4,4	0.38	0	6,6,6	0.14	0
4	SO4	A	1326	-	4,4,4	0.35	0	6,6,6	0.12	0
3	NAG	A	403	1	14,14,15	0.44	0	15,19,21	0.96	1 (6%)
3	NAG	A	404	1	14,14,15	0.47	0	15,19,21	0.94	1 (6%)
3	NAG	A	405	1	14,14,15	0.35	0	15,19,21	1.94	1 (6%)
4	SO4	B	1171	-	4,4,4	0.42	0	6,6,6	0.11	0
4	SO4	B	1172	-	4,4,4	0.41	0	6,6,6	0.12	0
4	SO4	B	1173	-	4,4,4	0.42	0	6,6,6	0.16	0
4	SO4	B	1174	-	4,4,4	0.34	0	6,6,6	0.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1123	1	-	0/6/23/26	0/1/1/1
4	SO4	A	1317	-	-	0/0/0/0	0/0/0/0
4	SO4	A	1318	-	-	0/0/0/0	0/0/0/0
4	SO4	A	1319	-	-	0/0/0/0	0/0/0/0
4	SO4	A	1320	-	-	0/0/0/0	0/0/0/0
4	SO4	A	1321	-	-	0/0/0/0	0/0/0/0
4	SO4	A	1322	-	-	0/0/0/0	0/0/0/0
4	SO4	A	1323	-	-	0/0/0/0	0/0/0/0
4	SO4	A	1324	-	-	0/0/0/0	0/0/0/0
4	SO4	A	1325	-	-	0/0/0/0	0/0/0/0
4	SO4	A	1326	-	-	0/0/0/0	0/0/0/0
3	NAG	A	403	1	-	0/6/23/26	0/1/1/1
3	NAG	A	404	1	-	0/6/23/26	0/1/1/1
3	NAG	A	405	1	-	0/6/23/26	0/1/1/1
4	SO4	B	1171	-	-	0/0/0/0	0/0/0/0
4	SO4	B	1172	-	-	0/0/0/0	0/0/0/0
4	SO4	B	1173	-	-	0/0/0/0	0/0/0/0
4	SO4	B	1174	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	A	404	NAG	O7-C7-C8	-2.22	117.98	122.06
3	A	403	NAG	C1-O5-C5	2.54	115.48	112.25
3	A	1123	NAG	C1-O5-C5	5.07	118.68	112.25
3	A	405	NAG	C1-O5-C5	6.70	120.75	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	317/321 (98%)	1.01	51 (16%) 3 4	20, 47, 125, 152	0
2	B	170/177 (96%)	1.08	31 (18%) 2 3	18, 61, 102, 110	0
All	All	487/498 (97%)	1.04	82 (16%) 2 3	18, 50, 118, 152	0

All (82) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	189	GLY	7.7
2	B	141	PHE	7.3
1	A	190	ASN	6.4
1	A	192	LEU	5.9
2	B	168	GLN	5.4
1	A	205	VAL	5.0
2	B	166	ALA	5.0
1	A	187	GLY	4.9
1	A	204	PHE	4.8
1	A	198	SER	4.6
2	B	164	GLU	4.5
1	A	241	ALA	4.5
1	A	194	THR	4.4
2	B	161	LYS	4.4
2	B	156	THR	4.4
1	A	193	VAL	4.2
2	B	167	MET	4.1
1	A	151	ALA	3.9
2	B	143	LYS	3.9
1	A	196	GLY	3.9
1	A	147	THR	3.8
1	A	216	GLY	3.7
1	A	236	PHE	3.6
1	A	242	PHE	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	162	TYR	3.5
2	B	59	THR	3.4
1	A	129	CYS	3.4
1	A	195	VAL	3.3
2	B	160	SER	3.2
1	A	186	TYR	3.2
1	A	212	PRO	3.1
1	A	239	ASN	3.1
2	B	147	ASP	3.1
2	B	29	ALA	3.1
1	A	122	THR	3.1
1	A	177	VAL	3.0
2	B	142	HIS	3.0
2	B	145	ASP	3.0
1	A	203	SER	2.9
2	B	165	GLU	2.9
1	A	183	THR	2.9
2	B	56	ILE	2.8
2	B	170	ARG	2.8
1	A	200	TYR	2.8
1	A	238	PHE	2.7
2	B	27	GLN	2.7
1	A	211	ARG	2.7
1	A	143	LEU	2.7
2	B	17	LEU	2.7
1	A	235	THR	2.7
1	A	215	ASN	2.6
1	A	214	VAL	2.6
2	B	163	ARG	2.6
2	B	159	HIS	2.5
1	A	121	ARG	2.5
1	A	191	LYS	2.5
1	A	149	ASN	2.5
1	A	152	PHE	2.5
1	A	182	GLN	2.5
1	A	118	SER	2.4
1	A	210	ALA	2.4
1	A	155	MET	2.4
1	A	120	ILE	2.4
1	A	234	VAL	2.4
1	A	197	SER	2.3
1	A	156	THR	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	237	SER	2.3
2	B	58	LYS	2.3
1	A	206	PRO	2.3
2	B	139	GLU	2.3
2	B	11	GLU	2.3
2	B	152	ILE	2.2
2	B	144	CYS	2.2
2	B	31	GLY	2.2
1	A	207	SER	2.2
1	A	154	GLN	2.2
1	A	125	THR	2.1
1	A	201	GLN	2.1
2	B	137	CYS	2.1
2	B	80	VAL	2.1
2	B	9	PHE	2.1
1	A	185	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	NAG	B	201	14/15	0.88	0.14	-1.96	38,46,53,56	0
5	NAG	B	202	14/15	0.77	0.22	-	58,64,68,69	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy

less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	SO4	A	1321	5/5	0.80	0.29	3.41	83,84,90,92	0
4	SO4	A	1326	5/5	0.81	0.46	3.14	99,101,103,105	0
4	SO4	B	1173	5/5	0.84	0.26	1.30	72,72,75,76	0
4	SO4	B	1171	5/5	0.86	0.21	-0.32	86,87,92,92	0
4	SO4	A	1320	5/5	0.74	0.37	-	112,116,119,119	0
4	SO4	A	1323	5/5	0.69	0.33	-	97,105,105,107	0
3	NAG	A	405	14/15	0.68	0.36	-	59,70,73,76	0
3	NAG	A	403	14/15	0.75	0.34	-	56,67,70,71	0
4	SO4	A	1322	5/5	0.88	0.19	-	65,66,71,71	0
4	SO4	A	1318	5/5	0.72	0.44	-	115,116,118,119	0
4	SO4	B	1174	5/5	0.74	0.59	-	95,99,115,115	3
3	NAG	A	404	14/15	0.67	0.46	-	59,71,75,76	0
4	SO4	B	1172	5/5	0.90	0.22	-	78,80,81,82	0
4	SO4	A	1325	5/5	0.84	0.35	-	89,91,93,93	0
4	SO4	A	1317	5/5	0.84	0.38	-	92,94,95,96	0
4	SO4	A	1319	5/5	0.77	0.34	-	108,108,111,112	0
3	NAG	A	1123	14/15	0.84	0.25	-	49,57,62,62	0
4	SO4	A	1324	5/5	0.89	0.13	-	90,92,95,95	0

6.5 Other polymers [i](#)

There are no such residues in this entry.