



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 09:53 PM GMT

PDB ID : 5BS3
Title : Crystal Structure of S.A. gyrase in complex with Compound 7
Authors : Lu, J.; Patel, S.; Soisson, S.
Deposited on : 2015-06-01
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3152 (2.70-2.62)
Clashscore	102246	3524 (2.70-2.62)
Ramachandran outliers	100387	3469 (2.70-2.62)
Sidechain outliers	100360	3469 (2.70-2.62)
RSRZ outliers	91569	3161 (2.70-2.62)
RNA backbone	2183	1001 (3.08-2.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	692	
1	D	692	
2	E	20	
2	F	20	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11537 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA gyrase subunit A and B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	670	Total	C	N	O	S	0	0	0
			5262	3280	948	1009	25			
1	D	670	Total	C	N	O	S	0	2	0
			5284	3292	952	1015	25			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	409	PRO	-	expression tag	UNP P0A0K8
B	544	THR	-	linker	UNP P0A0K8
B	545	GLY	-	linker	UNP P0A0K8
B	1123	PHE	TYR	engineered mutation	UNP P20831
B	1457	THR	ALA	engineered mutation	UNP P20831
D	409	PRO	-	expression tag	UNP P0A0K8
D	544	THR	-	linker	UNP P0A0K8
D	545	GLY	-	linker	UNP P0A0K8
D	1123	PHE	TYR	engineered mutation	UNP P20831
D	1457	THR	ALA	engineered mutation	UNP P20831

- Molecule 2 is a RNA chain called DNA/RNA (5'-R(P*AP*GP*CP*CP*G)-D(P*T)-R(P*A P*GP*GP*GP*CP*CP*C)-D(P*T)-R(P*AP*CP*GP*GP*C)-D(P*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	20	Total	C	N	O	P	0	0	0
			410	193	77	120	20			
2	F	19	Total	C	N	O	P	0	0	0
			390	183	75	113	19			

- Molecule 3 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mn	0	0
			1	1		
3	D	1	Total	Mn	0	0
			1	1		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	O	S	0	0
			5	4	1		
4	E	1	Total	O	S	0	0
			5	4	1		
4	F	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is (4R)-3-fluoro-4-hydroxy-4-([(1r,4R)-4-([(3-oxo-3,4-dihydro-2H-pyrido[3,2-b][1,4]oxazin-6-yl)methyl]amino)-2-oxabicyclo[2.2.2]oct-1-yl)methyl]-4,5-dihydro-7H-pyrrolo[3,2,1-de][1,5]naphthyridin-7-one (three-letter code: WCP) (formula: C₂₆H₂₆FN₅O₅).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	F	1	Total	C	F	N	O	0	0
			37	26	1	5	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	58	Total	O	0	0
			58	58		
6	D	61	Total	O	0	0
			61	61		
6	E	8	Total	O	0	0
			8	8		
6	F	10	Total	O	0	0
			10	10		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA gyrase subunit A and B





- Molecule 1: DNA gyrase subunit A and B



- Molecule 2: DNA/RNA (5'-R(P*AP*GP*CP*CP*G)-D(P*T)-R(P*AP*GP*GP*GP*CP*CP*C)-D(P*T)-R(P*AP*CP*GP*GP*C)-D(P*T)-3')



- Molecule 2: DNA/RNA (5'-R(P*AP*GP*CP*CP*G)-D(P*T)-R(P*AP*GP*GP*GP*CP*CP*C)-D(P*T)-R(P*AP*CP*GP*GP*C)-D(P*T)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	93.34Å 93.34Å 411.07Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	26.71 – 2.65 26.44 – 2.65	Depositor EDS
% Data completeness (in resolution range)	100.0 (26.71-2.65) 100.0 (26.44-2.65)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.32 (at 2.64Å)	Xtriage
Refinement program	BUSTER 2.9.7	Depositor
R, R_{free}	0.170 , 0.213 0.179 , 0.222	Depositor DCC
R_{free} test set	2952 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	45.7	Xtriage
Anisotropy	0.081	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 19.2	EDS
Estimated twinning fraction	0.165 for h,-h-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Outliers	0 of 58374 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11537	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: WCP, MN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.50	0/5333	0.71	0/7193
1	D	0.51	0/5355	0.72	0/7223
2	E	1.14	0/459	2.10	26/706 (3.7%)
2	F	1.08	0/437	1.97	16/672 (2.4%)
All	All	0.57	0/11584	0.91	42/15794 (0.3%)

There are no bond length outliers.

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	6	DT	O4'-C1'-N1	8.16	113.71	108.00
2	F	17	G	O4'-C1'-N9	-8.13	101.70	108.20
2	F	14	DT	O4'-C1'-N1	8.11	113.68	108.00
2	F	2	G	O4'-C1'-N9	8.05	114.64	108.20
2	E	1	A	OP1-P-OP2	-7.30	108.64	119.60
2	E	13	C	C4'-C3'-C2'	-7.27	95.33	102.60
2	E	10	G	P-O3'-C3'	7.21	128.35	119.70
2	F	16	C	C1'-O4'-C4'	-6.80	104.46	109.90
2	E	19	C	O4'-C1'-N1	6.78	113.62	108.20
2	E	14	DT	O4'-C4'-C3'	-6.76	101.79	104.50
2	E	14	DT	C4-C5-C7	6.73	123.04	119.00
2	E	4	C	O4'-C1'-N1	6.69	113.56	108.20
2	E	6	DT	N3-C4-O4	-6.69	115.89	119.90
2	F	11	C	P-O3'-C3'	6.64	127.67	119.70
2	E	16	C	C1'-O4'-C4'	-6.39	104.79	109.90
2	E	13	C	C1'-O4'-C4'	-6.26	104.89	109.90
2	E	7	A	N1-C2-N3	-6.14	126.23	129.30
2	F	10	G	P-O3'-C3'	6.12	127.04	119.70
2	F	1	A	OP1-P-OP2	-6.02	110.57	119.60
2	E	14	DT	C4'-C3'-C2'	-6.01	97.69	103.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	14	DT	C4'-C3'-C2'	-5.95	97.75	103.10
2	F	9	G	P-O3'-C3'	5.92	126.80	119.70
2	E	19	C	P-O3'-C3'	5.91	126.79	119.70
2	E	6	DT	O4'-C1'-N1	5.77	112.04	108.00
2	E	3	C	N1-C2-O2	5.75	122.35	118.90
2	E	2	G	P-O3'-C3'	5.72	126.56	119.70
2	E	15	A	N1-C2-N3	-5.71	126.44	129.30
2	F	13	C	N1-C2-O2	5.67	122.30	118.90
2	F	14	DT	N3-C4-O4	-5.67	116.50	119.90
2	E	20	DT	O4'-C1'-N1	5.56	111.89	108.00
2	E	15	A	C2-N3-C4	5.51	113.35	110.60
2	E	19	C	C4'-C3'-C2'	-5.50	97.10	102.60
2	F	9	G	C4'-C3'-C2'	-5.46	97.14	102.60
2	E	14	DT	N3-C4-O4	-5.44	116.64	119.90
2	F	12	C	C1'-O4'-C4'	-5.35	105.62	109.90
2	E	11	C	P-O3'-C3'	5.13	125.85	119.70
2	F	19	C	O4'-C1'-N1	5.09	112.27	108.20
2	E	18	G	O4'-C1'-N9	-5.09	104.13	108.20
2	E	13	C	N1-C2-O2	5.06	121.94	118.90
2	F	18	G	P-O3'-C3'	5.05	125.76	119.70
2	E	14	DT	C6-C5-C7	-5.04	119.88	122.90
2	E	17	G	O4'-C1'-N9	-5.04	104.17	108.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	5262	0	5256	24	0
1	D	5284	0	5273	38	0
2	E	410	0	224	2	0
2	F	390	0	212	3	0
3	B	1	0	0	0	0
3	D	1	0	0	0	0
4	D	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	E	5	0	0	0	0
4	F	5	0	0	0	0
5	F	37	0	26	5	0
6	B	58	0	0	0	0
6	D	61	0	0	0	0
6	E	8	0	0	0	0
6	F	10	0	0	0	0
All	All	11537	0	10991	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1058:MET:HG2	1:B:1065:LYS:HG3	1.74	0.68
5:F:101:WCP:H19	5:F:101:WCP:C10	2.25	0.66
1:D:1058:MET:HG2	1:D:1065:LYS:HG3	1.77	0.66
1:B:1403:THR:HG22	1:D:1436:LEU:HD22	1.78	0.64
5:F:101:WCP:H27	5:F:101:WCP:H19	1.83	0.61
1:B:503:ILE:HD13	1:B:531:LEU:HD21	1.82	0.60
1:B:502:LYS:HG2	1:B:538:TYR:CE1	2.37	0.58
1:B:1183:ILE:HG12	1:B:1335:MET:HG2	1.86	0.58
1:B:1109:ASN:HB3	1:B:1119:ALA:HB2	1.86	0.56
1:D:460:LYS:HG2	2:F:8:G:H1'	1.89	0.55
5:F:101:WCP:O5	5:F:101:WCP:H24	2.08	0.52
1:B:1102:PRO:O	1:B:1131:ILE:HD11	2.09	0.52
1:D:1234:TYR:O	1:D:1347:ASN:HB2	2.10	0.51
1:B:1054:ASN:HB2	1:B:1136:LEU:HD13	1.93	0.50
1:B:443:THR:HG22	1:B:454:ILE:HG12	1.93	0.50
1:D:1445:TYR:CE2	1:D:1449:LEU:HD11	2.46	0.50
1:B:588:ALA:HB1	1:D:1298:LEU:HD11	1.94	0.50
1:D:1242:GLN:HG3	1:D:1328:GLN:NE2	2.28	0.49
1:D:424:LYS:O	1:D:426:PRO:HD3	2.12	0.49
1:D:1325:THR:OG1	1:D:1327:LEU:HB2	2.13	0.48
1:B:461:ILE:HD11	1:B:477:GLU:HG2	1.95	0.48
1:D:464:VAL:HG21	1:D:523:PHE:HA	1.97	0.47
1:D:1242:GLN:HG3	1:D:1328:GLN:HE21	1.79	0.47
1:D:530:PRO:HA	1:D:533:GLU:HB2	1.96	0.47
1:D:506:MET:HG2	1:D:583:LEU:HD11	1.97	0.46
1:D:1224:ILE:HG21	1:D:1230:ILE:HD11	1.97	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:1225:LEU:HD21	1:D:1244:ARG:HD2	1.98	0.45
1:B:1445:TYR:CE1	1:B:1449:LEU:HD11	2.50	0.45
1:B:1075:MET:HE1	1:B:1083:ASP:HB3	1.99	0.45
1:D:1242:GLN:HE22	1:D:1330:SER:HB3	1.82	0.45
1:D:1135:LEU:HG	1:D:1164:PHE:CE2	2.52	0.45
1:D:433:LEU:HD23	1:D:455:LEU:HB3	1.99	0.45
1:D:443:THR:HG22	1:D:454:ILE:HG12	1.98	0.44
5:F:101:WCP:H27	5:F:101:WCP:C16	2.47	0.44
1:B:1066:LYS:HB2	1:B:1069:ARG:HG3	2.00	0.44
1:D:626:VAL:HG11	2:E:17:G:H3'	1.99	0.44
1:B:585:GLU:HG3	1:D:1123:PHE:O	2.18	0.44
1:D:437:ASP:HB2	2:F:10:G:H5''	1.99	0.44
1:B:1101:TYR:CZ	1:B:1188:LEU:HB2	2.53	0.43
1:D:1050:LEU:HD11	1:D:1103:LEU:HD13	2.00	0.43
1:D:1064:TYR:HB3	1:D:1125:GLU:HB3	2.01	0.43
1:D:460:LYS:HA	1:D:516:ILE:HG12	2.01	0.42
1:D:1269:ASN:HB3	1:D:1272:ARG:HB2	2.00	0.42
1:D:427:GLU:CD	1:D:427:GLU:H	2.22	0.42
1:B:425:SER:HB3	1:B:427:GLU:OE1	2.20	0.42
1:B:1385:ARG:HG3	1:B:1438:ARG:CZ	2.49	0.42
1:B:1430:LEU:HD12	1:D:1425:ILE:O	2.19	0.42
1:B:626:VAL:HG11	2:F:17:G:H3'	2.01	0.42
1:D:591:LEU:HD12	1:D:595:THR:CG2	2.50	0.42
1:B:1430:LEU:O	1:B:1433:LEU:HB2	2.19	0.41
1:D:1393:GLU:HB3	1:D:1415:PHE:HZ	1.85	0.41
2:E:11:C:N4	5:F:101:WCP:O5	2.53	0.41
1:D:426:PRO:HA	1:D:429:CYS:HB2	2.02	0.41
1:D:1054:ASN:HB2	1:D:1136:LEU:HD13	2.03	0.41
1:B:588:ALA:CB	1:D:1298:LEU:HD11	2.50	0.41
1:D:1231:ARG:NH2	1:D:1235:GLU:OE1	2.54	0.41
1:D:1234:TYR:CD2	1:D:1348:LEU:HD22	2.56	0.41
1:D:1029:VAL:HA	1:D:1033:ARG:HB3	2.03	0.41
1:D:444:LYS:HG2	1:D:454:ILE:HD13	2.03	0.40
1:B:1036:PRO:HG3	1:B:1172:ALA:CB	2.52	0.40
1:B:1426:LEU:HB3	1:D:1431:ARG:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	664/692 (96%)	642 (97%)	22 (3%)	0	100	100
1	D	666/692 (96%)	643 (96%)	22 (3%)	1 (0%)	52	77
All	All	1330/1384 (96%)	1285 (97%)	44 (3%)	1 (0%)	56	81

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	490	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	561/591 (95%)	541 (96%)	20 (4%)	42	70
1	D	564/591 (95%)	549 (97%)	15 (3%)	52	80
All	All	1125/1182 (95%)	1090 (97%)	35 (3%)	47	75

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	450	ARG
1	B	461	ILE
1	B	480	GLN
1	B	489	ILE
1	B	497	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	595	THR
1	B	609	GLU
1	B	1085	SER
1	B	1131	ILE
1	B	1138	ASP
1	B	1228	SER
1	B	1244	ARG
1	B	1292	LEU
1	B	1299	ARG
1	B	1300	THR
1	B	1327	LEU
1	B	1344	LYS
1	B	1396	SER
1	B	1410	SER
1	B	1448	LEU
1	D	420	ASP
1	D	427	GLU
1	D	624	ASP
1	D	1024	ASP
1	D	1084	SER
1	D	1148	ASP
1	D	1242	GLN
1	D	1285	LYS
1	D	1299	ARG
1	D	1400	GLU
1	D	1408	MET
1	D	1409	GLU
1	D	1413	GLN
1	D	1450	ASN
1	D	1465	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	501	HIS
1	B	541	GLN
1	B	605	GLN
1	B	631	GLN
1	B	1010	ASN
1	B	1153	ASN
1	D	480	GLN
1	D	590	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	605	GLN
1	D	1242	GLN
1	D	1368	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	E	0/20	-	-
2	F	0/20	-	-
All	All	0/40	-	-

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	D	1502	-	4,4,4	0.17	0	6,6,6	0.05	0
4	SO4	E	101	-	4,4,4	0.45	0	6,6,6	0.23	0
5	WCP	F	101	-	37,43,43	1.97	8 (21%)	30,68,68	1.70	6 (20%)
4	SO4	F	102	-	4,4,4	0.28	0	6,6,6	0.16	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SO4	D	1502	-	-	0/0/0/0	0/0/0/0
4	SO4	E	101	-	-	0/0/0/0	0/0/0/0
5	WCP	F	101	-	-	1/12/53/53	0/4/7/7
4	SO4	F	102	-	-	0/0/0/0	0/0/0/0

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	F	101	WCP	C25-C14	2.04	1.56	1.53
5	F	101	WCP	C8-N1	2.59	1.42	1.38
5	F	101	WCP	C6-C5	3.15	1.47	1.41
5	F	101	WCP	C15-C14	3.25	1.57	1.53
5	F	101	WCP	C11-C9	4.56	1.59	1.54
5	F	101	WCP	C10-C9	4.68	1.63	1.51
5	F	101	WCP	C21-C20	5.15	1.57	1.51
5	F	101	WCP	C4-C3	5.52	1.44	1.37

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	F	101	WCP	C6-C5-C1	-3.68	116.14	118.31
5	F	101	WCP	C18-C17-N3	-3.17	106.10	113.26
5	F	101	WCP	O5-C9-C2	-2.92	103.78	112.05
5	F	101	WCP	N5-C19-N4	2.40	119.50	116.30
5	F	101	WCP	C11-C9-C2	3.54	122.10	111.38
5	F	101	WCP	C6-C5-N2	3.83	124.93	118.54

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	F	101	WCP	C2-C9-C11-C12

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	F	101	WCP	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	1
1	D	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	D	545:GLY	C	580:TYR	N	3.23
1	B	545:GLY	C	580:TYR	N	3.08

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	670/692 (96%)	-0.16	1 (0%) 95 96	28, 46, 62, 76	0
1	D	670/692 (96%)	-0.15	1 (0%) 95 96	30, 45, 65, 81	0
2	E	20/20 (100%)	-0.73	0 100 100	30, 39, 51, 57	0
2	F	19/20 (95%)	-0.57	0 100 100	33, 39, 60, 85	0
All	All	1379/1424 (96%)	-0.17	2 (0%) 95 96	28, 45, 64, 85	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1433	LEU	2.1
1	D	490	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	WCP	F	101	37/37	0.90	0.20	1.64	42,50,65,67	0
4	SO4	E	101	5/5	0.91	0.15	1.44	101,106,106,107	0
3	MN	B	1501	1/1	1.00	0.17	0.97	45,45,45,45	0
4	SO4	F	102	5/5	0.92	0.14	0.41	93,97,98,98	0
3	MN	D	1501	1/1	0.98	0.16	0.03	44,44,44,44	0
4	SO4	D	1502	5/5	0.96	0.15	-	128,132,133,134	0

6.5 Other polymers [i](#)

There are no such residues in this entry.