



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 12, 2016 – 09:07 AM EDT

PDB ID : 5BUK
Title : Structure of flavin-dependent chlorinase Mpy16
Authors : Agarwal, V.; Louie, G.V.; Noel, J.P.; Moore, B.S.
Deposited on : 2015-06-03
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027674
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027674


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1833 (1.96-1.96)
Clashscore	102246	1953 (1.96-1.96)
Ramachandran outliers	100387	1936 (1.96-1.96)
Sidechain outliers	100360	1936 (1.96-1.96)
RSRZ outliers	91569	1835 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	450	
1	B	450	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	502	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7359 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called FADH2-dependent halogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	434	Total	C	N	O	S	0	4	0
			3449	2211	588	642	8			
1	B	426	Total	C	N	O	S	0	0	0
			3370	2158	576	628	8			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP J7H1A1
A	-1	GLY	-	expression tag	UNP J7H1A1
A	0	SER	-	expression tag	UNP J7H1A1
B	-2	SER	-	expression tag	UNP J7H1A1
B	-1	GLY	-	expression tag	UNP J7H1A1
B	0	SER	-	expression tag	UNP J7H1A1

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	215	Total	O	0	0
			215	215		
4	B	213	Total	O	0	0
			213	213		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: FADH2-dependent halogenase



- Molecule 1: FADH2-dependent halogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.83Å 73.28Å 189.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.08 – 1.95 38.08 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.8 (38.08-1.95) 99.8 (38.08-1.95)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.77 (at 1.95Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.1_743)	Depositor
R, R_{free}	0.170 , 0.212 0.166 , 0.208	Depositor DCC
R_{free} test set	3273 reflections (5.06%)	DCC
Wilson B-factor (Å ²)	29.7	Xtriage
Anisotropy	0.041	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7359	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/3545	0.52	0/4805
1	B	0.37	0/3454	0.51	0/4680
All	All	0.37	0/6999	0.51	0/9485

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3449	0	3386	29	0
1	B	3370	0	3290	28	0
2	A	53	0	31	3	0
2	B	53	0	31	3	0
3	A	6	0	8	1	0
4	A	215	0	0	5	0
4	B	213	0	0	10	1
All	All	7359	0	6746	58	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:346:ARG:NH1	4:A:601:HOH:O	2.00	0.94
1:A:198:ASN:HB3	4:A:795:HOH:O	1.86	0.74
1:A:4:GLN:HG2	1:A:29:LYS:HE2	1.78	0.66
1:B:346:ARG:NH1	4:B:601:HOH:O	2.24	0.66
1:B:417:ARG:HA	4:B:608:HOH:O	1.96	0.65
1:B:44:GLY:HA2	2:B:500:FAD:C4X	2.28	0.64
1:A:81:ASP:HB2	1:A:230:PRO:HG2	1.79	0.63
1:B:44:GLY:HA2	2:B:500:FAD:N5	2.15	0.62
1:B:375:ILE:HG12	4:B:608:HOH:O	1.98	0.61
1:A:44:GLY:HA2	2:A:501:FAD:C4X	2.31	0.61
1:A:179:GLY:HA2	1:A:314:ASP:HB2	1.82	0.60
1:B:438:LEU:C	4:B:632:HOH:O	2.39	0.60
1:A:274:SER:C	1:A:275:ARG:HG3	2.26	0.56
1:B:279:HIS:O	1:B:283:LYS:HG2	2.07	0.55
1:B:98:ARG:HD3	4:B:609:HOH:O	2.06	0.55
1:A:87:LYS:HZ1	1:A:96:ASP:HA	1.72	0.53
1:A:44:GLY:HA2	2:A:501:FAD:N5	2.24	0.52
1:B:3:PRO:HB3	1:B:170:PRO:HB2	1.92	0.51
1:B:179:GLY:HA2	1:B:314:ASP:HB2	1.92	0.51
1:A:148:ARG:NH2	4:A:607:HOH:O	2.44	0.50
1:A:65:GLU:OE2	1:A:116:TYR:OH	2.19	0.50
1:A:186:SER:O	1:A:189:LYS:HD3	2.11	0.50
1:B:366:VAL:HB	4:B:744:HOH:O	2.12	0.50
1:B:417:ARG:NE	4:B:608:HOH:O	2.44	0.50
1:A:230:PRO:HD2	4:A:788:HOH:O	2.10	0.49
1:A:87:LYS:NZ	1:A:96:ASP:HA	2.27	0.49
1:B:288:VAL:HG12	1:B:289:ARG:HG3	1.95	0.49
1:B:398:LYS:HE2	1:B:407:ASP:OD2	2.12	0.49
1:A:386:PHE:HB3	1:A:396:VAL:HG21	1.96	0.47
1:A:341[A]:ILE:HD13	1:A:355:PHE:CE1	2.49	0.47
1:B:149:VAL:HG22	1:B:158:LEU:HD23	1.96	0.47
1:B:254:LYS:HE3	1:B:258:LYS:HE3	1.97	0.47
1:A:426:ASP:OD1	1:A:428:ASN:HB2	2.15	0.46
1:B:314:ASP:OD2	2:B:500:FAD:H3'	2.16	0.46
1:B:432:HIS:N	1:B:433:PRO:CD	2.79	0.46
1:B:257:LEU:HD12	4:B:812:HOH:O	2.15	0.45
1:B:369:TRP:CZ2	1:B:402:GLY:HA2	2.53	0.44
1:A:263:ASP:HB2	4:A:791:HOH:O	2.18	0.44
1:A:337:ALA:O	1:A:341[A]:ILE:HG12	2.18	0.44
1:B:145:ARG:HD3	4:B:745:HOH:O	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:269:TRP:CZ3	1:B:283:LYS:HD3	2.54	0.43
1:B:417:ARG:CZ	4:B:608:HOH:O	2.67	0.43
1:A:173:MET:HE2	1:A:341[A]:ILE:HD11	2.01	0.43
1:A:214:LEU:HA	1:A:214:LEU:HD23	1.76	0.42
1:B:132:HIS:O	1:B:136:GLN:HG2	2.18	0.42
1:A:46:SER:O	1:A:324:SER:HB3	2.19	0.42
1:A:156:PRO:HD3	3:A:502:GOL:H11	2.01	0.42
1:B:283:LYS:HA	1:B:283:LYS:HD2	1.81	0.41
1:A:341[A]:ILE:HD13	1:A:355:PHE:HE1	1.85	0.41
1:B:128:LEU:HD23	1:B:128:LEU:C	2.41	0.41
1:A:207:ASP:HB2	1:A:288[A]:VAL:CG2	2.51	0.41
1:B:46:SER:O	1:B:324:SER:HB3	2.21	0.41
1:A:159:GLU:HB3	1:A:166:ARG:HH11	1.85	0.41
2:A:501:FAD:H1'1	2:A:501:FAD:H9	1.74	0.41
1:A:407:ASP:OD2	1:A:410:PRO:HA	2.22	0.40
1:A:438:LEU:HD23	1:A:438:LEU:C	2.42	0.40
1:B:421:LYS:HE3	1:B:421:LYS:HB3	1.90	0.40
1:A:273:GLY:C	1:A:275:ARG:N	2.75	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:669:HOH:O	4:B:800:HOH:O[4_455]	2.07	0.13

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	434/450 (96%)	426 (98%)	8 (2%)	0	100	100
1	B	422/450 (94%)	412 (98%)	10 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	856/900 (95%)	838 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	367/376 (98%)	362 (99%)	5 (1%)	74	70
1	B	357/376 (95%)	351 (98%)	6 (2%)	68	63
All	All	724/752 (96%)	713 (98%)	11 (2%)	72	68

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	ARG
1	A	110	SER
1	A	311	MET
1	A	413	LEU
1	A	421	LYS
1	B	40	ARG
1	B	101	GLU
1	B	258	LYS
1	B	277	GLU
1	B	311	MET
1	B	386	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	A	501	-	52,58,58	1.72	15 (28%)	52,89,89	2.51	11 (21%)
3	GOL	A	502	-	5,5,5	0.28	0	5,5,5	0.36	0
2	FAD	B	500	-	52,58,58	1.80	15 (28%)	52,89,89	2.51	14 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	501	-	-	0/30/50/50	0/6/6/6
3	GOL	A	502	-	-	0/4/4/4	0/0/0/0
2	FAD	B	500	-	-	0/30/50/50	0/6/6/6

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	500	FAD	C1'-N10	-4.92	1.43	1.48
2	A	501	FAD	C1'-N10	-4.09	1.44	1.48
2	B	500	FAD	PA-O1A	-3.72	1.37	1.51
2	A	501	FAD	C2-N3	-3.23	1.31	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	FAD	P-O2P	-2.99	1.42	1.55
2	A	501	FAD	PA-O2A	-2.99	1.42	1.55
2	A	501	FAD	P-O1P	-2.95	1.40	1.51
2	B	500	FAD	PA-O2A	-2.91	1.42	1.55
2	A	501	FAD	C6-C5X	-2.85	1.37	1.41
2	B	500	FAD	C6-C5X	-2.84	1.37	1.41
2	B	500	FAD	P-O1P	-2.75	1.41	1.51
2	A	501	FAD	PA-O1A	-2.66	1.41	1.51
2	B	500	FAD	P-O2P	-2.55	1.44	1.55
2	B	500	FAD	C2-N3	-2.48	1.33	1.38
2	B	500	FAD	O4B-C4B	-2.45	1.39	1.45
2	A	501	FAD	C2-N1	-2.36	1.33	1.38
2	B	500	FAD	C4A-N3A	-2.36	1.32	1.35
2	B	500	FAD	C10-N10	-2.30	1.36	1.39
2	A	501	FAD	C4A-N3A	-2.29	1.32	1.35
2	A	501	FAD	O4B-C4B	-2.27	1.39	1.45
2	B	500	FAD	C5A-N7A	-2.07	1.32	1.39
2	A	501	FAD	C5A-C4A	2.02	1.45	1.40
2	A	501	FAD	C4-C4X	2.20	1.45	1.41
2	B	500	FAD	C8-C7	2.26	1.47	1.41
2	A	501	FAD	C8-C7	2.36	1.47	1.41
2	A	501	FAD	C9A-C5X	2.40	1.47	1.42
2	B	500	FAD	C9A-C5X	2.60	1.48	1.42
2	B	500	FAD	C4-C4X	2.83	1.47	1.41
2	A	501	FAD	C4X-C10	2.85	1.46	1.40
2	B	500	FAD	C4X-C10	3.62	1.47	1.40

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	500	FAD	C4-C4X-C10	-8.69	114.38	119.94
2	A	501	FAD	C4-C4X-C10	-8.09	114.77	119.94
2	A	501	FAD	N3A-C2A-N1A	-5.27	124.73	128.87
2	B	500	FAD	N3-C2-N1	-3.83	121.24	127.69
2	A	501	FAD	C4X-C4-N3	-3.71	118.67	123.52
2	B	500	FAD	N3A-C2A-N1A	-3.67	125.99	128.87
2	A	501	FAD	N3-C2-N1	-3.64	121.57	127.69
2	B	500	FAD	C4X-C4-N3	-3.60	118.81	123.52
2	B	500	FAD	C4X-C10-N10	-3.07	118.29	120.52
2	B	500	FAD	C1B-N9A-C4A	-2.55	123.96	126.81
2	A	501	FAD	C1B-N9A-C4A	-2.55	123.96	126.81
2	A	501	FAD	C4X-C10-N10	-2.54	118.67	120.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	500	FAD	O2'-C2'-C3'	2.01	114.14	108.96
2	B	500	FAD	C1'-N10-C9A	2.09	121.25	118.83
2	A	501	FAD	C5X-C9A-N10	2.27	119.28	117.58
2	B	500	FAD	C5X-C9A-N10	2.43	119.40	117.58
2	B	500	FAD	C4B-O4B-C1B	2.55	112.35	109.64
2	B	500	FAD	N6A-C6A-N1A	2.63	122.94	118.52
2	A	501	FAD	C6-C5X-N5	3.28	123.01	118.92
2	B	500	FAD	C4X-N5-C5X	3.49	120.84	116.72
2	B	500	FAD	C4-C4X-N5	4.16	123.76	118.70
2	A	501	FAD	C4X-N5-C5X	4.30	121.79	116.72
2	A	501	FAD	C4-C4X-N5	4.63	124.33	118.70
2	A	501	FAD	C4-N3-C2	10.39	123.83	115.16
2	B	500	FAD	C4-N3-C2	10.62	124.02	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	FAD	3	0
3	A	502	GOL	1	0
2	B	500	FAD	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	434/450 (96%)	-0.12	2 (0%) 91 95	19, 29, 45, 61	0
1	B	426/450 (94%)	0.06	7 (1%) 74 83	20, 30, 49, 68	0
All	All	860/900 (95%)	-0.03	9 (1%) 84 89	19, 29, 47, 68	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	261	LYS	4.6
1	B	154	ALA	3.3
1	B	258	LYS	2.8
1	B	262	ASP	2.7
1	B	155	PHE	2.7
1	A	231	VAL	2.2
1	B	327	VAL	2.2
1	A	392	TYR	2.2
1	B	113	HIS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	GOL	A	502	6/6	0.91	0.15	4.70	28,38,40,55	0
2	FAD	B	500	53/53	0.98	0.13	0.20	19,22,25,27	0
2	FAD	A	501	53/53	0.98	0.10	-0.30	18,23,25,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.