



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:43 AM GMT

PDB ID : 3BXM
Title : Structure of an inactive mutant of human glutamate carboxypeptidase II [GCPII(E424A)] in complex with N-acetyl-Asp-Glu (NAAG)
Authors : Lubkowski, J.; Barinka, C.
Deposited on : 2008-01-14
Resolution : 1.71 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3998 (1.74-1.70)
Clashscore	102246	4425 (1.74-1.70)
Ramachandran outliers	100387	4360 (1.74-1.70)
Sidechain outliers	100360	4360 (1.74-1.70)
RSRZ outliers	91569	4010 (1.74-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	709	 6% 88% 9% ..
2	I	3	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	A	1758	-	-	-	X
3	NAG	A	1761	-	-	-	X
5	NAG	A	1763	-	-	-	X

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 6350 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate carboxypeptidase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	694	Total	C	N	O	S	0	25	0
			5653	3639	946	1046	22			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	42	ARG	-	EXPRESSION TAG	UNP Q04609
A	43	SER	-	EXPRESSION TAG	UNP Q04609
A	424	ALA	GLU	ENGINEERED	UNP Q04609

- Molecule 2 is a protein called N-Acetyl-Aspartyl-Glutamate (NAAG).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	I	3	Total	C	N	O	0	0	0
			21	11	2	8			

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	2	Total	C	N	O	0	0
			28	16	2	10		
3	A	2	Total	C	N	O	0	0
			28	16	2	10		
3	A	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 4 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is a polymer of unknown type called SUGAR (4-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	4	Total	C	N	O	0	0
			50	28	2	20		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	Zn	0	0
			2	2		

- Molecule 7 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Ca	0	0
			1	1		

- Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	Cl	0	0
			1	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	492	Total	O	0	0
			492	492		
9	I	4	Total	O	0	0
			4	4		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glutamate carboxypeptidase 2



• Molecule 2: N-Acetyl-Aspartyl-Glutamate (NAAG)



There are no outlier residues recorded for this chain.

4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	102.06 Å 129.80 Å 159.59 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 1.71 19.98 – 1.71	Depositor EDS
% Data completeness (in resolution range)	98.6 (15.00-1.71) 98.5 (19.98-1.71)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.73 (at 1.70 Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.184 , 0.213 0.184 , 0.208	Depositor DCC
R_{free} test set	1370 reflections (1.23%)	DCC
Wilson B-factor (Å ²)	25.0	Xtriage
Anisotropy	0.509	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 58.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 113292 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6350	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, BMA, NAG, ACE, CL, CA, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.05	5/5883 (0.1%)	0.93	6/7971 (0.1%)
2	I	1.54	0/18	0.87	0/22
All	All	1.05	5/5901 (0.1%)	0.93	6/7993 (0.1%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	733	TYR	CD1-CE1	6.55	1.49	1.39
1	A	434	TRP	CE3-CZ3	6.49	1.49	1.38
1	A	426	PHE	CE2-CZ	5.76	1.48	1.37
1	A	102	GLU	CG-CD	5.64	1.60	1.51
1	A	83	LEU	CG-CD2	5.12	1.70	1.51

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	536	ARG	NE-CZ-NH1	-7.18	116.71	120.30
1	A	369	ASP	CB-CG-OD1	6.59	124.23	118.30
1	A	370	ARG	NE-CZ-NH2	-6.54	117.03	120.30
1	A	370	ARG	NE-CZ-NH1	6.04	123.32	120.30
1	A	536	ARG	NE-CZ-NH2	5.94	123.27	120.30
1	A	673	ARG	NE-CZ-NH1	5.56	123.08	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5653	0	5507	32	1
2	I	21	0	13	0	0
3	A	84	0	75	2	0
4	A	42	0	39	1	0
5	A	50	0	43	0	1
6	A	2	0	0	0	0
7	A	1	0	0	0	0
8	A	1	0	0	0	0
9	A	492	0	0	10	0
9	I	4	0	0	0	0
All	All	6350	0	5677	34	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:660[A]:VAL:O	1:A:664[A]:MET:HG2	1.30	1.28
1:A:151:TYR:O	1:A:153:ASN:N	1.96	0.96
3:A:1756:NAG:H81	9:A:2014:HOH:O	1.89	0.73
1:A:620:GLN:NE2	9:A:2200:HOH:O	2.02	0.72
1:A:610:LYS:HE2	9:A:1948:HOH:O	1.91	0.70
1:A:610:LYS:CE	9:A:1948:HOH:O	2.46	0.62
1:A:617:LYS:NZ	9:A:2132:HOH:O	2.38	0.55
1:A:325:VAL:HB	1:A:326:PRO:HD2	1.89	0.54
1:A:347:HIS:HE1	9:A:2078:HOH:O	1.90	0.53
1:A:653:PHE:HZ	1:A:661[A]:LEU:HD13	1.74	0.52
1:A:92:LEU:O	1:A:96[B]:ILE:HG12	2.10	0.51
1:A:246:TRP:CD1	4:A:1760:NAG:H83	2.45	0.51
1:A:688[B]:ARG:NH2	9:A:1836:HOH:O	2.26	0.51
1:A:367:GLU:OE1	1:A:662[A]:ARG:NH1	2.43	0.50
1:A:517:SER:HB2	1:A:694:PRO:HG3	1.94	0.49
1:A:656[B]:SER:O	1:A:658[B]:PRO:HD3	2.12	0.49
1:A:612:TYR:CZ	1:A:616:MET:HG3	2.48	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:188:LEU:CD2	1:A:329:VAL:HG11	2.43	0.49
1:A:369:ASP:OD2	1:A:662[B]:ARG:NH2	2.35	0.48
1:A:258:ILE:HD13	1:A:294:VAL:HB	1.94	0.48
1:A:648:GLU:O	1:A:651[B]:GLN:HG2	2.13	0.47
1:A:551:LEU:HD22	1:A:556:TYR:HB2	1.95	0.47
3:A:1767:NAG:H83	9:A:2098:HOH:O	2.15	0.46
1:A:242:TYR:OH	9:A:2159:HOH:O	1.95	0.44
1:A:91:GLN:HE22	1:A:94:LYS:NZ	2.14	0.44
1:A:347:HIS:CE1	9:A:2078:HOH:O	2.69	0.44
1:A:489:GLU:H	1:A:489:GLU:CD	2.21	0.44
1:A:56:HIS:CD2	1:A:405[A]:LEU:HD22	2.54	0.43
1:A:71:LYS:HG3	1:A:570:PHE:CE2	2.55	0.42
1:A:505:GLU:HB2	1:A:506:PHE:CE1	2.55	0.41
1:A:516:GLY:O	1:A:526[B]:GLN:NE2	2.53	0.41
1:A:659[B]:ILE:O	1:A:663[B]:MET:HG3	2.21	0.40
1:A:468:PRO:CG	1:A:603:VAL:HG21	2.51	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:276:GLU:OE1	5:A:1765:BMA:O2[2_565]	2.10	0.10

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	714/709 (101%)	691 (97%)	21 (3%)	2 (0%)	46	26
2	I	1/3 (33%)	1 (100%)	0	0	100	100
All	All	715/712 (100%)	692 (97%)	21 (3%)	2 (0%)	46	26

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	152	GLU
1	A	382	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	615/604 (102%)	601 (98%)	14 (2%)	58	37
2	I	2/2 (100%)	2 (100%)	0	100	100
All	All	617/606 (102%)	603 (98%)	14 (2%)	63	37

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	171	GLU
1	A	174	LEU
1	A	189	GLU
1	A	239	VAL
1	A	279[A]	TYR
1	A	279[B]	TYR
1	A	303	GLN
1	A	486	GLU
1	A	519	ASN
1	A	537	TYR
1	A	600	TYR
1	A	662[A]	ARG
1	A	662[B]	ARG
1	A	673	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	91	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	136	ASN
1	A	618	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

10 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	1755	1,3	14,14,15	0.59	0	15,19,21	1.56	3 (20%)
3	NAG	A	1756	3	14,14,15	0.63	0	15,19,21	1.69	2 (13%)
3	NAG	A	1758	1,3	14,14,15	0.77	0	15,19,21	0.89	1 (6%)
3	NAG	A	1761	1,3	14,14,15	0.96	1 (7%)	15,19,21	0.74	0
3	NAG	A	1762	3	14,14,15	0.39	0	15,19,21	1.36	2 (13%)
5	NAG	A	1763	1,5	14,14,15	1.08	1 (7%)	15,19,21	1.51	3 (20%)
5	NAG	A	1764	5	14,14,15	0.59	0	15,19,21	1.39	2 (13%)
5	BMA	A	1765	5	11,11,12	0.99	1 (9%)	14,15,17	1.24	2 (14%)
5	MAN	A	1766	5	11,11,12	0.55	0	14,15,17	1.06	0
3	NAG	A	1767	3	14,14,15	0.35	0	15,19,21	1.74	5 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1755	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1756	3	-	0/6/23/26	0/1/1/1
3	NAG	A	1758	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1761	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1762	3	-	0/6/23/26	0/1/1/1
5	NAG	A	1763	1,5	-	0/6/23/26	0/1/1/1
5	NAG	A	1764	5	-	0/6/23/26	0/1/1/1
5	BMA	A	1765	5	-	0/2/19/22	0/1/1/1
5	MAN	A	1766	5	-	0/2/19/22	0/1/1/1
3	NAG	A	1767	3	-	0/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1765	BMA	O5-C1	-2.31	1.39	1.43
3	A	1761	NAG	O7-C7	2.42	1.28	1.23
5	A	1763	NAG	C1-C2	3.24	1.56	1.52

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1755	NAG	C2-N2-C7	-3.21	118.92	123.04
5	A	1764	NAG	C3-C4-C5	-2.99	104.99	110.20
3	A	1767	NAG	O7-C7-C8	-2.80	116.92	122.06
5	A	1765	BMA	O3-C3-C2	-2.73	105.07	110.00
5	A	1763	NAG	O4-C4-C5	-2.69	102.12	109.24
5	A	1763	NAG	C6-C5-C4	-2.63	106.52	113.02
3	A	1755	NAG	O3-C3-C4	-2.54	104.63	110.34
5	A	1765	BMA	O5-C1-C2	-2.40	106.97	110.86
3	A	1767	NAG	C4-C3-C2	-2.27	107.70	111.23
3	A	1758	NAG	C3-C2-N2	-2.12	105.48	110.56
5	A	1764	NAG	O4-C4-C5	-2.03	103.85	109.24
3	A	1767	NAG	C3-C4-C5	-2.01	106.70	110.20
3	A	1755	NAG	C4-C3-C2	2.36	114.89	111.23
3	A	1767	NAG	C2-N2-C7	2.42	126.15	123.04
3	A	1762	NAG	C8-C7-N2	2.69	121.25	116.11
5	A	1763	NAG	C1-O5-C5	2.75	115.74	112.25
3	A	1762	NAG	C1-O5-C5	3.09	116.17	112.25
3	A	1767	NAG	C8-C7-N2	3.20	122.23	116.11
3	A	1756	NAG	C2-N2-C7	3.86	128.00	123.04
3	A	1756	NAG	C1-O5-C5	4.66	118.16	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1756	NAG	1	0
5	A	1765	BMA	0	1
3	A	1767	NAG	1	0

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	1757	1	14,14,15	0.66	0	15,19,21	2.05	4 (26%)
4	NAG	A	1759	1	14,14,15	0.42	0	15,19,21	1.99	1 (6%)
4	NAG	A	1760	1	14,14,15	0.61	0	15,19,21	1.62	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1757	1	-	0/6/23/26	0/1/1/1
4	NAG	A	1759	1	-	0/6/23/26	0/1/1/1
4	NAG	A	1760	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	A	1760	NAG	C2-N2-C7	-2.74	119.52	123.04
4	A	1757	NAG	C2-N2-C7	-2.60	119.70	123.04
4	A	1757	NAG	O5-C5-C6	2.26	112.23	107.35
4	A	1757	NAG	C3-C4-C5	2.77	115.03	110.20
4	A	1760	NAG	C1-O5-C5	3.64	116.87	112.25
4	A	1757	NAG	C1-O5-C5	5.67	119.45	112.25
4	A	1759	NAG	C1-O5-C5	6.93	121.04	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1760	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	694/709 (97%)	0.20	41 (5%) 26 27	17, 27, 46, 63	0
2	I	2/3 (66%)	-0.65	0 100 100	26, 26, 26, 27	0
All	All	696/712 (97%)	0.20	41 (5%) 26 27	17, 27, 46, 63	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	55	LYS	6.2
1	A	541	TRP	6.1
1	A	153	ASN	6.0
1	A	506	PHE	5.3
1	A	155	SER	4.9
1	A	719	VAL	4.9
1	A	542	GLU	4.8
1	A	335	GLY	4.1
1	A	134	ASP	3.7
1	A	336	ASN	3.6
1	A	507	SER	3.6
1	A	133	GLU	3.4
1	A	487	GLY	3.3
1	A	543	THR	3.2
1	A	237	PRO	3.2
1	A	656[A]	SER	3.0
1	A	652	ASP	3.0
1	A	136	ASN	3.0
1	A	402	PHE	2.9
1	A	284	ALA	2.9
1	A	154	VAL	2.9
1	A	324	LYS	2.9
1	A	505	GLU	2.9
1	A	123	THR	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	152	GLU	2.8
1	A	193	LYS	2.8
1	A	135	GLY	2.7
1	A	219	LEU	2.5
1	A	409	GLY	2.5
1	A	283	ILE	2.4
1	A	746	LEU	2.4
1	A	339	THR	2.3
1	A	174	LEU	2.3
1	A	551	LEU	2.3
1	A	124	HIS	2.3
1	A	122	LYS	2.2
1	A	732	ILE	2.2
1	A	238	GLY	2.2
1	A	504	PRO	2.1
1	A	408	GLU	2.1
1	A	342	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	NAG	A	1763	14/15	0.87	0.13	6.90	24,30,38,48	0
3	NAG	A	1761	14/15	0.91	0.17	6.35	30,32,39,43	0
3	NAG	A	1758	14/15	0.85	0.16	3.07	40,44,48,52	0
5	MAN	A	1766	11/12	0.87	0.18	1.33	47,49,53,56	0
3	NAG	A	1756	14/15	0.68	0.32	-	42,51,55,57	0
3	NAG	A	1755	14/15	0.92	0.11	-	34,40,45,46	0
3	NAG	A	1762	14/15	0.90	0.23	-	34,41,50,50	0
5	BMA	A	1765	11/12	0.92	0.18	-	40,42,46,46	0
3	NAG	A	1767	14/15	0.73	0.31	-	55,59,64,64	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	NAG	A	1764	14/15	0.90	0.28	-	38,41,49,53	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
7	CA	A	1753	1/1	1.00	0.07	-0.23	18,18,18,18	0
8	CL	A	1754	1/1	1.00	0.05	-1.43	26,26,26,26	0
6	ZN	A	1752	1/1	1.00	0.04	-2.76	21,21,21,21	0
6	ZN	A	1751	1/1	1.00	0.03	-3.78	21,21,21,21	0
4	NAG	A	1760	14/15	0.80	0.17	-	30,43,46,49	0
4	NAG	A	1757	14/15	0.66	0.33	-	50,58,63,66	0
4	NAG	A	1759	14/15	0.83	0.35	-	63,68,71,72	0

6.5 Other polymers [i](#)

There are no such residues in this entry.