



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 12:54 AM GMT

PDB ID : 2C2V
Title : CRYSTAL STRUCTURE OF THE CHIP-UBC13-UEV1A COMPLEX
Authors : Zhang, M.; Roe, S.M.; Pearl, L.H.
Deposited on : 2005-09-30
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1451 (2.90-2.90)
Clashscore	102246	1668 (2.90-2.90)
Ramachandran outliers	100387	1630 (2.90-2.90)
Sidechain outliers	100360	1632 (2.90-2.90)
RSRZ outliers	91569	1456 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	154	
1	E	154	
1	H	154	
1	K	154	
2	C	142	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	142	<div><div><div></div><div></div><div></div><div></div><div></div></div><div>%15%31%33%18%•</div></div>
2	I	142	<div><div><div></div><div></div><div></div><div></div><div></div></div><div>%15%27%35%20%•</div></div>
2	L	142	<div><div><div></div><div></div><div></div><div></div><div></div></div><div>13%32%35%18%•</div></div>
3	S	78	<div><div><div></div><div></div><div></div><div></div><div></div></div><div>5%26%42%23%•</div></div>
3	T	78	<div><div><div></div><div></div><div></div><div></div><div></div></div><div>8%8%27%47%18%</div></div>
3	U	78	<div><div><div></div><div></div><div></div><div></div><div></div></div><div>%5%22%53%14%6%</div></div>
3	V	78	<div><div><div></div><div></div><div></div><div></div><div></div></div><div>%•29%29%29%9%</div></div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11636 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UBIQUITIN-CONJUGATING ENZYME E2 N.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	152	Total	C	N	O	S	0	0	0
			1202	769	208	221	4			
1	E	149	Total	C	N	O	S	0	0	0
			1187	761	205	217	4			
1	H	149	Total	C	N	O	S	0	0	0
			1187	761	205	217	4			
1	K	152	Total	C	N	O	S	0	0	0
			1202	769	208	221	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	5	SER	MET	CONFLICT	UNP P61089
K	5	SER	MET	CONFLICT	UNP P61089

- Molecule 2 is a protein called UBIQUITIN-CONJUGATING ENZYME E2 VARIANT 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	142	Total	C	N	O	S	0	0	0
			1123	709	195	211	8			
2	F	139	Total	C	N	O	S	0	0	0
			1109	701	192	208	8			
2	I	139	Total	C	N	O	S	0	0	0
			1109	701	192	208	8			
2	L	139	Total	C	N	O	S	0	0	0
			1109	701	192	208	8			

- Molecule 3 is a protein called CARBOXY TERMINUS OF HSP70-INTERACTING PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	S	75	Total	C	N	O	S	0	0	1
			598	379	101	114	4			
3	T	78	Total	C	N	O	S	0	0	0
			634	402	104	124	4			
3	U	73	Total	C	N	O	S	0	0	1
			577	365	98	110	4			
3	V	71	Total	C	N	O	S	0	0	1
			564	356	96	108	4			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	5	Total	O	0	0
			5	5		
4	C	5	Total	O	0	0
			5	5		
4	E	3	Total	O	0	0
			3	3		
4	F	7	Total	O	0	0
			7	7		
4	H	1	Total	O	0	0
			1	1		
4	I	1	Total	O	0	0
			1	1		
4	L	1	Total	O	0	0
			1	1		
4	S	6	Total	O	0	0
			6	6		
4	T	2	Total	O	0	0
			2	2		
4	U	1	Total	O	0	0
			1	1		
4	V	3	Total	O	0	0
			3	3		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: UBIQUITIN-CONJUGATING ENZYME E2 N




• Molecule 1: UBIQUITIN-CONJUGATING ENZYME E2 N




• Molecule 1: UBIQUITIN-CONJUGATING ENZYME E2 N








• Molecule 1: UBIQUITIN-CONJUGATING ENZYME E2 N




• Molecule 2: UBIQUITIN-CONJUGATING ENZYME E2 VARIANT 1




• Molecule 2: UBIQUITIN-CONJUGATING ENZYME E2 VARIANT 1




• Molecule 2: UBIQUITIN-CONJUGATING ENZYME E2 VARIANT 1







• Molecule 2: UBIQUITIN-CONJUGATING ENZYME E2 VARIANT 1




• Molecule 3: CARBOXY TERMINUS OF HSP70-INTERACTING PROTEIN



• Molecule 3: CARBOXY TERMINUS OF HSP70-INTERACTING PROTEIN



• Molecule 3: CARBOXY TERMINUS OF HSP70-INTERACTING PROTEIN



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	180.31Å 69.99Å 204.48Å 90.00° 106.95° 90.00°	Depositor
Resolution (Å)	196.12 – 2.90 37.40 – 2.90	Depositor EDS
% Data completeness (in resolution range)	95.3 (196.12-2.90) 95.4 (37.40-2.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.03 (at 2.90Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.214 , 0.297 0.214 , 0.290	Depositor DCC
R_{free} test set	2621 reflections (5.30%)	DCC
Wilson B-factor (Å ²)	64.0	Xtriage
Anisotropy	0.605	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 62.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 52095 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11636	wwPDB-VP
Average B, all atoms (Å ²)	4.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	B	3.41	160/1231 (13.0%)	2.68	86/1673 (5.1%)
1	E	3.07	115/1216 (9.5%)	2.33	62/1653 (3.8%)
1	H	2.87	99/1216 (8.1%)	2.29	61/1653 (3.7%)
1	K	3.07	114/1231 (9.3%)	2.29	65/1673 (3.9%)
2	C	3.59	140/1147 (12.2%)	2.73	88/1551 (5.7%)
2	F	3.13	108/1133 (9.5%)	2.43	75/1532 (4.9%)
2	I	2.61	68/1133 (6.0%)	2.17	49/1532 (3.2%)
2	L	2.82	92/1133 (8.1%)	2.32	62/1532 (4.0%)
3	S	3.46	83/612 (13.6%)	2.81	61/831 (7.3%)
3	T	3.20	78/649 (12.0%)	2.47	44/880 (5.0%)
3	U	3.14	60/589 (10.2%)	2.44	45/799 (5.6%)
3	V	3.32	74/576 (12.8%)	2.43	36/780 (4.6%)
All	All	3.13	1191/11866 (10.0%)	2.44	734/16089 (4.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	7
1	E	0	6
1	H	0	7
1	K	0	17
2	C	0	6
2	F	0	4
2	I	0	13
2	L	0	3
3	S	0	4
3	T	0	1
3	U	0	3
3	V	0	2
All	All	0	73

The worst 5 of 1191 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	131	ALA	CA-CB	30.67	2.16	1.52
1	B	142	GLU	CD-OE2	26.58	1.54	1.25
1	K	142	GLU	CD-OE2	22.60	1.50	1.25
1	H	65	GLU	CG-CD	20.68	1.82	1.51
3	S	243	GLU	CD-OE1	19.09	1.46	1.25

The worst 5 of 734 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	89	ARG	NE-CZ-NH2	-21.83	109.38	120.30
1	B	89	ARG	NE-CZ-NH1	20.31	130.45	120.30
2	I	43	ARG	NE-CZ-NH2	-19.23	110.69	120.30
1	B	74	ARG	NE-CZ-NH2	-18.82	110.89	120.30
1	B	74	ARG	NE-CZ-NH1	18.64	129.62	120.30

There are no chirality outliers.

5 of 73 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	29	ALA	Peptide
1	B	5	SER	Peptide
1	B	6	ALA	Peptide
1	B	76	MET	Peptide
1	B	97	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1202	0	1209	137	1
1	E	1187	0	1196	144	1
1	H	1187	0	1196	187	1
1	K	1202	0	1209	219	1
2	C	1123	0	1121	124	1
2	F	1109	0	1114	134	0
2	I	1109	0	1114	162	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	L	1109	0	1114	128	1
3	S	598	0	584	87	0
3	T	634	0	612	91	0
3	U	577	0	569	136	0
3	V	564	0	557	108	0
4	B	5	0	0	0	0
4	C	5	0	0	0	0
4	E	3	0	0	0	0
4	F	7	0	0	1	0
4	H	1	0	0	0	0
4	I	1	0	0	0	0
4	L	1	0	0	0	0
4	S	6	0	0	0	0
4	T	2	0	0	1	0
4	U	1	0	0	0	0
4	V	3	0	0	1	0
All	All	11636	0	11595	1587	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 68.

The worst 5 of 1587 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:96:LYS:CD	1:E:96:LYS:CE	1.74	1.65
3:U:228:ILE:CB	3:U:228:ILE:CG2	1.74	1.64
3:V:288:LYS:CG	3:V:288:LYS:CD	1.74	1.63
1:E:8:LEU:CD1	1:E:8:LEU:CG	1.75	1.63
1:H:110:LEU:CG	1:H:110:LEU:CD2	1.77	1.62

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:3:ALA:O	2:C:142:TYR:OH[2_555]	2.08	0.12
1:E:65:GLU:OE1	2:L:158:LYS:NZ[2_556]	2.16	0.04
1:H:142:GLU:OE2	1:K:145:ARG:NH1[3_434]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	148/154 (96%)	114 (77%)	28 (19%)	6 (4%)	3	14
1	E	145/154 (94%)	122 (84%)	13 (9%)	10 (7%)	1	4
1	H	145/154 (94%)	103 (71%)	23 (16%)	19 (13%)	0	1
1	K	148/154 (96%)	92 (62%)	27 (18%)	29 (20%)	0	0
2	C	140/142 (99%)	114 (81%)	12 (9%)	14 (10%)	1	2
2	F	137/142 (96%)	113 (82%)	12 (9%)	12 (9%)	1	2
2	I	137/142 (96%)	94 (69%)	20 (15%)	23 (17%)	0	0
2	L	137/142 (96%)	106 (77%)	17 (12%)	14 (10%)	1	2
3	S	73/78 (94%)	47 (64%)	15 (20%)	11 (15%)	0	0
3	T	76/78 (97%)	52 (68%)	14 (18%)	10 (13%)	0	1
3	U	71/78 (91%)	49 (69%)	14 (20%)	8 (11%)	0	1
3	V	69/78 (88%)	47 (68%)	10 (14%)	12 (17%)	0	0
All	All	1426/1496 (95%)	1053 (74%)	205 (14%)	168 (12%)	0	1

5 of 168 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	5	SER
1	B	137	GLU
2	C	34	THR
2	C	54	GLY
2	C	55	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	128/129 (99%)	91 (71%)	37 (29%)	0	1
1	E	127/129 (98%)	94 (74%)	33 (26%)	0	2
1	H	127/129 (98%)	97 (76%)	30 (24%)	1	2
1	K	128/129 (99%)	90 (70%)	38 (30%)	0	1
2	C	125/127 (98%)	100 (80%)	25 (20%)	1	5
2	F	125/127 (98%)	98 (78%)	27 (22%)	1	3
2	I	125/127 (98%)	85 (68%)	40 (32%)	0	1
2	L	125/127 (98%)	92 (74%)	33 (26%)	0	2
3	S	68/72 (94%)	49 (72%)	19 (28%)	0	1
3	T	72/72 (100%)	55 (76%)	17 (24%)	1	2
3	U	66/72 (92%)	56 (85%)	10 (15%)	3	10
3	V	65/72 (90%)	34 (52%)	31 (48%)	0	0
All	All	1281/1312 (98%)	941 (74%)	340 (26%)	0	2

5 of 340 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	I	38	VAL
2	I	166	PRO
3	V	240	LEU
2	I	66	ASP
2	I	103	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 41 such sidechains are listed below:

Mol	Chain	Res	Type
1	H	17	GLN
2	I	89	ASN
3	U	280	GLN
1	H	81	HIS
1	H	83	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	152/154 (98%)	-0.19	3 (1%) 68 64	2, 2, 12, 28	0
1	E	149/154 (96%)	-0.28	2 (1%) 79 78	2, 2, 18, 32	0
1	H	149/154 (96%)	-0.03	5 (3%) 49 41	2, 2, 20, 27	0
1	K	152/154 (98%)	0.46	17 (11%) 7 4	2, 3, 29, 38	0
2	C	142/142 (100%)	-0.00	2 (1%) 78 76	2, 2, 10, 28	0
2	F	139/142 (97%)	-0.26	1 (0%) 89 88	2, 2, 10, 20	0
2	I	139/142 (97%)	-0.01	2 (1%) 78 76	2, 2, 10, 17	0
2	L	139/142 (97%)	-0.27	0 100 100	2, 2, 10, 13	0
3	S	75/78 (96%)	-0.32	0 100 100	2, 2, 17, 21	0
3	T	78/78 (100%)	0.16	6 (7%) 16 11	2, 2, 50, 59	0
3	U	73/78 (93%)	-0.29	1 (1%) 78 76	2, 2, 11, 17	0
3	V	71/78 (91%)	-0.40	1 (1%) 78 76	2, 3, 14, 27	0
All	All	1458/1496 (97%)	-0.10	40 (2%) 58 52	2, 2, 18, 59	0

The worst 5 of 40 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	K	4	GLY	7.9
1	K	3	ALA	6.6
1	K	156	ILE	5.3
1	E	6	ALA	4.9
1	K	124	PRO	4.8

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.