



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:14 AM GMT

PDB ID : 2C7Z
Title : PLANT ENZYME CRYSTAL FORM II
Authors : Sundaramoorthy, R.; Micossi, E.; Alphey, M.S.; Leonard, G.A.; Hunter, W.N.
Deposited on : 2005-11-30
Resolution : 2.37 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4019 (2.40-2.36)
Clashscore	102246	4595 (2.40-2.36)
Ramachandran outliers	100387	4520 (2.40-2.36)
Sidechain outliers	100360	4522 (2.40-2.36)
RSRZ outliers	91569	4034 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	404	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3002 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-KETOACYL-COA THIOLASE 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	396	Total	C	N	O	S	34	0	0
			2900	1816	514	551	19			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	102	Total	O	0	0
			102	102		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 3-KETOACYL-COA THIOLASE 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	73.39Å 95.82Å 56.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	18.54 – 2.37 18.54 – 2.37	Depositor EDS
% Data completeness (in resolution range)	87.4 (18.54-2.37) 87.4 (18.54-2.37)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.21 (at 2.38Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.198 , 0.276 0.197 , 0.278	Depositor DCC
R_{free} test set	725 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	42.1	Xtriage
Anisotropy	0.358	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 66.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 14473 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3002	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	6/2944 (0.2%)	0.72	8/3983 (0.2%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	254	LYS	CB-CG	-21.54	0.94	1.52
1	A	206	ARG	CB-CG	-13.16	1.17	1.52
1	A	275	LYS	CB-CG	9.24	1.77	1.52
1	A	205	GLN	CG-CD	-9.20	1.29	1.51
1	A	282	LYS	CG-CD	-5.56	1.33	1.52
1	A	246	VAL	CB-CG1	5.12	1.63	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	254	LYS	CA-CB-CG	13.26	142.56	113.40
1	A	254	LYS	CB-CG-CD	10.19	138.10	111.60
1	A	282	LYS	CB-CG-CD	8.22	132.98	111.60
1	A	252	ASP	CB-CG-OD2	5.85	123.57	118.30
1	A	280	PHE	CB-CG-CD1	-5.66	116.84	120.80
1	A	280	PHE	CB-CG-CD2	5.49	124.64	120.80
1	A	205	GLN	CB-CG-CD	5.22	125.17	111.60
1	A	205	GLN	CG-CD-OE1	-5.15	111.29	121.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2900	0	2939	41	0
2	A	102	0	0	6	0
All	All	3002	0	2939	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:85:ALA:O	1:A:89:LYS:HD3	1.79	0.82
1:A:114:SER:HB2	1:A:175:TRP:HB3	1.65	0.79
1:A:184:LYS:N	2:A:2039:HOH:O	2.21	0.73
1:A:107:ALA:HB3	1:A:112:ARG:HG2	1.72	0.71
1:A:212:GLN:HG2	1:A:213:GLU:H	1.60	0.66
1:A:274:GLY:HA3	2:A:2059:HOH:O	1.95	0.64
1:A:371:CYS:SG	2:A:2074:HOH:O	2.55	0.64
1:A:50:ASP:O	1:A:306:ARG:NH1	2.31	0.64
1:A:187:ALA:O	1:A:191:ASN:HB2	1.98	0.64
1:A:65:SER:HB3	1:A:168:MET:HG3	1.80	0.62
1:A:82:VAL:HG12	1:A:300:ALA:HB3	1.81	0.61
1:A:212:GLN:CG	1:A:213:GLU:N	2.64	0.60
1:A:246:VAL:HG22	1:A:253:GLU:HG2	1.84	0.60
1:A:166:GLU:HG3	1:A:395:LEU:HB2	1.84	0.59
1:A:64:LYS:HE2	1:A:293:GLN:HB2	1.84	0.58
1:A:207:PHE:O	2:A:2042:HOH:O	2.17	0.58
1:A:212:GLN:CG	1:A:213:GLU:H	2.20	0.54
1:A:75:PRO:HG3	1:A:167:SER:HB3	1.89	0.54
1:A:200:SER:HB3	1:A:366:SER:OG	2.09	0.53
1:A:190:GLN:HB2	2:A:2040:HOH:O	2.09	0.52
1:A:378:ASP:HB3	1:A:381:LYS:HD2	1.91	0.52
1:A:335:ILE:HA	2:A:2074:HOH:O	2.10	0.51
1:A:399:GLY:O	1:A:403:VAL:HG23	2.11	0.51
1:A:65:SER:HB3	1:A:168:MET:CG	2.42	0.49
1:A:200:SER:OG	1:A:367:GLN:HB2	2.12	0.49
1:A:43:GLN:NE2	1:A:45:THR:OG1	2.46	0.49
1:A:406:LEU:HD22	1:A:422:VAL:HG23	1.95	0.49
1:A:287:THR:HG22	1:A:288:ALA:N	2.29	0.47
1:A:164:GLY:HA3	1:A:397:ALA:HA	1.98	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:379:PRO:HA	1:A:382:ILE:HD12	1.99	0.45
1:A:53:VAL:HG22	1:A:304:MET:O	2.17	0.44
1:A:118:MET:HG2	1:A:175:TRP:HB2	2.00	0.44
1:A:190:GLN:HA	1:A:193:LEU:HB2	2.00	0.43
1:A:58:HIS:NE2	1:A:89:LYS:HE3	2.33	0.43
1:A:201:GLU:OE2	1:A:284:GLY:HA3	2.19	0.42
1:A:328:VAL:HG11	1:A:336:GLY:HA2	2.01	0.42
1:A:107:ALA:CB	1:A:112:ARG:HG2	2.47	0.42
1:A:378:ASP:OD2	1:A:380:GLU:HB3	2.19	0.42
1:A:247:ASP:OD2	1:A:250:THR:HG23	2.20	0.41
1:A:55:VAL:HG23	1:A:302:LEU:HD23	2.02	0.40
1:A:87:ILE:HG23	1:A:92:LEU:O	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	392/404 (97%)	368 (94%)	22 (6%)	2 (0%)	34	46

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	109	GLY
1	A	395	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	300/307 (98%)	287 (96%)	13 (4%)	35 52

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	SER
1	A	44	ARG
1	A	51	ASP
1	A	89	LYS
1	A	93	ASN
1	A	188	GLN
1	A	212	GLN
1	A	244	LYS
1	A	254	LYS
1	A	277	LYS
1	A	308	VAL
1	A	320	PHE
1	A	333	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	43	GLN
1	A	93	ASN
1	A	212	GLN
1	A	214	GLN
1	A	361	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	396/404 (98%)	1.09	85 (21%) 1 1	35, 44, 63, 67	8 (2%)

All (85) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	174	ALA	9.4
1	A	273	LEU	8.4
1	A	110	SER	7.4
1	A	278	PRO	6.9
1	A	173	MET	6.7
1	A	67	ARG	6.2
1	A	281	LYS	6.1
1	A	172	PRO	6.0
1	A	209	VAL	5.8
1	A	208	GLY	5.5
1	A	274	GLY	5.4
1	A	108	PRO	5.1
1	A	204	ALA	5.1
1	A	197	GLY	5.1
1	A	207	PHE	4.9
1	A	284	GLY	4.9
1	A	186	PHE	4.3
1	A	199	THR	4.3
1	A	194	LEU	4.2
1	A	200	SER	4.1
1	A	313	GLY	3.9
1	A	111	GLN	3.9
1	A	190	GLN	3.9
1	A	276	LEU	3.8
1	A	268	THR	3.7
1	A	170	THR	3.6
1	A	212	GLN	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	112	ARG	3.5
1	A	255	PRO	3.4
1	A	283	ASP	3.4
1	A	206	ARG	3.4
1	A	198	VAL	3.4
1	A	277	LYS	3.3
1	A	109	GLY	3.3
1	A	203	VAL	3.3
1	A	260	VAL	3.2
1	A	196	MET	3.2
1	A	175	TRP	3.2
1	A	367	GLN	3.1
1	A	202	ASN	3.1
1	A	332	ILE	3.0
1	A	370	TYR	2.9
1	A	414	GLY	2.9
1	A	267	THR	2.8
1	A	275	LYS	2.8
1	A	280	PHE	2.7
1	A	302	LEU	2.7
1	A	74	TYR	2.7
1	A	51	ASP	2.7
1	A	366	SER	2.7
1	A	52	VAL	2.6
1	A	271	ALA	2.6
1	A	441	GLY	2.6
1	A	369	VAL	2.6
1	A	107	ALA	2.5
1	A	106	LEU	2.5
1	A	368	PHE	2.5
1	A	229	THR	2.5
1	A	251	GLY	2.4
1	A	176	GLU	2.4
1	A	427	GLY	2.4
1	A	311	GLN	2.4
1	A	287	THR	2.4
1	A	187	ALA	2.3
1	A	66	LYS	2.3
1	A	374	LYS	2.3
1	A	210	SER	2.3
1	A	282	LYS	2.3
1	A	320	PHE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	351	GLU	2.3
1	A	205	GLN	2.2
1	A	211	ARG	2.2
1	A	285	THR	2.2
1	A	303	LEU	2.2
1	A	242	LYS	2.2
1	A	50	ASP	2.1
1	A	335	ILE	2.1
1	A	249	LYS	2.1
1	A	201	GLU	2.1
1	A	217	ALA	2.1
1	A	252	ASP	2.0
1	A	415	LYS	2.0
1	A	83	LEU	2.0
1	A	165	LEU	2.0
1	A	436	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.