



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 12, 2016 – 08:23 PM EDT

PDB ID : 5C7J
Title : CRYSTAL STRUCTURE OF NEDD4 WITH A UB VARIANT
Authors : Walker, J.R.; Hu, J.; Dong, A.; Bountra, C.; Edwards, A.M.; Arrowsmith, C.H.; Tong, Y.; STRUCTURAL GENOMICS CONSORTIUM (SGC)
Deposited on : 2015-06-24
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.


The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027107
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027107

i

X-RAY DIFFRACTION

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	382	<div> <div>3%</div> <div>93%</div> <div>5%</div> </div>
1	B	382	<div> <div>2%</div> <div>92%</div> <div>5%</div> </div>
2	C	92	<div> <div>3%</div> <div>90%</div> <div>8%</div> </div>
2	D	92	<div> <div>7%</div> <div>88%</div> <div>8%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7161 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 ubiquitin-protein ligase NEDD4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	373	Total	C	N	O	S	0	1	0
			2962	1909	490	549	14			
1	B	373	Total	C	N	O	S	0	1	0
			2958	1908	491	545	14			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	519	GLY	-	expression tag	UNP P46934
B	519	GLY	-	expression tag	UNP P46934

- Molecule 2 is a protein called Polyubiquitin-C.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	85	Total	C	N	O	0	0	0
			616	386	110	120			
2	D	85	Total	C	N	O	0	1	0
			621	389	109	123			

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-17	ALA	-	expression tag	UNP P0CG48
C	-16	HIS	-	expression tag	UNP P0CG48
C	-15	HIS	-	expression tag	UNP P0CG48
C	-14	HIS	-	expression tag	UNP P0CG48
C	-13	HIS	-	expression tag	UNP P0CG48
C	-12	HIS	-	expression tag	UNP P0CG48
C	-11	HIS	-	expression tag	UNP P0CG48
C	-10	VAL	-	expression tag	UNP P0CG48
C	-9	THR	-	expression tag	UNP P0CG48
C	-8	SER	-	expression tag	UNP P0CG48

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-7	LEU	-	expression tag	UNP P0CG48
C	-6	TYR	-	expression tag	UNP P0CG48
C	-5	LYS	-	expression tag	UNP P0CG48
C	-4	LYS	-	expression tag	UNP P0CG48
C	-3	ALA	-	expression tag	UNP P0CG48
C	-2	GLY	-	expression tag	UNP P0CG48
C	-1	SER	-	expression tag	UNP P0CG48
C	0	THR	-	expression tag	UNP P0CG48
C	9	ALA	THR	engineered mutation	UNP P0CG48
C	11	TRP	LYS	engineered mutation	UNP P0CG48
C	12	GLY	THR	engineered mutation	UNP P0CG48
C	62	ARG	GLN	engineered mutation	UNP P0CG48
C	63	TYR	LYS	engineered mutation	UNP P0CG48
C	64	ASP	GLU	engineered mutation	UNP P0CG48
C	66	GLN	THR	engineered mutation	UNP P0CG48
C	71	GLY	LEU	engineered mutation	UNP P0CG48
D	-17	ALA	-	expression tag	UNP P0CG48
D	-16	HIS	-	expression tag	UNP P0CG48
D	-15	HIS	-	expression tag	UNP P0CG48
D	-14	HIS	-	expression tag	UNP P0CG48
D	-13	HIS	-	expression tag	UNP P0CG48
D	-12	HIS	-	expression tag	UNP P0CG48
D	-11	HIS	-	expression tag	UNP P0CG48
D	-10	VAL	-	expression tag	UNP P0CG48
D	-9	THR	-	expression tag	UNP P0CG48
D	-8	SER	-	expression tag	UNP P0CG48
D	-7	LEU	-	expression tag	UNP P0CG48
D	-6	TYR	-	expression tag	UNP P0CG48
D	-5	LYS	-	expression tag	UNP P0CG48
D	-4	LYS	-	expression tag	UNP P0CG48
D	-3	ALA	-	expression tag	UNP P0CG48
D	-2	GLY	-	expression tag	UNP P0CG48
D	-1	SER	-	expression tag	UNP P0CG48
D	0	THR	-	expression tag	UNP P0CG48
D	9	ALA	THR	engineered mutation	UNP P0CG48
D	11	TRP	LYS	engineered mutation	UNP P0CG48
D	12	GLY	THR	engineered mutation	UNP P0CG48
D	62	ARG	GLN	engineered mutation	UNP P0CG48
D	63	TYR	LYS	engineered mutation	UNP P0CG48
D	64	ASP	GLU	engineered mutation	UNP P0CG48
D	66	GLN	THR	engineered mutation	UNP P0CG48
D	71	GLY	LEU	engineered mutation	UNP P0CG48


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total 2	O 2	0	0
3	B	2	Total 2	O 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: E3 ubiquitin-protein ligase NEDD4




- Molecule 1: E3 ubiquitin-protein ligase NEDD4



- Molecule 2: Polyubiquitin-C



- Molecule 2: Polyubiquitin-C



4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	139.87Å 139.87Å 59.39Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.01 – 3.00 45.78 – 3.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (50.01-3.00) 100.0 (45.78-3.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.02 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.8.0123	Depositor
R, R_{free}	0.227 , 0.277 0.227 , 0.268	Depositor DCC
R_{free} test set	549 reflections (2.16%)	DCC
Wilson B-factor (Å ²)	97.5	Xtriage
Anisotropy	0.038	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 74.1	EDS
Estimated twinning fraction	0.476 for -h,-k,l 0.027 for h,-h-k,-l 0.028 for -k,-h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 26023 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7161	wwPDB-VP
Average B, all atoms (Å ²)	100.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/3044	0.57	0/4133
1	B	0.40	0/3041	0.57	0/4128
2	C	0.39	0/628	0.56	0/856
2	D	0.39	0/634	0.56	0/866
All	All	0.40	0/7347	0.57	0/9983

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2962	0	2641	8	0
1	B	2958	0	2638	9	0
2	C	616	0	528	1	0
2	D	621	0	524	2	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
All	All	7161	0	6331	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:642:ALA:HA	1:A:652:LEU:HD11	1.90	0.54
1:B:642:ALA:HA	1:B:652:LEU:HD11	1.90	0.53
1:B:561:MET:CE	1:B:640:ARG:HD2	2.44	0.47
1:A:561:MET:CE	1:A:640:ARG:HD2	2.44	0.46
1:A:658:ARG:N	1:A:659:PRO:HD2	2.31	0.45
1:B:704:GLU:OE1	1:B:744:ARG:NH2	2.50	0.45
1:B:658:ARG:N	1:B:659:PRO:HD2	2.32	0.44
1:A:704:GLU:OE1	1:A:744:ARG:NH2	2.50	0.44
1:A:855:TRP:HB3	1:A:872:ASP:HB3	2.01	0.43
1:A:780:LEU:HB2	1:A:833:VAL:HG11	2.01	0.43
1:A:757:PHE:HA	1:A:760:ILE:O	2.19	0.43
1:B:757:PHE:HA	1:B:760:ILE:O	2.19	0.42
1:B:827:VAL:HG11	1:B:850:PHE:CE1	2.55	0.42
1:B:855:TRP:HB3	1:B:872:ASP:HB3	2.01	0.42
2:C:-11:HIS:O	2:C:-8:SER:OG	2.36	0.42
1:B:627:CYS:HB3	2:D:70:VAL:HG21	2.02	0.41
1:A:827:VAL:HG11	1:A:850:PHE:CE1	2.55	0.41
2:D:-11:HIS:O	2:D:-8:SER:OG	2.36	0.41
1:B:702:ILE:HG21	1:B:739:TRP:CZ3	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	370/382 (97%)	360 (97%)	10 (3%)	0	100	100
1	B	370/382 (97%)	360 (97%)	10 (3%)	0	100	100
2	C	81/92 (88%)	77 (95%)	4 (5%)	0	100	100
2	D	82/92 (89%)	78 (95%)	4 (5%)	0	100	100
All	All	903/948 (95%)	875 (97%)	28 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	285/342 (83%)	282 (99%)	3 (1%)	80	94
1	B	283/342 (83%)	280 (99%)	3 (1%)	80	94
2	C	53/80 (66%)	53 (100%)	0	100	100
2	D	55/80 (69%)	54 (98%)	1 (2%)	66	91
All	All	676/844 (80%)	669 (99%)	7 (1%)	82	95

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	712	GLN
1	A	838	PHE
1	A	869	ASN
1	B	712	GLN
1	B	838	PHE
1	B	869	ASN
2	D	-15	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	854	GLN
1	B	854	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	373/382 (97%)	0.07	10 (2%) 58 28	65, 94, 136, 167	0
1	B	373/382 (97%)	0.08	8 (2%) 67 36	66, 95, 134, 168	0
2	C	85/92 (92%)	0.16	3 (3%) 48 21	84, 115, 152, 169	0
2	D	85/92 (92%)	0.17	6 (7%) 19 7	86, 115, 153, 166	0
All	All	916/948 (96%)	0.09	27 (2%) 55 26	65, 98, 147, 169	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	2	GLN	4.7
2	D	3	ILE	3.8
2	C	15	LEU	3.8
2	C	19	PRO	3.6
1	B	871	LEU	3.0
1	A	871	LEU	2.9
1	A	834	PRO	2.8
2	C	72	ARG	2.7
1	B	656	PHE	2.6
2	D	15	LEU	2.5
1	B	867	CYS	2.5
1	A	852	VAL	2.4
1	A	783	VAL	2.3
1	A	788	TRP	2.3
1	A	618	LEU	2.2
2	D	69	LEU	2.2
1	B	606	GLY	2.2
1	B	781	GLY	2.1
1	A	868	PHE	2.1
1	A	850	PHE	2.1
1	A	839	ALA	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	839	ALA	2.1
2	D	8	LEU	2.1
2	D	67	LEU	2.1
1	A	702	ILE	2.1
1	B	618	LEU	2.1
1	B	707	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.