



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:47 AM GMT

PDB ID : 3C87
Title : Crystal structure of the enterobactin esterase FES from *Shigella flexneri* in the presence of enterobactin
Authors : Kim, Y.; Maltseva, N.; Abergel, R.; Raymond, K.; Holzle, D.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2008-02-11
Resolution : 2.17 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	5130 (2.20-2.16)
Clashscore	102246	5965 (2.20-2.16)
Ramachandran outliers	100387	5863 (2.20-2.16)
Sidechain outliers	100360	5864 (2.20-2.16)
RSRZ outliers	91569	5142 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	403	<div> <div>4%</div> <div>83%</div> <div>13%</div> <div>• •</div> </div>
1	B	403	<div> <div>4%</div> <div>83%</div> <div>10%</div> <div>• •</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6697 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Enterochelin esterase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	388	Total	C	N	O	S	Se	0	5	0
			3178	2040	558	567	6	7			
1	B	385	Total	C	N	O	S	Se	0	10	0
			3211	2058	568	572	6	7			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	EXPRESSION TAG	UNP Q83SB9
A	-1	ASN	-	EXPRESSION TAG	UNP Q83SB9
A	0	ALA	-	EXPRESSION TAG	UNP Q83SB9
B	-2	SER	-	EXPRESSION TAG	UNP Q83SB9
B	-1	ASN	-	EXPRESSION TAG	UNP Q83SB9
B	0	ALA	-	EXPRESSION TAG	UNP Q83SB9

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Na	0	0
			1	1		

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	153	Total	O	0	0
			153	153		
4	B	149	Total	O	0	0
			149	149		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Enterochelin esterase



- Molecule 1: Enterochelin esterase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	47.30Å 113.25Å 151.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.03 – 2.17 46.03 – 2.17	Depositor EDS
% Data completeness (in resolution range)	98.9 (46.03-2.17) 98.9 (46.03-2.17)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.03 (at 2.18Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.190 , 0.234 0.189 , 0.234	Depositor DCC
R_{free} test set	2189 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	36.0	Xtriage
Anisotropy	0.026	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 44.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 43438 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6697	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.32 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.6917e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.81	0/3270	0.80	2/4449 (0.0%)
1	B	0.85	0/3304	0.80	4/4492 (0.1%)
All	All	0.83	0/6574	0.80	6/8941 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	233	ASP	CB-CG-OD1	7.06	124.65	118.30
1	B	395	LEU	CA-CB-CG	5.84	128.73	115.30
1	B	261	LEU	CA-CB-CG	5.57	128.10	115.30
1	A	233	ASP	CB-CG-OD2	-5.53	113.33	118.30
1	B	233	ASP	CB-CG-OD1	5.28	123.05	118.30
1	B	273	ASP	CB-CG-OD1	5.11	122.90	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3178	0	3078	44	0
1	B	3211	0	3093	47	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	5	0	0	0	0
4	A	153	0	0	4	0
4	B	149	0	0	2	0
All	All	6697	0	6171	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:22:ARG:HH22	1:A:148:GLN:HE21	1.02	1.00
1:A:22:ARG:HH22	1:A:148:GLN:NE2	1.63	0.95
1:B:22:ARG:HH22	1:B:148:GLN:HE21	0.95	0.94
1:A:336:ARG:HD3	4:A:584:HOH:O	1.68	0.92
1:B:59:HIS:HE1	1:B:344[B]:ARG:O	1.58	0.86
1:A:349:MSE:HE3	1:A:353:GLN:HG3	1.60	0.81
1:B:22:ARG:HH22	1:B:148:GLN:NE2	1.74	0.81
1:A:316:GLN:HE21	1:B:137:LEU:HD21	1.48	0.77
1:B:212:TRP:HB2	1:B:213:PRO:HD3	1.68	0.74
1:B:55:GLY:O	1:B:344[B]:ARG:HG2	1.91	0.70
1:B:16[A]:HIS:CE1	4:B:626:HOH:O	2.44	0.70
1:B:30:VAL:HG21	1:B:83:LEU:HD12	1.75	0.69
1:A:233:ASP:OD2	4:A:562:HOH:O	2.11	0.68
1:A:316:GLN:HA	1:B:117:ARG:HG2	1.74	0.68
1:A:349:MSE:CE	1:A:353:GLN:HG3	2.24	0.68
1:B:318:GLU:OE1	1:B:326:LYS:HE3	1.95	0.67
1:B:22:ARG:NH2	1:B:148:GLN:HE21	1.80	0.67
1:A:346:PRO:O	1:A:350:ARG:HG3	1.96	0.66
1:A:390:ASP:O	1:A:393[A]:GLN:HG2	1.97	0.65
1:A:22:ARG:NH2	1:A:148:GLN:HE21	1.85	0.64
1:A:349:MSE:HE1	1:A:368:TRP:CH2	2.34	0.62
1:B:192:GLU:O	1:B:193:GLU:HB2	2.00	0.62
1:B:27:MSE:HE3	1:B:82[A]:GLN:HB3	1.82	0.61
1:B:313:ARG:H	1:B:313:ARG:HD3	1.66	0.60
1:A:30:VAL:HG21	1:A:83:LEU:HD12	1.83	0.60
1:B:393:GLN:HE21	1:B:393:GLN:HA	1.67	0.60
1:A:349:MSE:CE	1:A:353:GLN:CG	2.80	0.58
1:A:349:MSE:HE1	1:A:368:TRP:CZ2	2.39	0.58
1:A:347:MSE:HE1	1:B:313:ARG:HH22	1.68	0.57
1:B:188:ASP:OD1	1:B:188:ASP:N	2.38	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:68:MSE:HG2	1:A:79:TRP:HB2	1.87	0.56
1:A:167:GLU:OE2	1:A:180:ARG:HG2	2.05	0.56
1:A:22:ARG:NH2	1:A:148:GLN:NE2	2.45	0.56
1:B:221[B]:ARG:C	1:B:222:GLN:HG2	2.28	0.54
1:A:108:ASP:HB3	1:A:111:GLU:HG3	1.90	0.54
1:A:121:PRO:HD3	4:A:567:HOH:O	2.08	0.53
1:B:171:LYS:HG2	1:B:178:SER:HB3	1.89	0.53
1:B:309[B]:TRP:HE3	1:B:347:MSE:SE	2.41	0.53
1:B:109:ARG:HD2	1:B:109:ARG:N	2.25	0.52
1:B:134:LYS:HG3	4:B:547:HOH:O	2.10	0.51
1:A:203:GLU:HG3	4:A:592:HOH:O	2.10	0.51
1:B:87:TRP:HE1	1:B:344[A]:ARG:CZ	2.23	0.51
1:B:68:MSE:HE2	1:B:78:GLN:HA	1.90	0.51
1:A:349:MSE:HE2	1:A:353:GLN:CG	2.41	0.51
1:B:313:ARG:HH21	1:B:313:ARG:HG2	1.75	0.51
1:B:68:MSE:HG2	1:B:79:TRP:HB2	1.92	0.50
1:B:344[B]:ARG:NH2	1:B:372:ASP:OD1	2.45	0.50
1:B:27:MSE:CE	1:B:82[B]:GLN:HG3	2.42	0.49
1:A:309:TRP:CH2	1:B:313:ARG:HD2	2.48	0.49
1:A:362:ILE:CG2	1:A:362:ILE:O	2.60	0.49
1:A:313:ARG:HE	1:B:313:ARG:HB3	1.78	0.48
1:A:369:ARG:NH1	1:A:390:ASP:OD2	2.35	0.48
1:B:221[A]:ARG:C	1:B:222:GLN:HG2	2.34	0.48
1:B:109:ARG:HD2	1:B:109:ARG:H	1.78	0.48
1:A:347:MSE:HE1	1:B:313:ARG:NH2	2.29	0.47
1:A:345:GLU:OE2	1:A:375:HIS:HD2	1.97	0.47
1:B:212:TRP:HB2	1:B:213:PRO:CD	2.39	0.47
1:A:376:ASP:O	1:A:379[B]:CYS:HB3	2.16	0.46
1:B:179:ARG:HD3	1:B:233:ASP:O	2.15	0.46
1:B:59:HIS:CE1	1:B:344[B]:ARG:O	2.51	0.45
1:A:281:SER:HA	1:A:305:SER:O	2.16	0.45
1:B:87:TRP:NE1	1:B:344[A]:ARG:CZ	2.79	0.45
1:A:316:GLN:NE2	1:B:137:LEU:HD21	2.23	0.45
1:A:313:ARG:HB2	1:B:313:ARG:HG2	1.99	0.45
1:A:345:GLU:OE2	1:A:375:HIS:CD2	2.70	0.44
1:B:326:LYS:HG2	1:B:358:GLN:CD	2.38	0.44
1:B:309[B]:TRP:HD1	1:B:319:GLY:H	1.66	0.44
1:B:87:TRP:HZ3	1:B:144:LEU:HB3	1.83	0.43
1:B:59:HIS:HE1	1:B:344[A]:ARG:O	2.01	0.43
1:A:349:MSE:SE	1:A:370:GLN:HE21	2.51	0.43
1:A:133:TRP:CD2	1:A:209:MSE:HE3	2.54	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:109:ARG:HH11	1:A:113:ARG:NH2	2.17	0.42
1:A:12:TRP:CH2	1:A:35:ARG:HB2	2.55	0.42
1:A:64:GLN:HA	1:A:65:PRO:HD3	1.86	0.42
1:B:155:TRP:CZ2	1:B:381:ARG:HB2	2.54	0.42
1:A:363:LYS:HB3	1:A:363:LYS:HE2	1.82	0.42
1:A:362:ILE:HA	1:A:362:ILE:HD13	1.75	0.42
1:A:313:ARG:CB	1:B:313:ARG:HG2	2.50	0.42
1:B:158:PRO:HB2	1:B:213:PRO:HB2	2.01	0.41
1:A:373:GLY:HA3	1:A:379[A]:CYS:SG	2.60	0.41
1:B:221[B]:ARG:O	1:B:222:GLN:HG2	2.19	0.41
1:A:59:HIS:ND1	1:A:59:HIS:N	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	385/403 (96%)	380 (99%)	5 (1%)	0	100	100
1	B	387/403 (96%)	380 (98%)	7 (2%)	0	100	100
All	All	772/806 (96%)	760 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	335/333 (101%)	322 (96%)	13 (4%)	39	45
1	B	337/333 (101%)	318 (94%)	19 (6%)	26	27
All	All	672/666 (101%)	640 (95%)	32 (5%)	33	35

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	SER
1	A	13[A]	GLN
1	A	13[B]	GLN
1	A	26	GLU
1	A	62	ASN
1	A	82[A]	GLN
1	A	82[B]	GLN
1	A	137	LEU
1	A	188	ASP
1	A	189	VAL
1	A	296	GLU
1	A	362	ILE
1	A	369	ARG
1	B	10	SER
1	B	64	GLN
1	B	78	GLN
1	B	109	ARG
1	B	117	ARG
1	B	118	LYS
1	B	137	LEU
1	B	166	LYS
1	B	171	LYS
1	B	173	GLU
1	B	188	ASP
1	B	192	GLU
1	B	193	GLU
1	B	222	GLN
1	B	313	ARG
1	B	322	LEU
1	B	339	LEU
1	B	393	GLN
1	B	395	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	21	GLN
1	A	148	GLN
1	A	207	GLN
1	A	242	HIS
1	A	316	GLN
1	A	352	ASN
1	A	370	GLN
1	A	375	HIS
1	B	59	HIS
1	B	84	ASN
1	B	139	HIS
1	B	148	GLN
1	B	358	GLN
1	B	393	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	B	502	-	4,4,4	0.17	0	6,6,6	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PO4	B	502	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	381/403 (94%)	0.14	17 (4%) 37 39	26, 37, 59, 74	0
1	B	378/403 (93%)	0.10	17 (4%) 37 39	24, 33, 62, 97	0
All	All	759/806 (94%)	0.12	34 (4%) 37 39	24, 35, 59, 97	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	190	THR	7.9
1	A	189	VAL	6.6
1	B	188	ASP	5.3
1	A	190	THR	5.1
1	A	62	ASN	4.9
1	A	334	GLY	4.1
1	A	58	ASP	3.6
1	B	313	ARG	3.5
1	B	4	LEU	3.4
1	B	189	VAL	3.3
1	B	109	ARG	3.1
1	A	4	LEU	3.1
1	B	113	ARG	3.0
1	A	43	TYR	3.0
1	A	59	HIS	2.9
1	B	63	SER	2.8
1	B	362	ILE	2.8
1	A	108	ASP	2.6
1	A	188	ASP	2.5
1	A	112	LEU	2.5
1	A	26	GLU	2.5
1	A	313	ARG	2.5
1	B	192	GLU	2.4
1	B	309[A]	TRP	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	112	LEU	2.3
1	A	63	SER	2.3
1	A	6	VAL	2.3
1	A	136	GLY	2.2
1	A	113	ARG	2.2
1	B	110	LEU	2.1
1	B	396	PHE	2.1
1	B	316	GLN	2.1
1	B	25	ASP	2.0
1	B	58[A]	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	PO4	B	502	5/5	0.94	0.12	-0.26	66,67,68,69	0
2	NA	A	501	1/1	0.95	0.14	-	43,43,43,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.