



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:22 PM GMT

PDB ID : 4CAN
Title : Structure of rat neuronal nitric oxide synthase heme domain in complex with
7-(2-(3-Fluorobenzylamino)ethyl)quinolin-2-amine
Authors : Li, H.; Poulos, T.L.
Deposited on : 2013-10-08
Resolution : 1.91 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	5832 (1.94-1.90)
Clashscore	102246	6540 (1.94-1.90)
Ramachandran outliers	100387	6464 (1.94-1.90)
Sidechain outliers	100360	6465 (1.94-1.90)
RSRZ outliers	91569	5846 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	<div> <div>18%</div> <div>82%</div> <div>14%</div> <div>..</div> </div>
1	B	422	<div> <div>7%</div> <div>85%</div> <div>11%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	M7K	B	800	-	-	-	X
5	ACT	A	860	-	-	-	X
5	ACT	B	860	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7090 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NITRIC OXIDE SYNTHASE, BRAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	407	Total	C	N	O	S	0	1	0
			3319	2125	566	607	21			
1	B	411	Total	C	N	O	S	0	2	0
			3351	2144	574	611	22			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	9	5	3		
3	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is 7-{2-[(3-FLUOROBENZYL)AMINO]ETHYL}QUINOLIN-2-AMINE (three-letter code: M7K) (formula: C₁₈H₁₈FN₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	F	N	0	0
			22	18	1	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	F	N	0	0
			22	18	1	3		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Zn	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	105	Total	O	0	0
			105	105		
7	B	142	Total	O	0	0
			142	142		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NITRIC OXIDE SYNTHASE, BRAIN



• Molecule 1: NITRIC OXIDE SYNTHASE, BRAIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.92Å 111.35Å 164.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.12 – 1.91 39.09 – 1.91	Depositor EDS
% Data completeness (in resolution range)	99.3 (39.12-1.91) 99.4 (39.09-1.91)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.77 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.196 , 0.239 0.195 , 0.238	Depositor DCC
R_{free} test set	3706 reflections (5.23%)	DCC
Wilson B-factor (Å ²)	36.3	Xtriage
Anisotropy	0.564	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 48.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	3 of 74865 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7090	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: M7K, HEM, ZN, H4B, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	0/3415	0.70	1/4633 (0.0%)
1	B	0.71	0/3450	0.80	2/4677 (0.0%)
All	All	0.65	0/6865	0.75	3/9310 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	410	ARG	NE-CZ-NH1	6.13	123.37	120.30
1	B	489	ASP	CB-CG-OD1	5.30	123.07	118.30
1	B	430	ARG	NE-CZ-NH1	5.05	122.83	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3319	0	3227	35	0
1	B	3351	0	3269	30	0
2	A	43	0	30	2	0
2	B	43	0	30	7	0
3	A	17	0	15	0	0
3	B	17	0	15	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	22	0	18	5	0
4	B	22	0	18	4	0
5	A	4	0	3	0	0
5	B	4	0	3	0	0
6	A	1	0	0	0	0
7	A	105	0	0	2	0
7	B	142	0	0	1	0
All	All	7090	0	6628	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:486:LYS:HE2	1:A:499:VAL:HG11	1.62	0.81
2:B:750:HEM:HH C	2:B:750:HEM:HBB2	1.65	0.78
1:A:350:THR:N	1:A:353:GLN:HE21	1.89	0.71
1:A:487:GLN:HB2	1:A:489:ASP:HB3	1.73	0.70
1:A:523:LEU:HD22	1:A:531:PRO:HB2	1.74	0.68
1:A:328:GLU:HB3	1:B:323:GLU:HG2	1.78	0.65
1:B:504:ILE:O	1:B:508:GLN:HG2	1.96	0.65
1:B:364:GLN:NE2	7:B:2012:HOH:O	2.31	0.61
1:A:462:PHE:HB2	1:A:581:ALA:HB3	1.83	0.59
2:B:750:HEM:HBA2	4:B:800:M7K:C09	2.32	0.59
1:A:350:THR:N	1:A:353:GLN:NE2	2.51	0.59
1:A:620:LYS:HB2	1:A:620:LYS:NZ	2.19	0.57
1:B:475:TRP:HB2	1:B:523:LEU:HB3	1.86	0.57
1:B:322:LEU:HD13	1:B:699:ARG:HH21	1.70	0.56
1:A:455:LEU:HD12	1:A:587:TRP:HB3	1.87	0.56
1:B:355:PHE:CE1	1:B:385:ASN:HB2	2.42	0.55
1:B:584:PHE:HE1	4:B:800:M7K:H06	1.72	0.54
1:B:607:LEU:HD13	1:B:626:LYS:HG2	1.89	0.54
2:B:750:HEM:HBA2	4:B:800:M7K:H09	1.90	0.54
1:B:328:GLU:H	1:B:328:GLU:CD	2.10	0.54
1:A:380:ARG:HD3	1:A:400:GLU:OE2	2.08	0.53
1:A:620:LYS:HB2	1:A:620:LYS:HZ2	1.74	0.52
1:B:567:VAL:HG21	4:B:800:M7K:C07	2.40	0.51
1:B:350:THR:HB	1:B:352:ASP:CB	2.41	0.51
1:B:572:LEU:HB3	1:B:579:PHE:HB2	1.94	0.50
1:B:462:PHE:HB2	1:B:581:ALA:HB3	1.94	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:524:LEU:O	1:A:531:PRO:HA	2.12	0.48
1:B:350:THR:HB	1:B:352:ASP:HB2	1.95	0.47
1:B:415:CYS:HB2	2:B:750:HEM:ND	2.29	0.47
1:A:330:ILE:HD11	1:B:696:LEU:HB3	1.95	0.47
1:B:352:ASP:HB3	1:B:353:GLN:HE21	1.80	0.47
1:A:465:ARG:HD2	1:A:578:GLU:OE1	2.15	0.47
1:A:371:ARG:HG3	1:A:371:ARG:HH11	1.80	0.47
1:A:686:SER:HA	1:A:691:PHE:CG	2.50	0.47
1:A:567:VAL:HG21	4:A:800:M7K:C07	2.45	0.46
1:B:584:PHE:CD1	2:B:750:HEM:HAC	2.51	0.46
1:B:478:GLN:HB2	1:B:481:ARG:HG3	1.97	0.46
1:A:387:GLU:OE2	1:A:394:TYR:HA	2.16	0.46
1:A:306:TRP:CD2	1:B:336:MET:HE3	2.51	0.46
1:B:409:TRP:CE3	1:B:421:TRP:HA	2.51	0.45
1:A:588:TYR:CD2	1:A:593:ILE:HD11	2.52	0.45
1:A:569:ASN:H	1:A:569:ASN:HD22	1.63	0.45
1:A:595:VAL:O	1:A:599:CYS:HB2	2.17	0.45
2:A:750:HEM:HBD1	4:A:800:M7K:H11	2.00	0.44
1:B:584:PHE:CD1	2:B:750:HEM:CAC	3.01	0.44
1:A:455:LEU:HD12	1:A:587:TRP:CB	2.48	0.44
1:A:415:CYS:HB3	1:A:418:ARG:HG3	1.99	0.44
2:A:750:HEM:HBA2	4:A:800:M7K:C09	2.48	0.43
1:B:675:ASP:O	1:B:679:ILE:HG12	2.18	0.43
1:A:605:ASN:ND2	7:A:2077:HOH:O	2.50	0.43
1:A:439:PHE:CZ	1:A:443:CYS:SG	3.12	0.43
1:B:302:LYS:HA	1:B:312:LEU:O	2.19	0.43
1:A:626:LYS:HB3	1:B:687:ILE:HD12	2.00	0.43
1:B:403:TYR:CE1	1:B:407:HIS:CE1	3.07	0.42
1:B:445:HIS:C	1:B:445:HIS:CD2	2.93	0.42
1:A:554:PHE:HB3	7:A:2063:HOH:O	2.20	0.42
1:B:388:ILE:O	1:B:392:SER:N	2.50	0.41
2:B:750:HEM:HHC	2:B:750:HEM:CBB	2.44	0.41
1:B:352:ASP:HB3	1:B:353:GLN:NE2	2.36	0.41
1:A:488:PRO:O	1:A:490:GLY:N	2.53	0.41
1:A:592:GLU:OE2	4:A:800:M7K:N01	2.53	0.41
1:A:569:ASN:H	1:A:569:ASN:ND2	2.19	0.41
1:A:478:GLN:HA	1:A:566:ALA:O	2.21	0.41
1:A:584:PHE:HE1	4:A:800:M7K:H06	1.86	0.40
1:A:473:ARG:NH2	1:A:710:PRO:HD3	2.37	0.40
1:B:362:LEU:HD12	1:B:381:LEU:HD23	2.03	0.40
1:A:676:TRP:CZ2	1:A:680:VAL:HG21	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	404/422 (96%)	386 (96%)	17 (4%)	1 (0%)	52	42
1	B	409/422 (97%)	395 (97%)	12 (3%)	2 (0%)	34	20
All	All	813/844 (96%)	781 (96%)	29 (4%)	3 (0%)	39	27

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	489	ASP
1	B	352	ASP
1	B	506	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	364/377 (97%)	347 (95%)	17 (5%)	32	19
1	B	368/377 (98%)	356 (97%)	12 (3%)	45	33
All	All	732/754 (97%)	703 (96%)	29 (4%)	38	25

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	319	LYS
1	A	337	LEU
1	A	350	THR
1	A	352	ASP
1	A	360	GLU
1	A	370	LYS
1	A	380	ARG
1	A	390	SER
1	A	469	LYS
1	A	507	GLN
1	A	523	LEU
1	A	547	ARG
1	A	555	LYS
1	A	569	ASN
1	A	645	LYS
1	A	663	GLU
1	A	715	VAL
1	B	332	MET
1	B	337	LEU
1	B	350	THR
1	B	353	GLN
1	B	390	SER
1	B	392	SER
1	B	481	ARG
1	B	486	LYS
1	B	503	GLU
1	B	511	LYS
1	B	547	ARG
1	B	620	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	353	GLN
1	A	454	ASN
1	A	507	GLN
1	A	527	ASN
1	A	569	ASN
1	A	628	GLN
1	A	642	GLN
1	A	697	ASN
1	B	353	GLN
1	B	364	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	385	ASN
1	B	425	GLN
1	B	454	ASN
1	B	507	GLN
1	B	535	GLN
1	B	601	ASN
1	B	605	ASN
1	B	642	GLN
1	B	697	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	HEM	A	750	1	30,50,50	2.03	7 (23%)	24,82,82	2.56	11 (45%)
3	H4B	A	760	-	13,18,18	1.29	2 (15%)	11,26,26	2.71	6 (54%)
4	M7K	A	800	-	24,24,24	1.20	2 (8%)	31,32,32	1.50	4 (12%)
5	ACT	A	860	-	1,3,3	1.54	0	0,3,3	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	B	750	1	30,50,50	2.21	6 (20%)	24,82,82	2.79	12 (50%)
3	H4B	B	760	-	13,18,18	1.10	1 (7%)	11,26,26	2.45	6 (54%)
4	M7K	B	800	-	24,24,24	1.09	1 (4%)	31,32,32	1.57	5 (16%)
5	ACT	B	860	-	1,3,3	1.30	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	750	1	-	0/10/54/54	0/0/8/8
3	H4B	A	760	-	-	0/8/17/17	0/2/2/2
4	M7K	A	800	-	-	0/7/7/7	0/3/3/3
5	ACT	A	860	-	-	0/0/0/0	0/0/0/0
2	HEM	B	750	1	-	0/10/54/54	0/0/8/8
3	H4B	B	760	-	-	0/8/17/17	0/2/2/2
4	M7K	B	800	-	-	0/7/7/7	0/3/3/3
5	ACT	B	860	-	-	0/0/0/0	0/0/0/0

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	750	HEM	C2C-C1C	-6.78	1.39	1.52
2	A	750	HEM	C2D-C3D	-6.15	1.36	1.54
2	B	750	HEM	C2D-C3D	-6.00	1.36	1.54
2	A	750	HEM	C2C-C1C	-5.94	1.41	1.52
2	B	750	HEM	C3D-C4D	-4.07	1.46	1.51
2	A	750	HEM	C3D-C4D	-2.96	1.47	1.51
2	B	750	HEM	C2B-C1B	-2.88	1.42	1.51
2	A	750	HEM	C2B-C1B	-2.12	1.44	1.51
2	B	750	HEM	C1C-NC	2.05	1.38	1.36
4	A	800	M7K	C04-C03	2.19	1.41	1.36
2	A	750	HEM	CHC-C1C	2.31	1.41	1.36
3	B	760	H4B	C7-N8	2.35	1.49	1.46
3	A	760	H4B	C4-N3	2.47	1.37	1.33
4	B	800	M7K	C02-N01	2.54	1.36	1.33
2	A	750	HEM	FE-NB	2.58	2.11	1.97
3	A	760	H4B	C2-N2	2.78	1.39	1.34
2	B	750	HEM	FE-NC	3.19	2.08	1.95
2	A	750	HEM	FE-NC	3.31	2.08	1.95
4	A	800	M7K	C02-N01	3.49	1.37	1.33

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	750	HEM	CBA-CAA-C2A	-5.90	101.96	112.53
2	A	750	HEM	CBA-CAA-C2A	-5.00	103.56	112.53
2	B	750	HEM	C1D-CHD-C4C	-4.36	118.53	125.82
3	A	760	H4B	N3-C2-N1	-4.06	118.87	125.53
2	A	750	HEM	C1D-CHD-C4C	-3.68	119.67	125.82
4	B	800	M7K	C03-C02-N01	-3.67	118.19	122.20
3	B	760	H4B	N3-C2-N1	-3.02	120.59	125.53
4	A	800	M7K	C03-C02-N01	-2.79	119.15	122.20
2	A	750	HEM	C3C-CAC-CBC	-2.77	120.21	124.46
4	B	800	M7K	C24-C23-C22	-2.68	119.89	123.35
2	B	750	HEM	C3B-C4B-NB	-2.62	106.63	111.63
2	B	750	HEM	CMA-C3A-C4A	-2.59	124.08	128.36
4	A	800	M7K	C24-C23-C22	-2.40	120.25	123.35
2	B	750	HEM	CAA-C2A-C1A	-2.07	124.76	127.01
2	A	750	HEM	CMA-C3A-C4A	-2.02	125.03	128.36
2	A	750	HEM	CAA-CBA-CGA	2.01	116.43	112.75
2	B	750	HEM	CHC-C4B-NB	2.26	129.97	124.52
3	B	760	H4B	N2-C2-N1	2.27	120.97	117.20
2	B	750	HEM	C2D-C3D-C4D	2.32	105.44	101.50
3	A	760	H4B	N2-C2-N1	2.33	121.06	117.20
3	B	760	H4B	C4A-C8A-N8	2.47	121.34	118.43
2	A	750	HEM	C2D-C3D-C4D	2.50	105.73	101.50
4	B	800	M7K	C14-N13-C12	2.52	122.75	113.48
3	B	760	H4B	C2-N1-C8A	2.58	120.33	114.54
2	A	750	HEM	CMD-C2D-C3D	2.81	126.78	114.35
3	A	760	H4B	C4A-C8A-N8	2.82	121.75	118.43
4	A	800	M7K	C02-N01-C10	3.09	121.47	117.78
2	B	750	HEM	CMD-C2D-C3D	3.20	128.49	114.35
3	A	760	H4B	C2-N1-C8A	3.48	122.36	114.54
2	A	750	HEM	CMB-C2B-C3B	3.59	125.48	116.53
2	A	750	HEM	CAD-C3D-C4D	3.81	125.91	112.47
2	B	750	HEM	CAD-C3D-C4D	3.85	126.04	112.47
4	B	800	M7K	C02-N01-C10	3.87	122.39	117.78
3	B	760	H4B	C4-N3-C2	3.94	121.40	115.94
2	B	750	HEM	CMB-C2B-C3B	4.03	126.59	116.53
3	A	760	H4B	C4-N3-C2	4.07	121.59	115.94
4	B	800	M7K	N02-C02-N01	4.10	120.70	118.03
2	B	750	HEM	CMC-C2C-C3C	4.20	127.02	116.53
4	A	800	M7K	N02-C02-N01	4.24	120.79	118.03
3	A	760	H4B	C4-C4A-C8A	4.24	118.40	114.56
2	A	750	HEM	CMC-C2C-C3C	4.41	127.55	116.53
3	B	760	H4B	C4-C4A-C8A	4.78	118.89	114.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	750	HEM	CAD-C3D-C2D	5.27	128.36	113.22
2	B	750	HEM	CAD-C3D-C2D	5.32	128.52	113.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	750	HEM	2	0
4	A	800	M7K	5	0
2	B	750	HEM	7	0
4	B	800	M7K	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	407/422 (96%)	0.99	78 (19%) 2 2	28, 56, 98, 131	0
1	B	411/422 (97%)	0.40	30 (7%) 18 20	27, 43, 74, 98	0
All	All	818/844 (96%)	0.69	108 (13%) 4 5	27, 48, 91, 131	0

All (108) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	300	PHE	8.3
1	A	716	TRP	8.3
1	A	715	VAL	7.5
1	A	352	ASP	5.7
1	A	488	PRO	5.6
1	A	713	THR	5.5
1	B	348	VAL	5.5
1	A	355	PHE	5.4
1	A	551	PHE	5.3
1	B	619	ARG	4.8
1	B	718	GLY	4.8
1	A	491	SER	4.7
1	A	490	GLY	4.6
1	A	486	LYS	4.5
1	A	351	LYS	4.5
1	B	338	PRO	4.3
1	A	712	ASN	4.3
1	A	506	ILE	4.3
1	B	677	VAL	4.2
1	B	350	THR	4.1
1	A	489	ASP	4.1
1	A	470	HIS	4.1
1	A	487	GLN	4.0
1	A	714	HIS	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	507	GLN	3.9
1	A	391	THR	3.9
1	A	338	PRO	3.8
1	A	504	ILE	3.7
1	A	390	SER	3.7
1	A	711	TRP	3.7
1	A	392	SER	3.7
1	A	386	LYS	3.6
1	A	503	GLU	3.6
1	A	679	ILE	3.6
1	A	677	VAL	3.6
1	A	553	TRP	3.5
1	B	680	VAL	3.5
1	A	300	PHE	3.5
1	B	299	ARG	3.5
1	A	480	ILE	3.4
1	A	388	ILE	3.4
1	A	393	THR	3.3
1	A	415	CYS	3.2
1	A	710	PRO	3.2
1	A	619	ARG	3.2
1	A	385	ASN	3.2
1	A	567	VAL	3.1
1	A	588	TYR	3.1
1	A	680	VAL	3.1
1	A	676	TRP	3.1
1	A	682	PRO	3.1
1	A	508	GLN	3.0
1	A	389	GLU	3.0
1	B	352	ASP	3.0
1	B	561	TRP	3.0
1	B	715	VAL	3.0
1	B	321	THR	3.0
1	A	554	PHE	2.9
1	A	382	GLU	2.9
1	A	350	THR	2.9
1	A	354	LEU	2.9
1	B	616	LEU	2.9
1	B	302	LYS	2.9
1	A	681	PRO	2.9
1	A	561	TRP	2.9
1	A	593	ILE	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	678	TRP	2.8
1	A	514	ARG	2.7
1	B	676	TRP	2.7
1	B	682	PRO	2.7
1	A	416	VAL	2.6
1	B	353	GLN	2.6
1	B	591	THR	2.6
1	A	499	VAL	2.6
1	B	679	ILE	2.6
1	B	667	ARG	2.6
1	A	322	LEU	2.6
1	A	469	LYS	2.6
1	A	591	THR	2.6
1	A	511	LYS	2.5
1	A	493	LEU	2.5
1	A	617	ASP	2.5
1	A	479	LEU	2.5
1	A	584	PHE	2.5
1	B	593	ILE	2.4
1	A	630	LEU	2.4
1	A	321	THR	2.4
1	B	691	PHE	2.4
1	A	531	PRO	2.3
1	B	620	LYS	2.3
1	B	567	VAL	2.3
1	A	565	PRO	2.3
1	B	681	PRO	2.3
1	A	566	ALA	2.3
1	A	685	GLY	2.3
1	B	588	TYR	2.2
1	A	564	LEU	2.2
1	A	384	VAL	2.2
1	A	691	PHE	2.2
1	A	492	THR	2.1
1	A	667	ARG	2.1
1	B	678	TRP	2.1
1	B	615	ASP	2.1
1	A	590	GLY	2.0
1	A	618	MET	2.0
1	B	608	GLU	2.0
1	A	452	LYS	2.0
1	A	467	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	ACT	B	860	4/4	0.94	0.24	7.05	66,69,69,70	0
5	ACT	A	860	4/4	0.95	0.25	6.91	75,76,78,80	0
4	M7K	B	800	22/22	0.94	0.23	2.23	29,47,101,102	0
2	HEM	B	750	43/43	0.97	0.17	0.72	27,30,43,50	0
3	H4B	B	760	17/17	0.95	0.19	0.60	35,38,41,42	0
2	HEM	A	750	43/43	0.98	0.19	0.35	32,34,43,45	0
3	H4B	A	760	17/17	0.94	0.16	0.15	35,39,42,42	0
4	M7K	A	800	22/22	0.93	0.19	0.02	25,42,82,82	0
6	ZN	A	1717	1/1	1.00	0.10	-0.34	43,43,43,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.