



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:28 PM GMT

PDB ID : 4CC1
Title : Notch ligand, Jagged-1, contains an N-terminal C2 domain
Authors : Chilakuri, C.R.; Sheppard, D.; Ilagan, M.X.G.; Holt, L.R.; Abbott, F.; Liang, S.; Kopan, R.; Handford, P.A.; Lea, S.M.
Deposited on : 2013-10-17
Resolution : 2.84 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	CL	A	1341	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 4798 atoms, of which 12 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN JAGGED-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	306	Total	C	N	O	S	0	0	0
			2363	1448	424	457	34			
1	B	306	Total	C	N	O	S	0	0	0
			2363	1448	424	457	34			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	336	VAL	-	EXPRESSION TAG	UNP P78504
A	337	ASP	-	EXPRESSION TAG	UNP P78504
A	338	HIS	-	EXPRESSION TAG	UNP P78504
A	339	HIS	-	EXPRESSION TAG	UNP P78504
A	340	HIS	-	EXPRESSION TAG	UNP P78504
A	341	HIS	-	EXPRESSION TAG	UNP P78504
A	342	HIS	-	EXPRESSION TAG	UNP P78504
A	343	HIS	-	EXPRESSION TAG	UNP P78504
B	336	VAL	-	EXPRESSION TAG	UNP P78504
B	337	ASP	-	EXPRESSION TAG	UNP P78504
B	338	HIS	-	EXPRESSION TAG	UNP P78504
B	339	HIS	-	EXPRESSION TAG	UNP P78504
B	340	HIS	-	EXPRESSION TAG	UNP P78504
B	341	HIS	-	EXPRESSION TAG	UNP P78504
B	342	HIS	-	EXPRESSION TAG	UNP P78504
B	343	HIS	-	EXPRESSION TAG	UNP P78504

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is SUGAR (ALPHA-L-FUCOSE) (three-letter code: FUC) (formula: $C_6H_{12}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	6	4		


- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Ca 1 1	0	0
4	A	1	Total Ca 1 1	0	0

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	2	Total Cl 2 2	0	0
5	A	4	Total Cl 4 4	0	0

- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total C H O 10 2 6 2	0	0
6	B	1	Total C H O 10 2 6 2	0	0

- Molecule 7 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total O 1 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	5	Total	O	0	0
			5	5		

- Molecule 1: PROTEIN JAGGED-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	55.95Å 60.56Å 63.03Å 92.89° 104.95° 105.92°	Depositor
Resolution (Å)	60.38 – 2.84 60.38 – 2.84	Depositor EDS
% Data completeness (in resolution range)	98.2 (60.38-2.84) 94.0 (60.38-2.84)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.39 (at 2.86Å)	Xtriage
Refinement program	BUSTER 2.11.4	Depositor
R, R_{free}	0.196 , 0.249 0.210 , 0.268	Depositor DCC
R_{free} test set	988 reflections (5.92%)	DCC
Wilson B-factor (Å ²)	60.7	Xtriage
Anisotropy	0.217	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 51.7	EDS
Estimated twinning fraction	0.016 for -h,-l,-k	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 17666 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4798	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, CA, EDO, NAG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/2425	0.75	0/3283
1	B	0.51	0/2425	0.75	0/3283
All	All	0.51	0/4850	0.75	0/6566

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2363	0	2145	10	0
1	B	2363	0	2146	14	0
2	A	14	0	13	0	0
2	B	14	0	13	0	0
3	A	10	0	10	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	4	0	0	0	0
5	B	2	0	0	1	0
6	B	8	12	12	4	0
7	A	1	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	5	0	0	0	0
All	All	4786	12	4339	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:59:ARG:H	6:B:1342:EDO:H21	1.58	0.69
1:B:180:GLU:HG3	6:B:1341:EDO:H22	1.81	0.63
1:B:260:LEU:HD23	1:B:261:TYR:CZ	2.43	0.53
1:A:260:LEU:HD23	1:A:261:TYR:CZ	2.44	0.52
1:A:272:VAL:O	1:A:293:CYS:HB2	2.12	0.50
1:A:199:PHE:CE1	1:A:201:ARG:HB2	2.47	0.50
1:B:180:GLU:HG3	6:B:1341:EDO:C2	2.43	0.48
1:A:171:LYS:HG2	1:A:180:GLU:HG2	1.96	0.48
1:B:272:VAL:O	1:B:293:CYS:HB2	2.13	0.48
1:B:68:ARG:HB2	1:B:69:ASP:H	1.55	0.47
1:B:210:TYR:CZ	1:B:218:LYS:HE3	2.50	0.47
1:A:32:SER:HB3	7:A:2001:HOH:O	2.15	0.47
1:A:163:PRO:HA	1:A:185:VAL:O	2.15	0.47
1:B:178:HIS:HB3	6:B:1341:EDO:H12	1.95	0.46
1:B:163:PRO:HA	1:B:185:VAL:O	2.15	0.46
1:B:230:ASN:HB2	5:B:1340:CL:CL	2.52	0.46
1:B:171:LYS:HG2	1:B:180:GLU:HG2	1.97	0.45
1:A:82:TYR:CE1	1:A:158:SER:HB3	2.52	0.44
1:B:190:TYR:HB2	1:B:199:PHE:CE1	2.54	0.42
1:A:131:SER:HA	1:A:159:GLY:O	2.20	0.42
1:B:153:GLU:HB2	1:B:177:ALA:HB3	2.03	0.41
1:A:213:ASP:OD2	1:A:217:ASN:HB2	2.21	0.41
1:A:218:LYS:HE2	1:A:228:GLU:OE1	2.21	0.41
1:B:153:GLU:HB3	1:B:179:PHE:HE2	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	304/312 (97%)	282 (93%)	22 (7%)	0	100	100
1	B	304/312 (97%)	277 (91%)	27 (9%)	0	100	100
All	All	608/624 (97%)	559 (92%)	49 (8%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/267 (98%)	248 (95%)	13 (5%)	30	62
1	B	261/267 (98%)	247 (95%)	14 (5%)	27	59
All	All	522/534 (98%)	495 (95%)	27 (5%)	29	61

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	69	ASP
1	A	84	SER
1	A	98	SER
1	A	109	LEU
1	A	144	ASP
1	A	146	VAL
1	A	165	ARG
1	A	197	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	201	ARG
1	A	230	ASN
1	A	236	GLN
1	A	263	ASP
1	A	323	SER
1	B	68	ARG
1	B	86	VAL
1	B	98	SER
1	B	109	LEU
1	B	142	SER
1	B	147	GLN
1	B	160	MET
1	B	174	THR
1	B	197	ASN
1	B	201	ARG
1	B	203	ARG
1	B	247	LEU
1	B	263	ASP
1	B	323	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 8 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	1338	1	14,14,15	0.30	0	15,19,21	0.41	0
3	FUC	A	1339	1	10,10,11	0.45	0	14,14,16	1.36	1 (7%)
2	NAG	B	1338	1	14,14,15	0.41	0	15,19,21	0.63	0
6	EDO	B	1341	-	3,3,3	1.03	0	2,2,2	0.40	0
6	EDO	B	1342	-	3,3,3	1.17	0	2,2,2	0.16	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1338	1	-	0/6/23/26	0/1/1/1
3	FUC	A	1339	1	-	0/0/17/20	0/1/1/1
2	NAG	B	1338	1	-	0/6/23/26	0/1/1/1
6	EDO	B	1341	-	-	0/1/1/1	0/0/0/0
6	EDO	B	1342	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1339	FUC	C1-O5-C5	4.67	119.59	112.38

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	1341	EDO	3	0
6	B	1342	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	306/312 (98%)	-0.12	2 (0%) 89 85	37, 63, 93, 111	0
1	B	306/312 (98%)	0.02	9 (2%) 55 45	33, 65, 131, 157	0
All	All	612/624 (98%)	-0.05	11 (1%) 71 63	33, 64, 110, 157	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	144	ASP	4.9
1	B	88	ALA	3.8
1	A	56	GLY	3.4
1	B	326	GLU	2.9
1	B	335	ILE	2.7
1	B	308	ASN	2.5
1	B	323	SER	2.4
1	A	68	ARG	2.1
1	B	292	LEU	2.1
1	B	309	GLY	2.1
1	B	85	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	CL	A	1341	1/1	0.98	0.34	4.18	17,17,17,17	0
5	CL	B	1340	1/1	0.96	0.26	1.71	55,55,55,55	0
6	EDO	B	1341	4/4	0.93	0.23	1.13	36,50,60,60	0
5	CL	A	1342	1/1	0.96	0.18	-0.36	42,42,42,42	0
4	CA	A	1340	1/1	0.93	0.11	-0.75	82,82,82,82	0
6	EDO	B	1342	4/4	0.84	0.12	-0.77	44,48,58,58	0
3	FUC	A	1339	10/11	0.97	0.12	-1.42	66,70,72,73	0
4	CA	B	1339	1/1	0.97	0.10	-1.88	85,85,85,85	0
5	CL	A	1344	1/1	0.95	0.17	-	49,49,49,49	0
5	CL	B	1343	1/1	0.89	0.11	-	51,51,51,51	0
2	NAG	A	1338	14/15	0.86	0.16	-	94,97,101,102	0
2	NAG	B	1338	14/15	0.81	0.20	-	105,107,108,109	0
5	CL	A	1343	1/1	0.96	0.20	-	51,51,51,51	0

6.5 Other polymers

There are no such residues in this entry.