



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 03:30 PM GMT

PDB ID : 4CCF
Title : Structure of Respiratory Syncytial Virus F protein head domain
Authors : Peat, T.S.
Deposited on : 2013-10-22
Resolution : 2.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION






The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3152 (2.70-2.62)
Clashscore	102246	3524 (2.70-2.62)
Ramachandran outliers	100387	3469 (2.70-2.62)
Sidechain outliers	100360	3469 (2.70-2.62)
RSRZ outliers	91569	3161 (2.70-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	574	
1	B	574	
1	C	574	
1	D	574	
1	E	574	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	574	 A horizontal bar chart showing the quality of the chain. The bar is divided into three segments: a green segment representing 41%, a yellow segment representing 5%, and a grey segment representing 54%. The percentages are labeled below the bar.

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 12234 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FUSION GLYCOPROTEIN F0.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	265	Total	C	N	O	S	0	0	0
			2036	1278	338	402	18			
1	B	258	Total	C	N	O	S	0	0	0
			1995	1251	331	395	18			
1	C	261	Total	C	N	O	S	0	1	0
			2026	1272	335	401	18			
1	D	268	Total	C	N	O	S	0	1	0
			2080	1309	343	410	18			
1	E	257	Total	C	N	O	S	0	2	0
			2005	1258	333	396	18			
1	F	265	Total	C	N	O	S	0	1	0
			2041	1281	338	404	18			

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	1	Total O S 5 4 1	0	0
2	D	1	Total O S 5 4 1	0	0
2	E	1	Total O S 5 4 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	8	Total O 8 8	0	0
3	B	6	Total O 6 6	0	0
3	C	6	Total O 6 6	0	0
3	D	3	Total O 3 3	0	0
3	E	8	Total O 8 8	0	0
3	F	5	Total O 5 5	0	0


3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: FUSION GLYCOPROTEIN F0




- Molecule 1: FUSION GLYCOPROTEIN F0







• Molecule 1: FUSION GLYCOPROTEIN F0



• Molecule 1: FUSION GLYCOPROTEIN F0





PHE
SER
ASN

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	68.38Å 133.50Å 116.42Å 90.00° 93.32° 90.00°	Depositor
Resolution (Å)	43.07 – 2.65 43.04 – 2.65	Depositor EDS
% Data completeness (in resolution range)	87.7 (43.07-2.65) 87.8 (43.04-2.65)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.65 (at 2.65Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.215 , 0.247 0.215 , 0.247	Depositor DCC
R_{free} test set	2763 reflections (5.48%)	DCC
Wilson B-factor (Å ²)	40.7	Xtriage
Anisotropy	0.137	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 14.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtriage
Outliers	0 of 53167 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	12234	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/2068	0.76	2/2811 (0.1%)
1	B	0.60	0/2026	0.74	0/2751
1	C	0.58	0/2057	0.75	1/2793 (0.0%)
1	D	0.54	0/2112	0.75	1/2868 (0.0%)
1	E	0.59	0/2036	0.75	1/2764 (0.0%)
1	F	0.57	0/2073	0.75	1/2817 (0.0%)
All	All	0.58	0/12372	0.75	6/16804 (0.0%)

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	448	ASP	CB-CG-OD1	9.11	126.50	118.30
1	F	264	MET	CG-SD-CE	7.83	112.73	100.20
1	C	392	ASP	CB-CG-OD1	5.85	123.56	118.30
1	A	364	ARG	CB-CG-CD	-5.80	96.52	111.60
1	A	392	ASP	CB-CG-OD1	5.66	123.39	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2036	0	2005	22	1
1	B	1995	0	1972	17	0
1	C	2026	0	2003	15	0
1	D	2080	0	2061	33	0
1	E	2005	0	1986	23	1
1	F	2041	0	2005	29	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
2	E	5	0	0	0	0
3	A	8	0	0	0	0
3	B	6	0	0	0	0
3	C	6	0	0	1	0
3	D	3	0	0	0	0
3	E	8	0	0	0	0
3	F	5	0	0	0	0
All	All	12234	0	12032	126	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 126 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:334:LEU:HB3	1:B:395:ILE:HD11	1.07	1.03
1:A:385:ASP:O	1:A:388:ASN:ND2	1.93	1.01
1:A:311:THR:HG21	1:A:344:ASP:O	1.65	0.96
1:E:311:THR:HG21	1:E:344:ASP:O	1.68	0.94
1:B:270:GLN:HG2	1:B:309:ILE:HD12	1.53	0.90

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:440:ASP:OD2	1:E:42:LYS:NZ[1_656]	2.10	0.10

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	259/574 (45%)	248 (96%)	11 (4%)	0	100	100
1	B	250/574 (44%)	242 (97%)	7 (3%)	1 (0%)	39	65
1	C	254/574 (44%)	247 (97%)	7 (3%)	0	100	100
1	D	261/574 (46%)	250 (96%)	10 (4%)	1 (0%)	39	65
1	E	251/574 (44%)	245 (98%)	5 (2%)	1 (0%)	39	65
1	F	260/574 (45%)	248 (95%)	12 (5%)	0	100	100
All	All	1535/3444 (45%)	1480 (96%)	52 (3%)	3 (0%)	52	77

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	437	ASN
1	B	437	ASN
1	E	437	ASN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/520 (46%)	238 (100%)	1 (0%)	93	98
1	B	237/520 (46%)	234 (99%)	3 (1%)	76	91
1	C	240/520 (46%)	240 (100%)	0	100	100
1	D	246/520 (47%)	245 (100%)	1 (0%)	93	98

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	238/520 (46%)	236 (99%)	2 (1%)	86	95
1	F	239/520 (46%)	238 (100%)	1 (0%)	93	98
All	All	1439/3120 (46%)	1431 (99%)	8 (1%)	92	97

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	454	ASN
1	F	291	ILE
1	E	388[A]	ASN
1	B	440	ASP
1	D	388	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	454	ASN
1	F	426	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	C	1466	-	4,4,4	0.21	0	6,6,6	1.28	1 (16%)
2	SO4	D	1470	-	4,4,4	0.63	0	6,6,6	0.39	0
2	SO4	E	1466	-	4,4,4	0.47	0	6,6,6	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	C	1466	-	-	0/0/0/0	0/0/0/0
2	SO4	D	1470	-	-	0/0/0/0	0/0/0/0
2	SO4	E	1466	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	C	1466	SO4	O2-S-O1	2.22	116.52	109.50

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	265/574 (46%)	-0.45	0	100 100	18, 38, 65, 107	1 (0%)
1	B	258/574 (44%)	-0.48	0	100 100	17, 37, 70, 91	0
1	C	261/574 (45%)	-0.52	0	100 100	16, 34, 64, 92	0
1	D	268/574 (46%)	-0.47	1 (0%)	93 94	21, 40, 66, 85	0
1	E	257/574 (44%)	-0.49	2 (0%)	87 87	17, 37, 65, 107	0
1	F	265/574 (46%)	-0.52	0	100 100	17, 36, 69, 97	0
All	All	1574/3444 (45%)	-0.49	3 (0%)	95 96	16, 37, 68, 107	1 (0%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	428	ASN	2.7
1	D	330	SER	2.4
1	E	429	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	SO4	C	1466	5/5	1.00	0.13	-0.01	26,26,27,28	0
2	SO4	E	1466	5/5	0.96	0.09	-2.51	53,56,58,64	0
2	SO4	D	1470	5/5	0.98	0.07	-4.59	57,59,64,69	0

6.5 Other polymers [i](#)

There are no such residues in this entry.