



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 01:18 AM GMT

PDB ID : 2CDH
Title : ARCHITECTURE OF THE THERMOMYCES LANUGINOSUS FUNGAL
FATTY ACID SYNTHASE AT 5 ANGSTROM RESOLUTION.
Authors : Jenni, S.; Leibundgut, M.; Maier, T.; Ban, N.
Deposited on : 2006-01-24
Resolution : 4.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.20 Å.





Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1140 (4.80-3.60)


























The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	0	226	 100%
1	1	226	 100%
1	2	226	 100%
1	3	226	 100%
1	Y	226	 100%
1	Z	226	 100%
2	4	305	 99% .
2	5	305	 99% .
2	6	305	 99% .



Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	7	305	 99% .
2	8	305	 99% .
2	9	305	 99% .
2	M	305	 99% .
2	N	305	 99% .
2	O	305	 99% .
2	P	305	 99% .
2	Q	305	 99% .
2	R	305	 99% .
3	A	406	 100%
3	B	406	 100%
3	C	406	 98% .
3	D	406	 98% .
3	E	406	 100%
3	F	406	 99% .
4	G	244	 100%
4	H	244	 100%
4	I	244	 100%
4	J	244	 100%
4	K	244	 100%
4	L	244	 100%
5	S	248	 100%
5	T	248	 100%
5	U	248	 100%
5	V	248	 100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	W	248	 100%
5	X	248	 100%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10404 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ENOYL REDUCTASE.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
1	0	226	Total C 226 226	0	0	226
1	1	226	Total C 226 226	0	0	226
1	2	226	Total C 226 226	0	0	226
1	3	226	Total C 226 226	0	0	226
1	Y	226	Total C 226 226	0	0	226
1	Z	226	Total C 226 226	0	0	226

- Molecule 2 is a protein called MALONYL/PALMITOYL TRANSFERASE.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
2	4	305	Total C 305 305	0	0	305
2	5	305	Total C 305 305	0	0	305
2	6	305	Total C 305 305	0	0	305
2	7	305	Total C 305 305	0	0	305
2	8	305	Total C 305 305	0	0	305
2	9	305	Total C 305 305	0	0	305
2	M	305	Total C 305 305	0	0	305
2	N	305	Total C 305 305	0	0	305

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
2	O	305	Total C 305 305	0	0	305
2	P	305	Total C 305 305	0	0	305
2	Q	305	Total C 305 305	0	0	305
2	R	305	Total C 305 305	0	0	305

- Molecule 3 is a protein called KETOACYL SYNTHASE.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
3	A	406	Total C 406 406	0	0	406
3	B	406	Total C 406 406	0	0	406
3	C	406	Total C 406 406	0	0	406
3	D	406	Total C 406 406	0	0	406
3	E	406	Total C 406 406	0	0	406
3	F	406	Total C 406 406	0	0	406

- Molecule 4 is a protein called KETOACYL REDUCTASE.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
4	G	244	Total C 244 244	0	0	244
4	H	244	Total C 244 244	0	0	244
4	I	244	Total C 244 244	0	0	244
4	J	244	Total C 244 244	0	0	244
4	K	244	Total C 244 244	0	0	244
4	L	244	Total C 244 244	0	0	244

- Molecule 5 is a protein called DEHYDRATASE.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
5	S	248	Total C 248 248	0	0	248
5	T	248	Total C 248 248	0	0	248
5	U	248	Total C 248 248	0	0	248
5	V	248	Total C 248 248	0	0	248
5	W	248	Total C 248 248	0	0	248
5	X	248	Total C 248 248	0	0	248

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: ENOYL REDUCTASE

Chain 0:  100%

There are no outlier residues recorded for this chain.

- Molecule 1: ENOYL REDUCTASE

Chain 1:  100%

There are no outlier residues recorded for this chain.

- Molecule 1: ENOYL REDUCTASE

Chain 2:  100%

There are no outlier residues recorded for this chain.

- Molecule 1: ENOYL REDUCTASE

Chain 3:  100%

There are no outlier residues recorded for this chain.

- Molecule 1: ENOYL REDUCTASE

Chain Y:  100%

There are no outlier residues recorded for this chain.


- Molecule 1: ENOYL REDUCTASE

Chain Z:  100%

There are no outlier residues recorded for this chain.


- Molecule 2: MALONYL/PALMITOYL TRANSFERASE

Chain 4:  99%




- Molecule 2: MALONYL/PALMITOYL TRANSFERASE

Chain 5:  99% .



- Molecule 2: MALONYL/PALMITOYL TRANSFERASE

Chain 6:  99% .



- Molecule 2: MALONYL/PALMITOYL TRANSFERASE

Chain 7:  99% .




- Molecule 2: MALONYL/PALMITOYL TRANSFERASE

Chain 8:  99% .



- Molecule 2: MALONYL/PALMITOYL TRANSFERASE

Chain 9:  99% .



- Molecule 2: MALONYL/PALMITOYL TRANSFERASE

Chain M:  99% .



- Molecule 2: MALONYL/PALMITOYL TRANSFERASE

Chain N:  99% .



- Molecule 2: MALONYL/PALMITOYL TRANSFERASE

Chain O:  99% .



- Molecule 2: MALONYL/PALMITOYL TRANSFERASE

Chain P:  99% .



- Molecule 2: MALONYL/PALMITOYL TRANSFERASE

Chain Q:  99% .



- Molecule 2: MALONYL/PALMITOYL TRANSFERASE

Chain R:  99% .



- Molecule 3: KETOACYL SYNTHASE

Chain A:  100% .

There are no outlier residues recorded for this chain.


- Molecule 3: KETOACYL SYNTHASE

Chain B:  100% .

There are no outlier residues recorded for this chain.

- Molecule 3: KETOACYL SYNTHASE

Chain C:  98% .



- Molecule 3: KETOACYL SYNTHASE

Chain D:  98% .




- Molecule 3: KETOACYL SYNTHASE

Chain E:  100%



- Molecule 3: KETOACYL SYNTHASE

Chain F:  99%



- Molecule 4: KETOACYL REDUCTASE

Chain G:  100%

There are no outlier residues recorded for this chain.

- Molecule 4: KETOACYL REDUCTASE

Chain H:  100%

There are no outlier residues recorded for this chain.

- Molecule 4: KETOACYL REDUCTASE

Chain I:  100%

There are no outlier residues recorded for this chain.

- Molecule 4: KETOACYL REDUCTASE

Chain J:  100%

There are no outlier residues recorded for this chain.

- Molecule 4: KETOACYL REDUCTASE

Chain K:  100%

There are no outlier residues recorded for this chain.

- Molecule 4: KETOACYL REDUCTASE

Chain L:  100%

There are no outlier residues recorded for this chain.

- Molecule 5: DEHYDRATASE

Chain S:  100%



- Molecule 5: DEHYDRATASE

Chain T:  100%



- Molecule 5: DEHYDRATASE

Chain U:  100%



- Molecule 5: DEHYDRATASE

Chain V:  100%

There are no outlier residues recorded for this chain.

- Molecule 5: DEHYDRATASE

Chain W:  100%

There are no outlier residues recorded for this chain.

- Molecule 5: DEHYDRATASE

Chain X:  100%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	217.00Å 415.00Å 222.00Å 90.00° 111.50° 90.00°	Depositor
Resolution (Å)	500.00 – 4.20	Depositor
% Data completeness (in resolution range)	(Not available) (500.00-4.20)	Depositor
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	?	Depositor
R, R_{free}	(Not available) , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	10404	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

There are no protein, RNA or DNA chains available to summarize Z scores of covalent bonds and angles.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	0	226	0	0	0	0
1	1	226	0	0	0	0
1	2	226	0	0	0	0
1	3	226	0	0	0	0
1	Y	226	0	0	0	0
1	Z	226	0	0	0	0
2	4	305	0	0	2	0
2	5	305	0	0	2	0
2	6	305	0	0	2	0
2	7	305	0	0	1	0
2	8	305	0	0	1	0
2	9	305	0	0	2	0
2	M	305	0	0	1	0
2	N	305	0	0	1	0
2	O	305	0	0	1	0
2	P	305	0	0	1	0
2	Q	305	0	0	1	0
2	R	305	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	406	0	0	0	0
3	B	406	0	0	0	0
3	C	406	0	0	10	0
3	D	406	0	0	10	0
3	E	406	0	0	3	0
3	F	406	0	0	3	0
4	G	244	0	0	0	0
4	H	244	0	0	0	0
4	I	244	0	0	0	0
4	J	244	0	0	0	0
4	K	244	0	0	0	0
4	L	244	0	0	0	0
5	S	248	0	0	1	0
5	T	248	0	0	1	0
5	U	248	0	0	1	0
5	V	248	0	0	0	0
5	W	248	0	0	0	0
5	X	248	0	0	1	0
All	All	10404	0	0	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 29 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:138:MET:CA	3:D:107:GLY:CA	1.82	1.56
3:C:107:GLY:CA	3:D:138:MET:CA	1.95	1.44
3:C:115:PHE:CA	3:D:114:VAL:CA	2.16	1.24
3:E:266:GLY:CA	3:F:153:HIS:CA	2.31	1.09
3:C:200:GLU:CA	3:D:131:PRO:CA	2.44	0.96

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.