



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 03:32 PM GMT

PDB ID : 4CFQ
Title : Ca-bound truncated (delta13C) and C3S, C81S and C86S mutated S100A4 complexed with non-muscle myosin IIA
Authors : Duelli, A.; Kiss, B.; Lundholm, I.; Bodor, A.; Radnai, L.; Petoukhov, M.; Svergun, D.; Nyitray, L.; Katona, G.
Deposited on : 2013-11-19
Resolution : 1.37 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1918 (1.40-1.36)
Clashscore	102246	2042 (1.40-1.36)
Ramachandran outliers	100387	1993 (1.40-1.36)
Sidechain outliers	100360	1992 (1.40-1.36)
RSRZ outliers	91569	1917 (1.40-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	91	<div> <div>10%</div> <div>84%</div> <div>10%</div> <div>7%</div> </div>
1	B	91	<div> <div>4%</div> <div>84%</div> <div>9%</div> <div>8%</div> </div>
1	C	91	<div> <div>15%</div> <div>85%</div> <div>9%</div> <div>7%</div> </div>
1	D	91	<div> <div>7%</div> <div>85%</div> <div>7%</div> <div>8%</div> </div>
2	Q	45	<div> <div>16%</div> <div>42%</div> <div>16%</div> <div>40%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	R	45	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CA	B	502	-	-	-	X
3	CA	D	502	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6614 atoms, of which 3138 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN S100-A4.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	85	Total	C	H	N	O	S	18	1	0
			1354	430	675	108	135	6			
1	B	84	Total	C	H	N	O	S	4	1	0
			1357	429	680	110	132	6			
1	C	85	Total	C	H	N	O	S	30	1	0
			1347	429	667	108	137	6			
1	D	84	Total	C	H	N	O	S	0	2	0
			1358	429	680	110	134	5			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	EXPRESSION TAG	UNP P26447
A	-1	SER	-	EXPRESSION TAG	UNP P26447
A	0	HIS	-	EXPRESSION TAG	UNP P26447
A	3	SER	CYS	ENGINEERED MUTATION	UNP P26447
A	81	SER	CYS	ENGINEERED MUTATION	UNP P26447
A	86	SER	CYS	ENGINEERED MUTATION	UNP P26447
B	-2	GLY	-	EXPRESSION TAG	UNP P26447
B	-1	SER	-	EXPRESSION TAG	UNP P26447
B	0	HIS	-	EXPRESSION TAG	UNP P26447
B	3	SER	CYS	ENGINEERED MUTATION	UNP P26447
B	81	SER	CYS	ENGINEERED MUTATION	UNP P26447
B	86	SER	CYS	ENGINEERED MUTATION	UNP P26447
C	-2	GLY	-	EXPRESSION TAG	UNP P26447
C	-1	SER	-	EXPRESSION TAG	UNP P26447
C	0	HIS	-	EXPRESSION TAG	UNP P26447
C	3	SER	CYS	ENGINEERED MUTATION	UNP P26447
C	81	SER	CYS	ENGINEERED MUTATION	UNP P26447
C	86	SER	CYS	ENGINEERED MUTATION	UNP P26447
D	-2	GLY	-	EXPRESSION TAG	UNP P26447
D	-1	SER	-	EXPRESSION TAG	UNP P26447
D	0	HIS	-	EXPRESSION TAG	UNP P26447

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	3	SER	CYS	ENGINEERED MUTATION	UNP P26447
D	81	SER	CYS	ENGINEERED MUTATION	UNP P26447
D	86	SER	CYS	ENGINEERED MUTATION	UNP P26447

- Molecule 2 is a protein called MYOSIN-9.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	Q	27	Total	C	H	N	O	S	16	0	0
			420	127	209	40	43	1			
2	R	29	Total	C	H	N	O	S	12	0	0
			452	137	227	42	45	1			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Q	1893	TYR	ARG	ENGINEERED MUTATION	UNP P35579
R	1893	TYR	ARG	ENGINEERED MUTATION	UNP P35579

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	2	Total	Ca	0	0
			2	2		
3	A	2	Total	Ca	0	0
			2	2		
3	D	2	Total	Ca	0	0
			2	2		
3	C	2	Total	Ca	0	0
			2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	88	Total	O	0	0
			88	88		
4	B	65	Total	O	0	0
			65	65		
4	C	71	Total	O	0	0
			71	71		
4	D	68	Total	O	0	0
			68	68		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	Q	15	Total	O	0	0
			15	15		
4	R	11	Total	O	0	0
			11	11		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PROTEIN S100-A4




- Molecule 1: PROTEIN S100-A4




- Molecule 1: PROTEIN S100-A4




- Molecule 1: PROTEIN S100-A4




- Molecule 2: MYOSIN-9



- Molecule 2: MYOSIN-9





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	38.72Å 38.79Å 71.43Å 88.65° 75.54° 72.17°	Depositor
Resolution (Å)	31.77 – 1.37 36.87 – 1.37	Depositor EDS
% Data completeness (in resolution range)	98.6 (31.77-1.37) 96.7 (36.87-1.37)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.62 (at 1.37Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.161 , 0.193 0.161 , 0.193	Depositor DCC
R_{free} test set	2378 reflections (3.00%)	DCC
Wilson B-factor (Å ²)	10.6	Xtriage
Anisotropy	0.034	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.45 , 51.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 79285 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6614	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/691	0.60	0/923
1	B	0.55	0/690	0.63	0/921
1	C	0.42	0/692	0.57	0/924
1	D	0.54	0/694	0.72	1/927 (0.1%)
2	Q	0.51	0/212	1.03	3/283 (1.1%)
2	R	0.58	0/226	1.04	1/303 (0.3%)
All	All	0.50	0/3205	0.70	5/4281 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	R	0	3

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	57	LYS	CD-CE-NZ	10.12	134.98	111.70
2	R	1926	LEU	CA-CB-CG	6.63	130.56	115.30
2	Q	1926	LEU	CA-CB-CG	5.40	127.73	115.30
2	Q	1912	ARG	NE-CZ-NH1	-5.37	117.61	120.30
2	Q	1926	LEU	CB-CG-CD2	5.14	119.73	111.00

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	R	1924	GLY	Peptide
2	R	1925	ASP	Peptide
2	R	1926	LEU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	679	675	676	4	0
1	B	677	680	681	4	0
1	C	680	667	668	4	0
1	D	678	680	682	7	1
2	Q	211	209	209	4	0
2	R	225	227	227	9	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
4	A	88	0	0	0	0
4	B	65	0	0	1	1
4	C	71	0	0	0	1
4	D	68	0	0	0	0
4	Q	15	0	0	2	0
4	R	11	0	0	0	1
All	All	3476	3138	3143	27	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 27 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:57:LYS:HE2	2:R:1926:LEU:HD12	1.66	0.78
1:B:61:ASN:ND2	4:B:2046:HOH:O	2.19	0.75
2:Q:1923:ARG:NH1	4:Q:2013:HOH:O	2.26	0.64
2:R:1923:ARG:N	2:R:1924:GLY:HA3	2.14	0.62
2:Q:1919:ASN:OD1	4:Q:2008:HOH:O	2.16	0.61

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:2049:HOH:O	4:R:2002:HOH:O[1_644]	2.17	0.03
1:D:48:LYS:NZ	4:C:2068:HOH:O[1_655]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	82/91 (90%)	82 (100%)	0	0	100	100
1	B	83/91 (91%)	81 (98%)	2 (2%)	0	100	100
1	C	82/91 (90%)	82 (100%)	0	0	100	100
1	D	84/91 (92%)	82 (98%)	2 (2%)	0	100	100
2	Q	25/45 (56%)	21 (84%)	3 (12%)	1 (4%)	4	0
2	R	27/45 (60%)	24 (89%)	0	3 (11%)	0	0
All	All	383/454 (84%)	372 (97%)	7 (2%)	4 (1%)	17	3

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	R	1925	ASP
2	Q	1926	LEU
2	R	1927	PRO
2	R	1924	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	78/82 (95%)	76 (97%)	2 (3%)	54	16
1	B	77/82 (94%)	75 (97%)	2 (3%)	54	16
1	C	78/82 (95%)	77 (99%)	1 (1%)	76	46
1	D	78/82 (95%)	77 (99%)	1 (1%)	76	46
2	Q	23/40 (58%)	20 (87%)	3 (13%)	5	0
2	R	25/40 (62%)	22 (88%)	3 (12%)	6	0
All	All	359/408 (88%)	347 (97%)	12 (3%)	44	11

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	73	GLN
2	Q	1904	THR
2	R	1906	THR
1	C	85	MET
2	Q	1926	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	73	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	85/91 (93%)	0.69	9 (10%) 8 8	9, 17, 33, 47	3 (3%)
1	B	84/91 (92%)	0.47	4 (4%) 34 33	6, 12, 27, 42	2 (2%)
1	C	85/91 (93%)	0.70	14 (16%) 2 3	9, 18, 34, 44	5 (5%)
1	D	84/91 (92%)	0.50	6 (7%) 19 18	6, 12, 30, 42	1 (1%)
2	Q	27/45 (60%)	1.76	7 (25%) 1 1	8, 19, 39, 42	5 (18%)
2	R	29/45 (64%)	2.33	12 (41%) 0 1	7, 21, 40, 40	7 (24%)
All	All	394/454 (86%)	0.80	52 (13%) 4 4	6, 15, 38, 47	23 (5%)

The worst 5 of 52 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	R	1926	LEU	9.6
2	Q	1902	ASP	8.6
2	R	1923	ARG	7.5
2	Q	1928	PHE	6.6
1	D	55	PHE	6.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	CA	B	502	1/1	1.00	0.12	5.34	5,5,5,5	0
3	CA	D	502	1/1	1.00	0.12	4.42	5,5,5,5	0
3	CA	B	501	1/1	1.00	0.10	0.60	5,5,5,5	0
3	CA	D	501	1/1	1.00	0.11	0.44	5,5,5,5	0
3	CA	A	502	1/1	1.00	0.10	0.03	9,9,9,9	0
3	CA	C	502	1/1	1.00	0.09	-0.13	9,9,9,9	0
3	CA	C	501	1/1	0.99	0.08	-0.99	26,26,26,26	0
3	CA	A	501	1/1	1.00	0.07	-1.15	24,24,24,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.