



wwPDB EM Map/Model Validation Report ⓘ

Apr 10, 2016 – 01:46 PM BST

PDB ID : 2CGT
EMDB ID: : EMD-1202
Title : GROEL-ADP-gp31 COMPLEX
Authors : Clare, D.K.; Bakkes, P.J.; van Heerikhuizen, H.; van der Vies, S.M.; Saibil, H.R.
Deposited on : 2006-03-09
Resolution : 8.20 Å(reported)
Based on PDB ID : 1A0N,1G31

This is a wwPDB EM Map/Model Validation Report for a publicly released PDB/EMDB entry.
For rigid body fitted models, validation errors reported here could stem from errors in the original structure(s) used in the fitting.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/EMValidationReportHelp>


MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : trunk27241

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 8.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.















| Metric | Whole archive (#Entries) | EM structures (#Entries) |
|-----------------------|--------------------------|--------------------------|
| Clashscore | 114402 | 924 |
| Ramachandran outliers | 111179 | 726 |
| Sidechain outliers | 111093 | 686 |

The table below summarises the geometric issues observed across the polymeric chains. The red, orange, yellow and green segments on the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1 | A | 547 | 62% 27% 6% . |
| 1 | B | 547 | 62% 27% 6% . |
| 1 | C | 547 | 62% 27% 6% . |
| 1 | D | 547 | 63% 26% 6% . |
| 1 | E | 547 | 62% 27% 6% . |
| 1 | F | 547 | 63% 27% 6% . |
| 1 | G | 547 | 62% 27% 6% . |
| 1 | H | 547 | 56% 32% 5% . 6% |
| 1 | I | 547 | 55% 33% 5% . 6% |

Continued on next page...

Continued from previous page...

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|--|
| 1 | J | 547 |  55% 33% 5% • 6% |
| 1 | K | 547 |  55% 32% 6% • 6% |
| 1 | L | 547 |  55% 33% 5% • 6% |
| 1 | M | 547 |  55% 33% 6% • 6% |
| 1 | N | 547 |  55% 33% 5% • 6% |
| 2 | O | 111 |  47% 27% • 25% |
| 2 | P | 111 |  47% 26% •• 25% |
| 2 | Q | 111 |  47% 26% •• 25% |
| 2 | R | 111 |  45% 28% •• 25% |
| 2 | S | 111 |  45% 29% • 25% |
| 2 | T | 111 |  45% 29% • 25% |
| 2 | U | 111 |  45% 29% • 25% |

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 57953 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 60 KDA GROEL.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 1 | A | 524 | Total | C | N | O | S | 0 | 0 |
| | | | 3855 | 2397 | 665 | 773 | 20 | | |
| 1 | B | 524 | Total | C | N | O | S | 0 | 0 |
| | | | 3855 | 2397 | 665 | 773 | 20 | | |
| 1 | C | 524 | Total | C | N | O | S | 0 | 0 |
| | | | 3855 | 2397 | 665 | 773 | 20 | | |
| 1 | D | 524 | Total | C | N | O | S | 0 | 0 |
| | | | 3855 | 2397 | 665 | 773 | 20 | | |
| 1 | E | 524 | Total | C | N | O | S | 0 | 0 |
| | | | 3855 | 2397 | 665 | 773 | 20 | | |
| 1 | F | 524 | Total | C | N | O | S | 0 | 0 |
| | | | 3855 | 2397 | 665 | 773 | 20 | | |
| 1 | G | 524 | Total | C | N | O | S | 0 | 0 |
| | | | 3855 | 2397 | 665 | 773 | 20 | | |
| 1 | H | 515 | Total | C | N | O | S | 0 | 0 |
| | | | 3783 | 2354 | 656 | 753 | 20 | | |
| 1 | I | 515 | Total | C | N | O | S | 0 | 0 |
| | | | 3783 | 2354 | 656 | 753 | 20 | | |
| 1 | J | 515 | Total | C | N | O | S | 0 | 0 |
| | | | 3783 | 2354 | 656 | 753 | 20 | | |
| 1 | K | 515 | Total | C | N | O | S | 0 | 0 |
| | | | 3783 | 2354 | 656 | 753 | 20 | | |
| 1 | L | 515 | Total | C | N | O | S | 0 | 0 |
| | | | 3783 | 2354 | 656 | 753 | 20 | | |
| 1 | M | 515 | Total | C | N | O | S | 0 | 0 |
| | | | 3783 | 2354 | 656 | 753 | 20 | | |
| 1 | N | 515 | Total | C | N | O | S | 0 | 0 |
| | | | 3783 | 2354 | 656 | 753 | 20 | | |

- Molecule 2 is a protein called CAPSID ASSEMBLY PROTEIN GP31.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 2 | O | 83 | Total | C | N | O | S | 0 | 0 |
| | | | 641 | 417 | 106 | 114 | 4 | | |

Continued on next page...


Continued from previous page...

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|--------------|----------|----------|----------|--------|---------|-------|
| 2 | P | 83 | Total 641 | C 417 | N 106 | O 114 | S 4 | 0 | 0 |
| 2 | Q | 83 | Total 641 | C 417 | N 106 | O 114 | S 4 | 0 | 0 |
| 2 | R | 83 | Total 641 | C 417 | N 106 | O 114 | S 4 | 0 | 0 |
| 2 | S | 83 | Total 641 | C 417 | N 106 | O 114 | S 4 | 0 | 0 |
| 2 | T | 83 | Total 641 | C 417 | N 106 | O 114 | S 4 | 0 | 0 |
| 2 | U | 83 | Total 641 | C 417 | N 106 | O 114 | S 4 | 0 | 0 |


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 60 KDA GROEL




• Molecule 1: 60 KDA GROEL



ALA
GLY
GLY
MET
GLY
GLY
MET
GLY
GLY
MET
MET

• Molecule 1: 60 KDA GROEL


Chain C:  62% 27% 6% •



ALA
GLY
GLY
MET
GLY
GLY
MET
GLY
GLY
MET
MET

• Molecule 1: 60 KDA GROEL


Chain D:  63% 26% 6% •




GLY
GLY
MET
GLY
GLY
MET
GLY
GLY
MET
MET

• Molecule 1: 60 KDA GROEL


Chain E:  62% 27% 6% •






• Molecule 1: 60 KDA GROEL

Chain H: 56% 32% 5% • 6%




• Molecule 1: 60 KDA GROEL


Chain I: 55% 33% 5% • 6%




- Molecule 1: 60 KDA GROEL




- Molecule 1: 60 KDA GROEL




- Molecule 1: 60 KDA GROEL






• Molecule 1: 60 KDA GROEL



Chain M: 55% 33% 6% • 6%





• Molecule 1: 60 KDA GROEL

Chain N: 55% 33% 5% • 6%







• Molecule 2: CAPSID ASSEMBLY PROTEIN GP31




• Molecule 2: CAPSID ASSEMBLY PROTEIN GP31




• Molecule 2: CAPSID ASSEMBLY PROTEIN GP31




• Molecule 2: CAPSID ASSEMBLY PROTEIN GP31




• Molecule 2: CAPSID ASSEMBLY PROTEIN GP31






● Molecule 2: CAPSID ASSEMBLY PROTEIN GP31



● Molecule 2: CAPSID ASSEMBLY PROTEIN GP31



4 Experimental information

| Property | Value | Source |
|--------------------------------------|--------------------------------------|-----------|
| Reconstruction method | SINGLE PARTICLE | Depositor |
| Imposed symmetry | POINT, Not provided | Depositor |
| Number of images | Not provided | Depositor |
| Resolution determination method | Not provided | Depositor |
| CTF correction method | FULL CORRECTION ON 2D CLASS AVERAGES | Depositor |
| Microscope | FEI TECNAI F20 | Depositor |
| Voltage (kV) | 200 | Depositor |
| Electron dose ($e^-/\text{\AA}^2$) | Not provided | Depositor |
| Minimum defocus (nm) | 1300 | Depositor |
| Maximum defocus (nm) | 3300 | Depositor |
| Magnification | 50000 | Depositor |
| Image detector | KODAK SO163 FILM | Depositor |

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|-----------------|-------------|------------------|
| | | RMSZ | # $ Z > 2$ | RMSZ | # $ Z > 2$ |
| 1 | A | 1.07 | 3/3882 (0.1%) | 1.23 | 27/5240 (0.5%) |
| 1 | B | 1.07 | 3/3882 (0.1%) | 1.23 | 27/5240 (0.5%) |
| 1 | C | 1.07 | 3/3882 (0.1%) | 1.23 | 27/5240 (0.5%) |
| 1 | D | 1.07 | 3/3882 (0.1%) | 1.23 | 27/5240 (0.5%) |
| 1 | E | 1.07 | 3/3882 (0.1%) | 1.23 | 27/5240 (0.5%) |
| 1 | F | 1.07 | 3/3882 (0.1%) | 1.23 | 27/5240 (0.5%) |
| 1 | G | 1.07 | 3/3882 (0.1%) | 1.23 | 27/5240 (0.5%) |
| 1 | H | 0.68 | 1/3806 (0.0%) | 1.04 | 15/5130 (0.3%) |
| 1 | I | 0.69 | 1/3806 (0.0%) | 1.04 | 15/5130 (0.3%) |
| 1 | J | 0.69 | 1/3806 (0.0%) | 1.04 | 15/5130 (0.3%) |
| 1 | K | 0.69 | 1/3806 (0.0%) | 1.04 | 15/5130 (0.3%) |
| 1 | L | 0.69 | 1/3806 (0.0%) | 1.04 | 15/5130 (0.3%) |
| 1 | M | 0.69 | 1/3806 (0.0%) | 1.04 | 15/5130 (0.3%) |
| 1 | N | 0.69 | 1/3806 (0.0%) | 1.04 | 15/5130 (0.3%) |
| 2 | O | 0.59 | 0/657 | 0.79 | 1/894 (0.1%) |
| 2 | P | 0.59 | 0/657 | 0.79 | 1/894 (0.1%) |
| 2 | Q | 0.59 | 0/657 | 0.79 | 1/894 (0.1%) |
| 2 | R | 0.59 | 0/657 | 0.79 | 1/894 (0.1%) |
| 2 | S | 0.59 | 0/657 | 0.79 | 1/894 (0.1%) |
| 2 | T | 0.59 | 0/657 | 0.79 | 1/894 (0.1%) |
| 2 | U | 0.59 | 0/657 | 0.79 | 1/894 (0.1%) |
| All | All | 0.88 | 28/58415 (0.0%) | 1.12 | 301/78848 (0.4%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 1 | A | 0 | 1 |
| 1 | B | 0 | 1 |
| 1 | C | 0 | 1 |
| 1 | D | 0 | 1 |

Continued on next page...

Continued from previous page...

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 1 | E | 0 | 1 |
| 1 | F | 0 | 1 |
| 1 | G | 0 | 1 |
| 1 | H | 0 | 1 |
| 1 | I | 0 | 1 |
| 1 | J | 0 | 1 |
| 1 | K | 0 | 1 |
| 1 | L | 0 | 1 |
| 1 | M | 0 | 1 |
| 1 | N | 0 | 1 |
| All | All | 0 | 14 |

The worst 5 of 28 bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|-------|--------|-------------|----------|
| 1 | F | 191 | GLU | C-N | -42.62 | 0.56 | 1.33 |
| 1 | A | 191 | GLU | C-N | -42.60 | 0.56 | 1.33 |
| 1 | C | 191 | GLU | C-N | -42.60 | 0.56 | 1.33 |
| 1 | E | 191 | GLU | C-N | -42.60 | 0.56 | 1.33 |
| 1 | D | 191 | GLU | C-N | -42.59 | 0.56 | 1.33 |

The worst 5 of 301 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-------|--------|-------------|----------|
| 1 | H | 373 | ALA | O-C-N | -33.61 | 66.06 | 123.20 |
| 1 | J | 373 | ALA | O-C-N | -33.61 | 66.06 | 123.20 |
| 1 | N | 373 | ALA | O-C-N | -33.61 | 66.06 | 123.20 |
| 1 | K | 373 | ALA | O-C-N | -33.60 | 66.07 | 123.20 |
| 1 | L | 373 | ALA | O-C-N | -33.60 | 66.07 | 123.20 |

There are no chirality outliers.

5 of 14 planarity outliers are listed below:

| Mol | Chain | Res | Type | Group |
|-----|-------|-----|------|-----------|
| 1 | A | 191 | GLU | Mainchain |
| 1 | B | 191 | GLU | Mainchain |
| 1 | C | 191 | GLU | Mainchain |
| 1 | D | 191 | GLU | Mainchain |
| 1 | E | 191 | GLU | Mainchain |

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | A | 3855 | 0 | 3971 | 193 | 0 |
| 1 | B | 3855 | 0 | 3971 | 194 | 0 |
| 1 | C | 3855 | 0 | 3971 | 197 | 0 |
| 1 | D | 3855 | 0 | 3971 | 189 | 0 |
| 1 | E | 3855 | 0 | 3971 | 199 | 0 |
| 1 | F | 3855 | 0 | 3971 | 196 | 0 |
| 1 | G | 3855 | 0 | 3971 | 198 | 0 |
| 1 | H | 3783 | 0 | 3916 | 226 | 0 |
| 1 | I | 3783 | 0 | 3916 | 233 | 0 |
| 1 | J | 3783 | 0 | 3916 | 229 | 0 |
| 1 | K | 3783 | 0 | 3916 | 234 | 0 |
| 1 | L | 3783 | 0 | 3916 | 228 | 0 |
| 1 | M | 3783 | 0 | 3916 | 236 | 0 |
| 1 | N | 3783 | 0 | 3916 | 234 | 0 |
| 2 | O | 641 | 0 | 652 | 60 | 0 |
| 2 | P | 641 | 0 | 652 | 61 | 0 |
| 2 | Q | 641 | 0 | 652 | 61 | 0 |
| 2 | R | 641 | 0 | 652 | 61 | 0 |
| 2 | S | 641 | 0 | 652 | 60 | 0 |
| 2 | T | 641 | 0 | 652 | 60 | 0 |
| 2 | U | 641 | 0 | 652 | 61 | 0 |
| All | All | 57953 | 0 | 59773 | 2885 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.

The worst 5 of 2885 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|----------------|-----------------|--------------------------|-------------------|
| 1:L:138:CYS:CB | 1:L:410:GLY:HA2 | 1.38 | 1.52 |
| 1:K:138:CYS:CB | 1:K:410:GLY:HA2 | 1.38 | 1.51 |
| 1:N:138:CYS:CB | 1:N:410:GLY:HA2 | 1.38 | 1.51 |
| 1:M:138:CYS:CB | 1:M:410:GLY:HA2 | 1.38 | 1.50 |
| 1:H:138:CYS:CB | 1:H:410:GLY:HA2 | 1.38 | 1.49 |

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|-----------------|------------|----------|----------|-------------|-----|
| 1 | A | 520/547 (95%) | 483 (93%) | 31 (6%) | 6 (1%) | 16 | 61 |
| 1 | B | 520/547 (95%) | 483 (93%) | 31 (6%) | 6 (1%) | 16 | 61 |
| 1 | C | 520/547 (95%) | 483 (93%) | 31 (6%) | 6 (1%) | 16 | 61 |
| 1 | D | 520/547 (95%) | 483 (93%) | 31 (6%) | 6 (1%) | 16 | 61 |
| 1 | E | 520/547 (95%) | 483 (93%) | 31 (6%) | 6 (1%) | 16 | 61 |
| 1 | F | 520/547 (95%) | 483 (93%) | 31 (6%) | 6 (1%) | 16 | 61 |
| 1 | G | 520/547 (95%) | 483 (93%) | 31 (6%) | 6 (1%) | 16 | 61 |
| 1 | H | 505/547 (92%) | 447 (88%) | 45 (9%) | 13 (3%) | 7 | 45 |
| 1 | I | 505/547 (92%) | 447 (88%) | 45 (9%) | 13 (3%) | 7 | 45 |
| 1 | J | 505/547 (92%) | 447 (88%) | 45 (9%) | 13 (3%) | 7 | 45 |
| 1 | K | 505/547 (92%) | 447 (88%) | 45 (9%) | 13 (3%) | 7 | 45 |
| 1 | L | 505/547 (92%) | 447 (88%) | 45 (9%) | 13 (3%) | 7 | 45 |
| 1 | M | 505/547 (92%) | 447 (88%) | 45 (9%) | 13 (3%) | 7 | 45 |
| 1 | N | 505/547 (92%) | 447 (88%) | 45 (9%) | 13 (3%) | 7 | 45 |
| 2 | O | 79/111 (71%) | 76 (96%) | 3 (4%) | 0 | 100 | 100 |
| 2 | P | 79/111 (71%) | 76 (96%) | 3 (4%) | 0 | 100 | 100 |
| 2 | Q | 79/111 (71%) | 76 (96%) | 3 (4%) | 0 | 100 | 100 |
| 2 | R | 79/111 (71%) | 76 (96%) | 3 (4%) | 0 | 100 | 100 |
| 2 | S | 79/111 (71%) | 76 (96%) | 3 (4%) | 0 | 100 | 100 |
| 2 | T | 79/111 (71%) | 76 (96%) | 3 (4%) | 0 | 100 | 100 |
| 2 | U | 79/111 (71%) | 76 (96%) | 3 (4%) | 0 | 100 | 100 |
| All | All | 7728/8435 (92%) | 7042 (91%) | 553 (7%) | 133 (2%) | 16 | 55 |

5 of 133 Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | A | 44 | PHE |
| 1 | A | 137 | PRO |
| 1 | A | 337 | GLY |
| 1 | B | 44 | PHE |
| 1 | B | 137 | PRO |

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|---------------|-----------|----------|-------------|----|
| 1 | A | 404/414 (98%) | 352 (87%) | 52 (13%) | 5 | 28 |
| 1 | B | 404/414 (98%) | 352 (87%) | 52 (13%) | 5 | 28 |
| 1 | C | 404/414 (98%) | 352 (87%) | 52 (13%) | 5 | 28 |
| 1 | D | 404/414 (98%) | 352 (87%) | 52 (13%) | 5 | 28 |
| 1 | E | 404/414 (98%) | 352 (87%) | 52 (13%) | 5 | 28 |
| 1 | F | 404/414 (98%) | 352 (87%) | 52 (13%) | 5 | 28 |
| 1 | G | 404/414 (98%) | 352 (87%) | 52 (13%) | 5 | 28 |
| 1 | H | 396/414 (96%) | 347 (88%) | 49 (12%) | 6 | 30 |
| 1 | I | 396/414 (96%) | 347 (88%) | 49 (12%) | 6 | 30 |
| 1 | J | 396/414 (96%) | 347 (88%) | 49 (12%) | 6 | 30 |
| 1 | K | 396/414 (96%) | 347 (88%) | 49 (12%) | 6 | 30 |
| 1 | L | 396/414 (96%) | 347 (88%) | 49 (12%) | 6 | 30 |
| 1 | M | 396/414 (96%) | 347 (88%) | 49 (12%) | 6 | 30 |
| 1 | N | 396/414 (96%) | 347 (88%) | 49 (12%) | 6 | 30 |
| 2 | O | 73/96 (76%) | 71 (97%) | 2 (3%) | 52 | 79 |
| 2 | P | 73/96 (76%) | 71 (97%) | 2 (3%) | 52 | 79 |
| 2 | Q | 73/96 (76%) | 71 (97%) | 2 (3%) | 52 | 79 |
| 2 | R | 73/96 (76%) | 71 (97%) | 2 (3%) | 52 | 79 |
| 2 | S | 73/96 (76%) | 71 (97%) | 2 (3%) | 52 | 79 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|-----------------|------------|-----------|-------------|----|
| 2 | T | 73/96 (76%) | 71 (97%) | 2 (3%) | 52 | 79 |
| 2 | U | 73/96 (76%) | 71 (97%) | 2 (3%) | 52 | 79 |
| All | All | 6111/6468 (94%) | 5390 (88%) | 721 (12%) | 11 | 32 |

5 of 721 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | G | 147 | VAL |
| 1 | H | 422 | VAL |
| 1 | N | 74 | VAL |
| 1 | G | 225 | LYS |
| 1 | H | 23 | LEU |

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 152 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | I | 97 | GLN |
| 1 | K | 146 | GLN |
| 2 | S | 81 | HIS |
| 1 | I | 153 | ASN |
| 1 | J | 146 | GLN |

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.