



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 10:09 PM GMT

PDB ID : 5CIT
Title : Ran GDP wild type monoclinic crystal form
Authors : Vetter, I.R.; Brucker, S.
Deposited on : 2015-07-13
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1609 (1.76-1.76)
Clashscore	102246	1730 (1.76-1.76)
Ramachandran outliers	100387	1711 (1.76-1.76)
Sidechain outliers	100360	1711 (1.76-1.76)
RSRZ outliers	91569	1610 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	216	
1	B	216	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3328 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GTP-binding nuclear protein Ran.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	199	Total	C	N	O	S	0	0	0
			1600	1036	277	281	6			
1	B	197	Total	C	N	O	S	0	0	0
			1584	1027	274	277	6			

- Molecule 2 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Mg 1	0	0
3	A	1	Total 1	Mg 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	39	Total 39	O 39	0	0
4	B	47	Total 47	O 47	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: GTP-binding nuclear protein Ran



• Molecule 1: GTP-binding nuclear protein Ran



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.01Å 59.23Å 64.77Å 90.00° 96.95° 90.00°	Depositor
Resolution (Å)	19.81 – 1.75 19.84 – 1.75	Depositor EDS
% Data completeness (in resolution range)	96.2 (19.81-1.75) 96.3 (19.84-1.75)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.09 (at 1.76Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.216 , 0.262 0.216 , 0.261	Depositor DCC
R_{free} test set	2123 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	29.5	Xtriage
Anisotropy	0.058	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 51.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 42450 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3328	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.89	0/1640	1.05	5/2222 (0.2%)
1	B	1.02	0/1624	1.11	9/2200 (0.4%)
All	All	0.96	0/3264	1.08	14/4422 (0.3%)

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	92	VAL	CG1-CB-CG2	9.22	125.65	110.90
1	B	27	VAL	CG1-CB-CG2	7.74	123.28	110.90
1	A	148	ASP	CB-CG-OD2	7.22	124.80	118.30
1	B	27	VAL	CB-CA-C	-6.53	99.00	111.40
1	B	144	LEU	CA-CB-CG	6.44	130.11	115.30
1	A	189	MET	CG-SD-CE	6.39	110.43	100.20
1	A	65	ASP	CB-CG-OD1	6.31	123.98	118.30
1	A	129	ARG	NE-CZ-NH1	-6.21	117.19	120.30
1	B	148	ASP	CB-CG-OD1	-5.77	113.11	118.30
1	B	81	ILE	CB-CA-C	-5.76	100.08	111.60
1	B	29	ARG	NE-CZ-NH2	-5.52	117.54	120.30
1	B	92	VAL	CG1-CB-CG2	5.50	119.71	110.90
1	B	81	ILE	CG1-CB-CG2	5.33	123.12	111.40
1	B	27	VAL	CA-CB-CG2	5.28	118.82	110.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1600	0	1615	35	0
1	B	1584	0	1600	42	0
2	A	28	0	12	1	0
2	B	28	0	12	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	39	0	0	5	0
4	B	47	0	0	4	0
All	All	3328	0	3239	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:134:LYS:HD2	1:B:134:LYS:H	1.33	0.93
1:B:55:ASN:ND2	1:B:175:GLU:H	1.76	0.83
1:B:53:HIS:HD2	4:B:442:HOH:O	1.60	0.82
1:B:179:MET:HG3	1:B:180:PRO:HD2	1.63	0.80
1:B:117:ILE:HB	1:B:144:LEU:HD22	1.63	0.79
1:B:103:ASN:ND2	1:B:106:ARG:HH22	1.82	0.77
1:B:103:ASN:HD22	1:B:106:ARG:NH2	1.83	0.76
1:A:199:HIS:HD2	4:A:433:HOH:O	1.69	0.75
1:B:142:LYS:HB2	1:B:144:LEU:HG	1.69	0.73
1:B:134:LYS:CD	1:B:134:LYS:H	2.05	0.70
1:A:166:ARG:NH1	1:A:175:GLU:OE1	2.25	0.69
1:B:137:VAL:O	1:B:140:ARG:HG3	1.92	0.69
1:A:71:LYS:HE2	4:A:403:HOH:O	1.95	0.66
1:B:92:VAL:HG13	1:B:129:ARG:HG3	1.78	0.65
1:A:55:ASN:H	1:A:55:ASN:HD22	1.45	0.63
1:B:48:HIS:HE1	4:B:401:HOH:O	1.83	0.60
1:B:156:ASN:OD1	1:B:159:LYS:HE2	2.03	0.59
1:A:53:HIS:HD2	4:A:436:HOH:O	1.86	0.59
1:B:53:HIS:CD2	4:B:442:HOH:O	2.43	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:55:ASN:HD21	1:A:174:LEU:HA	1.69	0.57
1:A:105:HIS:O	1:A:109:VAL:HG12	2.05	0.56
1:B:93:THR:HG21	1:B:126:ILE:HG21	1.87	0.56
1:B:103:ASN:ND2	1:B:106:ARG:NH2	2.45	0.55
1:A:126:ILE:HD11	2:A:301:GDP:N2	2.22	0.55
1:B:179:MET:HG3	1:B:180:PRO:CD	2.35	0.54
1:B:137:VAL:HA	1:B:140:ARG:HH11	1.73	0.54
1:B:55:ASN:HD22	1:B:175:GLU:H	1.54	0.53
1:B:105:HIS:O	1:B:109:VAL:HG12	2.09	0.52
1:B:95:ARG:HB3	1:B:95:ARG:NH2	2.25	0.52
1:A:115:ILE:O	1:A:117:ILE:HG13	2.10	0.51
1:B:99:LYS:O	1:B:102:PRO:HD2	2.11	0.51
1:A:190:ASP:HB3	1:A:193:LEU:HB2	1.94	0.50
1:B:48:HIS:HD2	4:B:405:HOH:O	1.95	0.49
1:B:95:ARG:HB3	1:B:95:ARG:HH21	1.78	0.49
1:A:159:LYS:HB2	1:A:160:PRO:HD3	1.94	0.48
1:A:78:GLY:HA2	1:A:81:ILE:HG13	1.95	0.48
1:B:13:LEU:HD23	1:B:13:LEU:C	2.34	0.48
1:B:55:ASN:HD21	1:B:174:LEU:HA	1.78	0.48
1:A:199:HIS:CD2	4:A:433:HOH:O	2.55	0.47
1:A:137:VAL:HG12	1:A:140:ARG:NH1	2.30	0.47
1:A:48:HIS:HD2	4:A:409:HOH:O	1.98	0.47
1:A:55:ASN:ND2	1:A:175:GLU:H	2.12	0.46
1:B:99:LYS:HB3	1:B:99:LYS:HE2	1.56	0.46
1:A:139:HIS:CE1	1:A:140:ARG:HD3	2.51	0.46
1:B:56:ARG:HH11	1:B:171:ASP:HB2	1.81	0.46
1:A:110:ARG:HH22	1:A:111:VAL:HG22	1.81	0.46
1:B:145:GLN:HB2	1:B:163:TRP:CE2	2.51	0.45
1:B:137:VAL:HB	1:B:140:ARG:HD2	1.99	0.45
1:A:75:LEU:HG	1:A:75:LEU:H	1.55	0.45
1:A:145:GLN:HA	1:A:145:GLN:OE1	2.17	0.45
1:B:86:ALA:CB	1:B:108:LEU:HD21	2.47	0.45
1:A:196:GLN:NE2	1:A:200:ASP:OD2	2.50	0.44
1:A:40:VAL:HG11	1:A:43:LEU:HD12	1.98	0.44
1:B:55:ASN:HD22	1:B:55:ASN:H	1.66	0.44
1:A:92:VAL:HG13	1:A:129:ARG:HG3	1.99	0.44
1:A:69:GLN:HG2	1:A:75:LEU:CD2	2.48	0.43
1:A:193:LEU:HD13	1:A:197:TYR:CE1	2.54	0.43
1:B:92:VAL:HB	1:B:122:ASN:O	2.18	0.43
1:A:69:GLN:HG2	1:A:75:LEU:HD22	2.01	0.43
1:B:127:LYS:HB3	1:B:127:LYS:HE2	1.50	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:167:LYS:HA	1:B:167:LYS:HD2	1.83	0.42
1:B:184:PRO:HA	1:B:185:PRO:HD3	1.94	0.42
1:A:156:ASN:OD1	1:A:159:LYS:HE2	2.19	0.42
1:A:168:LEU:HD23	1:A:168:LEU:HA	1.85	0.42
1:A:54:THR:HA	1:A:175:GLU:O	2.20	0.42
1:A:118:VAL:HG22	1:A:163:TRP:CE3	2.55	0.42
1:A:55:ASN:H	1:A:55:ASN:ND2	2.16	0.41
1:A:142:LYS:HD2	1:A:142:LYS:HA	1.81	0.41
1:B:117:ILE:CB	1:B:144:LEU:HD22	2.43	0.41
1:B:93:THR:HG21	1:B:126:ILE:CG2	2.51	0.41
1:A:11:PHE:CG	1:A:168:LEU:HD13	2.55	0.41
1:B:117:ILE:HB	1:B:144:LEU:CD2	2.42	0.41
1:B:152:LYS:HD3	1:B:186:GLU:OE2	2.21	0.41
1:A:73:GLY:O	1:A:74:GLY:C	2.59	0.40
1:B:190:ASP:HB3	1:B:193:LEU:HB2	2.02	0.40
1:B:132:LYS:HB2	1:B:134:LYS:HD3	2.03	0.40
1:A:110:ARG:HB3	1:A:110:ARG:NH2	2.35	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	197/216 (91%)	188 (95%)	7 (4%)	2 (1%)	19	5
1	B	195/216 (90%)	186 (95%)	8 (4%)	1 (0%)	34	14
All	All	392/432 (91%)	374 (95%)	15 (4%)	3 (1%)	24	8

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	113	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	133	ALA
1	A	74	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	172/185 (93%)	157 (91%)	15 (9%)	13	1
1	B	170/185 (92%)	156 (92%)	14 (8%)	14	2
All	All	342/370 (92%)	313 (92%)	29 (8%)	13	2

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	LEU
1	A	50	LEU
1	A	55	ASN
1	A	56	ARG
1	A	75	LEU
1	A	77	ASP
1	A	92	VAL
1	A	96	VAL
1	A	110	ARG
1	A	132	LYS
1	A	134	LYS
1	A	135	SER
1	A	188	VAL
1	A	193	LEU
1	A	196	GLN
1	B	15	LEU
1	B	27	VAL
1	B	55	ASN
1	B	56	ARG
1	B	69	GLN
1	B	81	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	92	VAL
1	B	95	ARG
1	B	99	LYS
1	B	127	LYS
1	B	130	LYS
1	B	134	LYS
1	B	142	LYS
1	B	193	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	48	HIS
1	A	55	ASN
1	A	103	ASN
1	A	196	GLN
1	A	199	HIS
1	B	10	GLN
1	B	48	HIS
1	B	53	HIS
1	B	55	ASN
1	B	62	ASN
1	B	103	ASN
1	B	114	ASN
1	B	143	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	GDP	A	301	3	24,30,30	1.34	4 (16%)	26,47,47	1.98	7 (26%)
2	GDP	B	301	3	24,30,30	1.66	4 (16%)	26,47,47	1.44	5 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GDP	A	301	3	-	0/12/32/32	0/3/3/3
2	GDP	B	301	3	-	0/12/32/32	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	GDP	C2'-C1'	-2.63	1.49	1.53
2	A	301	GDP	PB-O3B	-2.31	1.46	1.54
2	A	301	GDP	PB-O1B	2.48	1.58	1.50
2	B	301	GDP	C6-N1	2.70	1.37	1.33
2	A	301	GDP	C5-C4	2.90	1.47	1.40
2	A	301	GDP	C6-C5	2.96	1.47	1.41
2	B	301	GDP	C5-C4	3.86	1.49	1.40
2	B	301	GDP	C6-C5	4.21	1.49	1.41

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	GDP	C6-C5-C4	-4.71	115.48	120.86
2	A	301	GDP	N3-C2-N1	-4.47	121.47	127.56
2	A	301	GDP	C1'-N9-C4	-3.81	122.55	126.81

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	GDP	C6-C5-C4	-3.23	117.17	120.86
2	B	301	GDP	C5-C6-N1	-3.03	119.56	123.52
2	A	301	GDP	C5-C6-N1	-2.31	120.50	123.52
2	A	301	GDP	O3'-C3'-C2'	-2.04	105.26	111.86
2	B	301	GDP	O2A-PA-O3A	2.04	114.02	105.27
2	B	301	GDP	C6-N1-C2	2.12	118.36	115.88
2	B	301	GDP	O2B-PB-O1B	2.44	118.59	110.63
2	A	301	GDP	O3B-PB-O2B	2.45	116.44	107.44
2	A	301	GDP	C6-N1-C2	4.24	120.85	115.88

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	GDP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	199/216 (92%)	0.83	37 (18%) 2 2	21, 37, 66, 97	0
1	B	197/216 (91%)	0.52	18 (9%) 11 14	19, 32, 58, 86	0
All	All	396/432 (91%)	0.67	55 (13%) 4 6	19, 35, 62, 97	0

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	140	ARG	5.4
1	A	74	GLY	5.1
1	B	188	VAL	5.0
1	B	192	ALA	4.7
1	A	73	GLY	4.4
1	A	72	PHE	4.4
1	B	141	LYS	4.0
1	A	188	VAL	3.9
1	A	41	ALA	3.8
1	A	81	ILE	3.8
1	B	204	ALA	3.7
1	B	199	HIS	3.7
1	A	39	TYR	3.5
1	B	191	PRO	3.4
1	A	141	LYS	3.3
1	B	9	VAL	3.2
1	A	114	ASN	3.1
1	B	137	VAL	3.0
1	A	13	LEU	3.0
1	A	75	LEU	3.0
1	A	143	ASN	2.9
1	B	134	LYS	2.8
1	B	72	PHE	2.8
1	B	203	VAL	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	149	ILE	2.7
1	B	50	LEU	2.7
1	A	15	LEU	2.7
1	A	90	PHE	2.7
1	A	106	ARG	2.6
1	A	140	ARG	2.6
1	B	95	ARG	2.6
1	A	96	VAL	2.6
1	A	50	LEU	2.6
1	A	16	VAL	2.6
1	A	95	ARG	2.5
1	A	63	VAL	2.4
1	A	8	GLN	2.4
1	B	88	ILE	2.4
1	A	142	LYS	2.3
1	A	9	VAL	2.3
1	B	138	PHE	2.3
1	A	87	ILE	2.3
1	A	88	ILE	2.3
1	A	40	VAL	2.2
1	A	56	ARG	2.2
1	A	92	VAL	2.2
1	A	161	PHE	2.2
1	A	61	PHE	2.1
1	A	205	GLN	2.1
1	B	87	ILE	2.1
1	A	89	MET	2.1
1	A	82	GLN	2.1
1	B	92	VAL	2.1
1	A	76	ARG	2.1
1	A	110	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	GDP	B	301	28/28	0.98	0.08	-0.61	22,28,33,35	0
2	GDP	A	301	28/28	0.98	0.07	-0.80	24,30,36,40	0
3	MG	A	302	1/1	0.98	0.04	-1.58	29,29,29,29	0
3	MG	B	302	1/1	0.98	0.05	-2.18	25,25,25,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.