



Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2016 – 10:25 PM EDT

PDB ID : 5CKR
Title : Crystal Structure of MraY in complex with Muraymycin D2
Authors : Lee, S.Y.; Chung, B.C.; Mashalidis, E.H.; Tanino, T.; Kim, M.; Hong, J.;
Ichikawa, S.
Deposited on : 2015-07-15
Resolution : 2.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027674
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027674


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2184 (3.00-2.92)
Clashscore	102246	2552 (3.00-2.92)
Ramachandran outliers	100387	2468 (3.00-2.92)
Sidechain outliers	100360	2471 (3.00-2.92)
RSRZ outliers	91569	2201 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	365	<div> <div>10%</div> <div>63%</div> <div>26%</div> <div>• 10%</div> </div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2638 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phospho-N-acetylmuramoyl-pentapeptide-transferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	329	2565	1738	395	424	8	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	GLY	-	expression tag	UNP O66465
A	-4	PRO	-	expression tag	UNP O66465
A	-3	ALA	-	expression tag	UNP O66465
A	-2	VAL	-	expression tag	UNP O66465
A	-1	PRO	-	expression tag	UNP O66465
A	0	ARG	-	expression tag	UNP O66465

- Molecule 2 is Muraymycin D2 (three-letter code: 57M) (formula: $C_{37}H_{61}N_{11}O_{16}$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			64	37	11	16		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	9	Total	O	0	0
			9	9		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phospho-N-acetylmuramoyl-pentapeptide-transferase



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	94.48Å 102.05Å 135.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	51.02 – 2.95 69.33 – 2.95	Depositor EDS
% Data completeness (in resolution range)	99.3 (51.02-2.95) 99.4 (69.33-2.95)	Depositor EDS
R_{merge}	0.27	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 2.96Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.247 , 0.261 0.247 , 0.261	Depositor DCC
R_{free} test set	702 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	67.4	Xtrriage
Anisotropy	0.040	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 76.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	2638	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 57M

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/2633	0.55	1/3587 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	329	LEU	CA-CB-CG	5.41	127.75	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2565	0	2673	94	0
2	A	64	0	0	7	0
3	A	9	0	0	2	0
All	All	2638	0	2673	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:120:VAL:HG21	1:A:128:ILE:HD11	1.09	1.05
1:A:120:VAL:CG2	1:A:128:ILE:HD11	1.94	0.97
1:A:120:VAL:HG21	1:A:128:ILE:CD1	1.94	0.97
1:A:98:LYS:HA	1:A:101:TRP:HD1	1.40	0.87
1:A:193:ASP:OD1	3:A:501:HOH:O	1.96	0.83
1:A:192:THR:O	1:A:198:LEU:HD23	1.78	0.82
1:A:197:GLY:O	1:A:342:TRP:NE1	2.16	0.78
1:A:322:PRO:HD2	1:A:325:HIS:HD2	1.50	0.77
1:A:148:ILE:O	1:A:152:ALA:O	2.07	0.72
1:A:213:VAL:HG22	1:A:356:LEU:HD21	1.71	0.71
1:A:195:LEU:HB2	1:A:198:LEU:HD22	1.73	0.70
1:A:117:ASP:HA	1:A:128:ILE:HG12	1.73	0.69
1:A:352:ALA:HA	1:A:355:MET:HG3	1.73	0.69
1:A:97:ILE:O	1:A:98:LYS:HG2	1.93	0.68
1:A:97:ILE:HG22	1:A:99:TYR:CD2	2.29	0.67
1:A:263:MET:HG2	1:A:267:GLY:HA3	1.77	0.67
1:A:98:LYS:HA	1:A:101:TRP:CD1	2.26	0.67
1:A:192:THR:HG1	1:A:295:PHE:HD1	1.43	0.66
1:A:191:LEU:HD21	1:A:322:PRO:HG3	1.78	0.65
1:A:116:TRP:CE3	1:A:128:ILE:HG13	2.31	0.65
1:A:322:PRO:HD2	1:A:325:HIS:CD2	2.32	0.65
1:A:93:MET:SD	1:A:100:THR:HG21	2.38	0.64
1:A:75:THR:HG23	1:A:76:MET:HG3	1.80	0.64
1:A:55:PHE:HE2	1:A:123:LYS:HG2	1.61	0.63
1:A:120:VAL:HG11	1:A:128:ILE:HD13	1.81	0.63
1:A:22:ILE:HD11	1:A:235:GLY:HA3	1.82	0.62
1:A:321:ALA:O	2:A:401:57M:N47	2.33	0.62
1:A:263:MET:HB2	3:A:503:HOH:O	2.00	0.60
1:A:194:GLY:HA2	2:A:401:57M:C34	2.31	0.60
1:A:97:ILE:CG2	1:A:99:TYR:CD2	2.85	0.58
1:A:75:THR:HA	1:A:262:PHE:HB2	1.86	0.58
1:A:119:TYR:CE2	1:A:123:LYS:HD2	2.39	0.58
1:A:22:ILE:HG23	1:A:23:THR:HG23	1.88	0.56
1:A:264:GLY:N	2:A:401:57M:O23	2.29	0.56
1:A:191:LEU:HD21	1:A:322:PRO:CG	2.36	0.55
1:A:191:LEU:C	1:A:191:LEU:HD23	2.26	0.55
1:A:45:ILE:HG12	1:A:261:MET:HE3	1.89	0.55
1:A:93:MET:SD	1:A:100:THR:CG2	2.94	0.55
1:A:98:LYS:O	1:A:102:VAL:HG23	2.07	0.54
1:A:98:LYS:HB2	1:A:151:TRP:CH2	2.43	0.54
1:A:319:LYS:O	1:A:320:ARG:NH2	2.42	0.53
1:A:118:ASP:OD2	1:A:265:ASP:N	2.41	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:196:ASP:OD1	2:A:401:57M:N30	2.42	0.52
1:A:159:TYR:CZ	1:A:229:PRO:HB3	2.44	0.52
1:A:197:GLY:HA2	1:A:200:ILE:HG22	1.92	0.52
1:A:250:GLY:O	1:A:253:TRP:HB3	2.10	0.51
1:A:194:GLY:HA2	2:A:401:57M:O26	2.11	0.51
1:A:116:TRP:CZ3	1:A:128:ILE:HD12	2.45	0.51
1:A:192:THR:O	1:A:198:LEU:CD2	2.55	0.50
1:A:192:THR:OG1	1:A:295:PHE:HD1	1.93	0.50
1:A:55:PHE:CE2	1:A:123:LYS:HG2	2.45	0.50
1:A:194:GLY:HA2	2:A:401:57M:N28	2.26	0.50
1:A:191:LEU:HD23	1:A:191:LEU:O	2.12	0.50
1:A:115:PHE:CD2	1:A:115:PHE:C	2.85	0.49
1:A:93:MET:CE	1:A:100:THR:HB	2.42	0.48
1:A:121:LYS:HA	1:A:126:LYS:O	2.14	0.48
1:A:284:SER:HB2	1:A:287:ILE:HB	1.94	0.47
1:A:351:ILE:O	1:A:354:SER:HB2	2.15	0.46
1:A:100:THR:HA	1:A:103:VAL:HG12	1.97	0.46
1:A:100:THR:O	1:A:103:VAL:HG12	2.16	0.45
1:A:287:ILE:HG12	1:A:355:MET:HE2	1.98	0.45
1:A:121:LYS:O	1:A:125:LYS:N	2.47	0.45
1:A:71:LYS:HE3	1:A:71:LYS:HB3	1.58	0.45
1:A:305:GLN:NE2	2:A:401:57M:O56	2.37	0.45
1:A:306:ILE:O	1:A:310:ARG:HG2	2.17	0.44
1:A:202:PRO:HB2	1:A:295:PHE:CZ	2.53	0.44
1:A:195:LEU:HD22	1:A:334:GLU:HG3	2.00	0.44
1:A:116:TRP:O	1:A:120:VAL:HG23	2.18	0.44
1:A:319:LYS:HG3	1:A:326:HIS:HD2	1.82	0.43
1:A:157:ILE:HD12	1:A:167:TYR:CZ	2.53	0.43
1:A:235:GLY:O	1:A:238:THR:HB	2.19	0.43
1:A:41:SER:O	1:A:45:ILE:HG13	2.19	0.43
1:A:139:LEU:HA	1:A:139:LEU:HD23	1.87	0.43
1:A:170:LEU:HD12	1:A:174:TYR:CD1	2.54	0.43
1:A:22:ILE:O	1:A:25:ARG:HB3	2.19	0.42
1:A:218:GLY:HA2	1:A:231:VAL:O	2.20	0.42
1:A:353:ILE:HA	1:A:356:LEU:HD23	2.00	0.42
1:A:160:PHE:HB2	1:A:163:PHE:O	2.19	0.42
1:A:195:LEU:HB2	1:A:198:LEU:CD2	2.45	0.42
1:A:141:ALA:O	1:A:145:SER:OG	2.32	0.42
1:A:209:ALA:O	1:A:213:VAL:HG23	2.20	0.41
1:A:115:PHE:HA	1:A:266:VAL:HG22	2.01	0.41
1:A:197:GLY:O	1:A:342:TRP:CE2	2.73	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:191:LEU:CD2	1:A:322:PRO:HG3	2.47	0.41
1:A:287:ILE:HA	1:A:355:MET:HE2	2.02	0.41
1:A:177:PHE:O	1:A:181:VAL:HG23	2.20	0.41
1:A:349:ALA:O	1:A:353:ILE:HG13	2.20	0.41
1:A:104:LEU:HA	1:A:104:LEU:HD23	1.84	0.41
1:A:97:ILE:HG13	1:A:236:GLU:HG3	2.03	0.41
1:A:38:LEU:HA	1:A:38:LEU:HD12	1.81	0.41
1:A:97:ILE:CG2	1:A:99:TYR:HD2	2.32	0.41
1:A:299:THR:O	1:A:303:ILE:HG13	2.21	0.41
1:A:309:PHE:CE2	1:A:314:GLY:HA2	2.57	0.40
1:A:98:LYS:HD2	1:A:151:TRP:CZ2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	325/365 (89%)	322 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	271/309 (88%)	268 (99%)	3 (1%)	80 93

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	174	TYR
1	A	329	LEU
1	A	356	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	325	HIS
1	A	326	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	57M	A	401	-	56,67,67	3.38	20 (35%)	66,95,95	1.85	14 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	57M	A	401	-	-	1/54/108/108	0/4/4/4

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	57M	C58-N59	-5.23	1.36	1.47
2	A	401	57M	C63-C64	-4.82	1.31	1.49
2	A	401	57M	C37-C27	-4.13	1.47	1.53
2	A	401	57M	C16-C21	-4.04	1.47	1.53
2	A	401	57M	C46-N47	-3.27	1.38	1.45
2	A	401	57M	C21-C22	-2.73	1.46	1.53
2	A	401	57M	C36-C37	-2.49	1.46	1.53
2	A	401	57M	C36-C25	-2.24	1.48	1.53
2	A	401	57M	C22-C18	-2.15	1.47	1.53
2	A	401	57M	O17-C18	2.29	1.50	1.45
2	A	401	57M	C64-N61	3.06	1.54	1.47
2	A	401	57M	C48-N50	3.52	1.45	1.35
2	A	401	57M	C48-N47	3.61	1.45	1.35
2	A	401	57M	O26-C25	4.04	1.50	1.44
2	A	401	57M	O17-C16	4.20	1.49	1.41
2	A	401	57M	C46-C58	5.13	1.61	1.54
2	A	401	57M	C06-N08	5.48	1.45	1.33
2	A	401	57M	C44-N43	5.48	1.46	1.34
2	A	401	57M	O26-C27	6.58	1.50	1.41
2	A	401	57M	C60-N61	16.78	1.54	1.29

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	57M	C63-C58-C46	-3.82	108.97	112.77
2	A	401	57M	C09-N08-C06	-3.07	116.47	122.62
2	A	401	57M	O07-C06-N08	-2.74	117.60	123.04
2	A	401	57M	C55-C51-N50	-2.24	109.11	113.40
2	A	401	57M	C34-N28-C29	-2.14	117.83	121.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	57M	N62-C60-N59	2.21	121.92	116.67
2	A	401	57M	C10-C09-N08	2.33	118.89	112.29
2	A	401	57M	C16-O15-C14	2.45	124.51	118.00
2	A	401	57M	C53-C52-C51	2.58	113.66	111.30
2	A	401	57M	O26-C27-N28	2.97	113.75	108.10
2	A	401	57M	C54-C52-C51	3.34	114.35	111.30
2	A	401	57M	C05-C06-N08	3.52	123.86	116.66
2	A	401	57M	N47-C48-N50	4.77	120.94	115.15
2	A	401	57M	C31-N30-C29	6.87	121.44	114.21

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	57M	N47-C46-C58-N59

There are no ring outliers.

1 monomer is involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	57M	7	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	329/365 (90%)	0.90	38 (11%) 6 3	28, 58, 106, 148	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	56	GLY	8.2
1	A	124	ASN	4.4
1	A	133	LYS	4.4
1	A	52	GLN	4.2
1	A	55	PHE	4.2
1	A	128	ILE	4.2
1	A	123	LYS	4.0
1	A	359	ARG	3.8
1	A	57	GLY	3.7
1	A	129	SER	3.7
1	A	137	GLN	3.4
1	A	122	LEU	3.4
1	A	100	THR	3.1
1	A	135	LEU	3.1
1	A	131	LYS	2.9
1	A	119	TYR	2.9
1	A	44	PHE	2.8
1	A	224	GLN	2.6
1	A	132	THR	2.6
1	A	141	ALA	2.6
1	A	315	LYS	2.5
1	A	311	TRP	2.5
1	A	48	LEU	2.4
1	A	233	TYR	2.4
1	A	236	GLU	2.4
1	A	74	PRO	2.4
1	A	168	VAL	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	223	ALA	2.3
1	A	344	ILE	2.3
1	A	167	TYR	2.2
1	A	166	LEU	2.1
1	A	332	LEU	2.1
1	A	69	VAL	2.1
1	A	158	LEU	2.1
1	A	337	ILE	2.1
1	A	120	VAL	2.0
1	A	229	PRO	2.0
1	A	157	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	57M	A	401	64/64	0.92	0.24	-0.04	43,61,80,98	0

6.5 Other polymers [i](#)

There are no such residues in this entry.