



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 01:22 AM GMT

PDB ID : 2CSB
Title : Crystal structure of Topoisomerase V from Methanopyrus kandleri (61 kDa fragment)
Authors : Taneja, B.; Patel, A.; Slesarev, A.; Mondragon, A.
Deposited on : 2005-05-21
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	519	
1	B	519	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 9026 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Topoisomerase V.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	517	Total	C	N	O	S	Se	0	12	0
			4265	2667	768	819	5	6			
1	B	517	Total	C	N	O	S	Se	0	8	0
			4244	2655	765	813	5	6			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	GB 20094872
A	40	MSE	MET	MODIFIED RESIDUE	GB 20094872
A	155	MSE	MET	MODIFIED RESIDUE	GB 20094872
A	166	MSE	MET	MODIFIED RESIDUE	GB 20094872
A	286	MSE	MET	MODIFIED RESIDUE	GB 20094872
A	325	MSE	MET	MODIFIED RESIDUE	GB 20094872
A	391	MSE	MET	MODIFIED RESIDUE	GB 20094872
B	1	MSE	MET	MODIFIED RESIDUE	GB 20094872
B	40	MSE	MET	MODIFIED RESIDUE	GB 20094872
B	155	MSE	MET	MODIFIED RESIDUE	GB 20094872
B	166	MSE	MET	MODIFIED RESIDUE	GB 20094872
B	286	MSE	MET	MODIFIED RESIDUE	GB 20094872
B	325	MSE	MET	MODIFIED RESIDUE	GB 20094872
B	391	MSE	MET	MODIFIED RESIDUE	GB 20094872

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	277	Total 277	O 277	0	0
3	B	239	Total 239	O 239	0	0


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Topoisomerase V



• Molecule 1: Topoisomerase V





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	70.69 Å 89.77 Å 189.07 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.44 – 2.30 27.77 – 2.25	Depositor EDS
% Data completeness (in resolution range)	97.9 (26.44-2.30) 100.0 (27.77-2.25)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1841.48 (at 2.24 Å)	Xtriage
Refinement program	REFMAC 5.2.0003	Depositor
R, R_{free}	0.217 , 0.292 0.231 , 0.303	Depositor DCC
R_{free} test set	2742 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	33.9	Xtriage
Anisotropy	0.208	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 57919 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	9026	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	0/4375	0.94	14/5884 (0.2%)
1	B	0.84	4/4338 (0.1%)	0.92	14/5835 (0.2%)
All	All	0.83	4/8713 (0.0%)	0.93	28/11719 (0.2%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	453[A]	GLU	CD-OE2	14.97	1.42	1.25
1	B	453[B]	GLU	CD-OE2	14.97	1.42	1.25
1	B	453[A]	GLU	CD-OE1	6.01	1.32	1.25
1	B	453[B]	GLU	CD-OE1	6.01	1.32	1.25

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	102	ARG	NE-CZ-NH2	-7.79	116.41	120.30
1	A	182	ASP	CB-CG-OD2	7.64	125.17	118.30
1	B	368	ASP	CB-CG-OD2	7.49	125.04	118.30
1	A	124	ASP	CB-CG-OD2	7.36	124.92	118.30
1	A	219	ASP	CB-CG-OD2	7.24	124.82	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4265	0	4271	74	0
1	B	4244	0	4258	104	0
2	A	1	0	0	0	0
3	A	277	0	0	10	0
3	B	239	0	0	12	0
All	All	9026	0	8529	177	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 177 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:90:ARG:HG3	1:B:93[B]:ARG:NH1	1.53	1.22
1:A:388:ILE:HA	1:A:391:MSE:HE3	1.45	0.98
1:B:90:ARG:CG	1:B:93[B]:ARG:NH1	2.28	0.96
1:B:90:ARG:HG3	1:B:93[B]:ARG:HH12	1.32	0.93
1:B:61:LEU:CD2	1:B:155:MSE:HE1	1.99	0.91

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	527/519 (102%)	514 (98%)	13 (2%)	0	100	100
1	B	523/519 (101%)	501 (96%)	20 (4%)	2 (0%)	39	48
All	All	1050/1038 (101%)	1015 (97%)	33 (3%)	2 (0%)	52	64

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	336	ASP
1	B	364	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	460/442 (104%)	440 (96%)	20 (4%)	35	47
1	B	456/442 (103%)	421 (92%)	35 (8%)	16	20
All	All	916/884 (104%)	861 (94%)	55 (6%)	24	31

5 of 55 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	148	ARG
1	B	270	ARG
1	B	480	ARG
1	B	155	MSE
1	B	205	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	ASN
1	A	235	ASN
1	A	373	HIS
1	B	78	GLN
1	B	511	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	511/519 (98%)	0.78	53 (10%) 8 12	22, 35, 47, 67	0
1	B	511/519 (98%)	1.22	99 (19%) 1 2	27, 42, 57, 66	0
All	All	1022/1038 (98%)	1.00	152 (14%) 3 5	22, 38, 54, 67	0

The worst 5 of 152 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	208	GLY	8.9
1	B	450	ALA	8.8
1	B	14	ARG	7.2
1	B	48	ASN	6.3
1	B	459	VAL	6.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	MG	A	3052	1/1	0.96	0.16	-	30,30,30,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.