



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 08:01 AM GMT

PDB ID : 3CSY
Title : Crystal structure of the trimeric prefusion Ebola virus glycoprotein in complex with a neutralizing antibody from a human survivor
Authors : Lee, J.E.; Fusco, M.L.; Hessel, A.J.; Oswald, W.B.; Burton, D.R.; Saphire, E.O.
Deposited on : 2008-04-10
Resolution : 3.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1611 (3.50-3.30)
Ramachandran outliers	100387	1571 (3.50-3.30)
Sidechain outliers	100360	1571 (3.50-3.30)
RSRZ outliers	91569	1485 (3.50-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	226	
1	C	226	
1	E	226	
1	G	226	
2	B	217	
2	D	217	
2	F	217	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	217	
3	I	334	
3	K	334	
3	M	334	
3	O	334	
4	J	131	
4	L	131	
4	N	131	
4	P	131	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	NAG	K	351	-	-	-	X

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 23539 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab KZ52 heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	226	Total	C	N	O	S	Se	0	0	0
			1687	1059	286	334	4	4			
1	C	226	Total	C	N	O	S	Se	0	0	0
			1687	1059	286	334	4	4			
1	E	226	Total	C	N	O	S	Se	0	0	0
			1687	1059	286	334	4	4			
1	G	226	Total	C	N	O	S	Se	0	0	0
			1687	1059	286	334	4	4			

- Molecule 2 is a protein called Fab KZ52 light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	217	Total	C	N	O	S	Se	0	0	0
			1682	1056	281	340	4	1			
2	D	217	Total	C	N	O	S	Se	0	0	0
			1682	1056	281	340	4	1			
2	F	217	Total	C	N	O	S	Se	0	0	0
			1682	1056	281	340	4	1			
2	H	217	Total	C	N	O	S	Se	0	0	0
			1682	1056	281	340	4	1			

- Molecule 3 is a protein called Envelope glycoprotein GP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	I	236	Total	C	N	O	S	0	0	0
			1707	1085	293	325	4			
3	K	232	Total	C	N	O	S	0	0	0
			1687	1073	289	321	4			
3	M	230	Total	C	N	O	S	0	0	0
			1677	1067	287	319	4			
3	O	225	Total	C	N	O	S	0	0	0
			1651	1052	282	313	4			

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	16	TYR	-	EXPRESSION TAG	UNP Q05320
I	17	PRO	-	EXPRESSION TAG	UNP Q05320
I	18	TYR	-	EXPRESSION TAG	UNP Q05320
I	19	ASP	-	EXPRESSION TAG	UNP Q05320
I	20	VAL	-	EXPRESSION TAG	UNP Q05320
I	21	PRO	-	EXPRESSION TAG	UNP Q05320
I	22	ASP	-	EXPRESSION TAG	UNP Q05320
I	23	TYR	-	EXPRESSION TAG	UNP Q05320
I	24	ALA	-	EXPRESSION TAG	UNP Q05320
I	25	ILE	-	EXPRESSION TAG	UNP Q05320
I	26	GLU	-	EXPRESSION TAG	UNP Q05320
I	27	GLY	-	EXPRESSION TAG	UNP Q05320
I	28	ARG	-	EXPRESSION TAG	UNP Q05320
I	29	GLY	-	EXPRESSION TAG	UNP Q05320
I	30	ALA	-	EXPRESSION TAG	UNP Q05320
I	31	ARG	-	EXPRESSION TAG	UNP Q05320
I	42	VAL	THR	ENGINEERED	UNP Q05320
I	230	VAL	THR	ENGINEERED	UNP Q05320
K	16	TYR	-	EXPRESSION TAG	UNP Q05320
K	17	PRO	-	EXPRESSION TAG	UNP Q05320
K	18	TYR	-	EXPRESSION TAG	UNP Q05320
K	19	ASP	-	EXPRESSION TAG	UNP Q05320
K	20	VAL	-	EXPRESSION TAG	UNP Q05320
K	21	PRO	-	EXPRESSION TAG	UNP Q05320
K	22	ASP	-	EXPRESSION TAG	UNP Q05320
K	23	TYR	-	EXPRESSION TAG	UNP Q05320
K	24	ALA	-	EXPRESSION TAG	UNP Q05320
K	25	ILE	-	EXPRESSION TAG	UNP Q05320
K	26	GLU	-	EXPRESSION TAG	UNP Q05320
K	27	GLY	-	EXPRESSION TAG	UNP Q05320
K	28	ARG	-	EXPRESSION TAG	UNP Q05320
K	29	GLY	-	EXPRESSION TAG	UNP Q05320
K	30	ALA	-	EXPRESSION TAG	UNP Q05320
K	31	ARG	-	EXPRESSION TAG	UNP Q05320
K	42	VAL	THR	ENGINEERED	UNP Q05320
K	230	VAL	THR	ENGINEERED	UNP Q05320
M	16	TYR	-	EXPRESSION TAG	UNP Q05320
M	17	PRO	-	EXPRESSION TAG	UNP Q05320
M	18	TYR	-	EXPRESSION TAG	UNP Q05320
M	19	ASP	-	EXPRESSION TAG	UNP Q05320
M	20	VAL	-	EXPRESSION TAG	UNP Q05320
M	21	PRO	-	EXPRESSION TAG	UNP Q05320

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
M	22	ASP	-	EXPRESSION TAG	UNP Q05320
M	23	TYR	-	EXPRESSION TAG	UNP Q05320
M	24	ALA	-	EXPRESSION TAG	UNP Q05320
M	25	ILE	-	EXPRESSION TAG	UNP Q05320
M	26	GLU	-	EXPRESSION TAG	UNP Q05320
M	27	GLY	-	EXPRESSION TAG	UNP Q05320
M	28	ARG	-	EXPRESSION TAG	UNP Q05320
M	29	GLY	-	EXPRESSION TAG	UNP Q05320
M	30	ALA	-	EXPRESSION TAG	UNP Q05320
M	31	ARG	-	EXPRESSION TAG	UNP Q05320
M	42	VAL	THR	ENGINEERED	UNP Q05320
M	230	VAL	THR	ENGINEERED	UNP Q05320
O	16	TYR	-	EXPRESSION TAG	UNP Q05320
O	17	PRO	-	EXPRESSION TAG	UNP Q05320
O	18	TYR	-	EXPRESSION TAG	UNP Q05320
O	19	ASP	-	EXPRESSION TAG	UNP Q05320
O	20	VAL	-	EXPRESSION TAG	UNP Q05320
O	21	PRO	-	EXPRESSION TAG	UNP Q05320
O	22	ASP	-	EXPRESSION TAG	UNP Q05320
O	23	TYR	-	EXPRESSION TAG	UNP Q05320
O	24	ALA	-	EXPRESSION TAG	UNP Q05320
O	25	ILE	-	EXPRESSION TAG	UNP Q05320
O	26	GLU	-	EXPRESSION TAG	UNP Q05320
O	27	GLY	-	EXPRESSION TAG	UNP Q05320
O	28	ARG	-	EXPRESSION TAG	UNP Q05320
O	29	GLY	-	EXPRESSION TAG	UNP Q05320
O	30	ALA	-	EXPRESSION TAG	UNP Q05320
O	31	ARG	-	EXPRESSION TAG	UNP Q05320
O	42	VAL	THR	ENGINEERED	UNP Q05320
O	230	VAL	THR	ENGINEERED	UNP Q05320

- Molecule 4 is a protein called Envelope glycoprotein GP2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	J	98	Total	C	N	O	S	0	0	0
			746	477	130	136	3			
4	L	93	Total	C	N	O	S	0	0	0
			727	466	126	132	3			
4	N	95	Total	C	N	O	S	0	0	0
			732	469	128	132	3			
4	P	93	Total	C	N	O	S	0	0	0
			719	462	125	129	3			

- Molecule 5 is a polymer of unknown type called SUGAR (3-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	I	3	Total	C	N	O	0	0
			39	22	2	15		
5	M	3	Total	C	N	O	0	0
			39	22	2	15		
5	O	3	Total	C	N	O	0	0
			39	22	2	15		

- Molecule 6 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	J	5	Total	C	N	O	0	0
			61	34	2	25		
6	N	5	Total	C	N	O	0	0
			61	34	2	25		

- Molecule 7 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	K	2	Total	C	N	O	0	0
			28	16	2	10		


- Molecule 8 is a polymer of unknown type called SUGAR (6-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	L	6	Total	C	N	O	0	0
			75	42	3	30		
8	P	6	Total	C	N	O	0	0
			75	42	3	30		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fab KZ52 heavy chain




- Molecule 1: Fab KZ52 heavy chain




- Molecule 1: Fab KZ52 heavy chain








• Molecule 1: Fab KZ52 heavy chain



• Molecule 2: Fab KZ52 light chain




• Molecule 2: Fab KZ52 light chain




• Molecule 2: Fab KZ52 light chain








• Molecule 2: Fab KZ52 light chain




• Molecule 3: Envelope glycoprotein GP1




• Molecule 3: Envelope glycoprotein GP1







• Molecule 3: Envelope glycoprotein GP1




• Molecule 3: Envelope glycoprotein GP1








• Molecule 4: Envelope glycoprotein GP2




• Molecule 4: Envelope glycoprotein GP2



• Molecule 4: Envelope glycoprotein GP2



• Molecule 4: Envelope glycoprotein GP2



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	273.71 Å 273.71 Å 409.43 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.37 – 3.40 48.37 – 3.40	Depositor EDS
% Data completeness (in resolution range)	93.4 (48.37-3.40) 96.9 (48.37-3.40)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.67 (at 3.40 Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.261 , 0.302 (Not available) , (Not available)	Depositor DCC
R_{free} test set	NotAvailable	DCC
Wilson B-factor (Å ²)	95.5	Xtriage
Anisotropy	0.075	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 168.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 151305 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	23539	wwPDB-VP
Average B, all atoms (Å ²)	121.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.53	0/1721	0.56	0/2333
1	C	0.49	1/1721 (0.1%)	0.50	0/2333
1	E	0.49	1/1721 (0.1%)	0.52	0/2333
1	G	0.48	0/1721	0.52	0/2333
2	B	0.41	0/1718	0.50	0/2331
2	D	0.37	0/1718	0.49	0/2331
2	F	0.42	0/1718	0.49	0/2331
2	H	0.37	0/1718	0.48	0/2331
3	I	0.49	0/1743	0.57	0/2379
3	K	0.44	0/1723	0.54	0/2351
3	M	0.47	0/1712	0.57	1/2334 (0.0%)
3	O	0.41	0/1686	0.53	0/2298
4	J	0.54	0/762	0.60	0/1038
4	L	0.47	0/742	0.56	0/1010
4	N	0.50	0/747	0.61	0/1016
4	P	0.49	0/734	0.61	0/999
All	All	0.46	2/23605 (0.0%)	0.53	1/32081 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
3	I	0	1
3	K	0	1
4	N	0	1
All	All	0	4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	108	MSE	CG-SE	-5.33	1.77	1.95
1	E	34	MSE	CG-SE	-5.19	1.77	1.95

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	M	265	LYS	N-CA-C	5.20	125.03	111.00

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	125	ALA	Peptide
3	I	55	ASP	Peptide
3	K	54	ARG	Peptide
4	N	536	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1687	0	1660	78	0
1	C	1687	0	1660	51	0
1	E	1687	0	1660	59	0
1	G	1687	0	1660	53	0
2	B	1682	0	1638	48	0
2	D	1682	0	1638	39	0
2	F	1682	0	1638	38	0
2	H	1682	0	1638	44	0
3	I	1707	0	1548	81	0
3	K	1687	0	1540	73	0
3	M	1677	0	1535	68	0
3	O	1651	0	1520	73	0
4	J	746	0	722	41	0
4	L	727	0	712	44	0
4	N	732	0	716	39	0
4	P	719	0	706	26	0
5	I	39	0	34	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	M	39	0	34	0	0
5	O	39	0	34	6	0
6	J	61	0	52	2	0
6	N	61	0	52	3	0
7	K	28	0	25	2	0
8	L	75	0	64	3	0
8	P	75	0	64	4	0
All	All	23539	0	22550	750	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 750 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:N:563:ASN:ND2	6:N:701:NAG:C1	1.67	1.54
3:O:257:ASN:HD21	5:O:351:NAG:C1	0.89	1.53
3:I:257:ASN:ND2	5:I:351:NAG:C1	1.68	1.51
3:K:257:ASN:ND2	7:K:351:NAG:C1	1.68	1.50
4:L:563:ASN:ND2	8:L:701:NAG:C1	1.70	1.49

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	224/226 (99%)	192 (86%)	29 (13%)	3 (1%)	15	57
1	C	224/226 (99%)	194 (87%)	27 (12%)	3 (1%)	15	57
1	E	224/226 (99%)	196 (88%)	26 (12%)	2 (1%)	21	65
1	G	224/226 (99%)	202 (90%)	21 (9%)	1 (0%)	39	79

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	215/217 (99%)	181 (84%)	25 (12%)	9 (4%)	3	29
2	D	215/217 (99%)	183 (85%)	25 (12%)	7 (3%)	5	37
2	F	215/217 (99%)	183 (85%)	26 (12%)	6 (3%)	6	41
2	H	215/217 (99%)	179 (83%)	27 (13%)	9 (4%)	3	29
3	I	230/334 (69%)	184 (80%)	33 (14%)	13 (6%)	2	20
3	K	226/334 (68%)	184 (81%)	33 (15%)	9 (4%)	4	31
3	M	222/334 (66%)	181 (82%)	29 (13%)	12 (5%)	2	22
3	O	217/334 (65%)	171 (79%)	26 (12%)	20 (9%)	1	9
4	J	96/131 (73%)	80 (83%)	14 (15%)	2 (2%)	9	47
4	L	89/131 (68%)	79 (89%)	8 (9%)	2 (2%)	8	46
4	N	91/131 (70%)	80 (88%)	10 (11%)	1 (1%)	17	61
4	P	89/131 (68%)	74 (83%)	13 (15%)	2 (2%)	8	46
All	All	3016/3632 (83%)	2543 (84%)	372 (12%)	101 (3%)	5	37

5 of 101 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	100(B)	TYR
2	B	110	VAL
2	D	27(F)	SER
2	D	78	LEU
2	D	169	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	190/186 (102%)	175 (92%)	15 (8%)	15	52
1	C	190/186 (102%)	177 (93%)	13 (7%)	20	60
1	E	190/186 (102%)	179 (94%)	11 (6%)	25	65
1	G	190/186 (102%)	175 (92%)	15 (8%)	15	52

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	191/190 (100%)	185 (97%)	6 (3%)	47	81
2	D	191/190 (100%)	188 (98%)	3 (2%)	70	89
2	F	191/190 (100%)	187 (98%)	4 (2%)	61	86
2	H	191/190 (100%)	186 (97%)	5 (3%)	54	84
3	I	163/282 (58%)	156 (96%)	7 (4%)	35	74
3	K	163/282 (58%)	154 (94%)	9 (6%)	27	66
3	M	163/282 (58%)	153 (94%)	10 (6%)	23	63
3	O	162/282 (57%)	152 (94%)	10 (6%)	23	63
4	J	73/110 (66%)	69 (94%)	4 (6%)	27	66
4	L	74/110 (67%)	71 (96%)	3 (4%)	37	75
4	N	73/110 (66%)	70 (96%)	3 (4%)	37	75
4	P	72/110 (66%)	69 (96%)	3 (4%)	36	74
All	All	2467/3072 (80%)	2346 (95%)	121 (5%)	31	70

5 of 121 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	18	LEU
2	H	27(F)	SER
3	O	81	SER
1	G	34	MSE
1	G	75	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	155	GLN
3	I	110	ASN
4	J	595	GLN
3	O	257	ASN
4	P	563	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

33 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	NAG	I	351	5	14,14,15	0.73	0	15,19,21	2.23	3 (20%)
5	NAG	I	352	5	14,14,15	0.80	0	15,19,21	1.12	1 (6%)
5	BMA	I	353	5	11,11,12	1.41	1 (9%)	14,15,17	2.11	4 (28%)
6	NAG	J	701	6	14,14,15	0.93	1 (7%)	15,19,21	1.20	2 (13%)
6	NAG	J	702	6	14,14,15	0.70	0	15,19,21	2.05	4 (26%)
6	BMA	J	703	6	11,11,12	1.12	1 (9%)	14,15,17	2.25	2 (14%)
6	MAN	J	704	6	11,11,12	0.77	0	14,15,17	1.65	3 (21%)
6	MAN	J	705	6	11,11,12	0.68	0	14,15,17	1.57	2 (14%)
7	NAG	K	351	7	14,14,15	0.75	0	15,19,21	1.80	2 (13%)
7	NAG	K	352	7	14,14,15	0.65	0	15,19,21	1.50	2 (13%)
8	NAG	L	701	8	14,14,15	0.99	1 (7%)	15,19,21	2.45	4 (26%)
8	NAG	L	702	8	14,14,15	0.52	0	15,19,21	1.63	2 (13%)
8	BMA	L	703	8	11,11,12	1.38	2 (18%)	14,15,17	1.92	3 (21%)
8	MAN	L	704	8	11,11,12	0.98	1 (9%)	14,15,17	1.44	2 (14%)
8	MAN	L	705	8	11,11,12	0.84	1 (9%)	14,15,17	1.73	3 (21%)
8	NAG	L	706	8	14,14,15	0.68	0	15,19,21	1.61	2 (13%)
5	NAG	M	351	5	14,14,15	0.87	0	15,19,21	2.34	3 (20%)
5	NAG	M	352	5	14,14,15	0.56	0	15,19,21	1.54	3 (20%)
5	BMA	M	353	5	11,11,12	1.13	0	14,15,17	1.27	2 (14%)
6	NAG	N	701	6	14,14,15	0.89	1 (7%)	15,19,21	1.91	2 (13%)
6	NAG	N	702	6	14,14,15	0.97	1 (7%)	15,19,21	1.52	4 (26%)
6	BMA	N	703	6	11,11,12	1.20	1 (9%)	14,15,17	2.02	1 (7%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	MAN	N	704	6	11,11,12	0.86	1 (9%)	14,15,17	1.52	2 (14%)
6	MAN	N	705	6	11,11,12	0.52	0	14,15,17	1.25	1 (7%)
5	NAG	O	351	5	14,14,15	0.72	0	15,19,21	2.21	2 (13%)
5	NAG	O	352	5	14,14,15	0.86	0	15,19,21	1.35	2 (13%)
5	BMA	O	353	5	11,11,12	1.66	2 (18%)	14,15,17	1.84	1 (7%)
8	NAG	P	701	8	14,14,15	0.90	1 (7%)	15,19,21	1.94	5 (33%)
8	NAG	P	702	8	14,14,15	0.70	0	15,19,21	1.73	3 (20%)
8	BMA	P	703	8	11,11,12	1.21	1 (9%)	14,15,17	1.64	3 (21%)
8	MAN	P	704	8	11,11,12	1.17	2 (18%)	14,15,17	1.28	2 (14%)
8	MAN	P	705	8	11,11,12	0.65	0	14,15,17	1.28	2 (14%)
8	NAG	P	706	8	14,14,15	0.59	0	15,19,21	1.97	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	I	351	5	-	0/6/23/26	0/1/1/1
5	NAG	I	352	5	-	0/6/23/26	0/1/1/1
5	BMA	I	353	5	-	0/2/19/22	0/1/1/1
6	NAG	J	701	6	-	0/6/23/26	0/1/1/1
6	NAG	J	702	6	-	0/6/23/26	0/1/1/1
6	BMA	J	703	6	-	0/2/19/22	0/1/1/1
6	MAN	J	704	6	-	0/2/19/22	0/1/1/1
6	MAN	J	705	6	-	0/2/19/22	0/1/1/1
7	NAG	K	351	7	-	0/6/23/26	0/1/1/1
7	NAG	K	352	7	-	0/6/23/26	0/1/1/1
8	NAG	L	701	8	-	0/6/23/26	0/1/1/1
8	NAG	L	702	8	-	0/6/23/26	0/1/1/1
8	BMA	L	703	8	-	0/2/19/22	0/1/1/1
8	MAN	L	704	8	-	0/2/19/22	0/1/1/1
8	MAN	L	705	8	-	0/2/19/22	0/1/1/1
8	NAG	L	706	8	-	0/6/23/26	0/1/1/1
5	NAG	M	351	5	-	0/6/23/26	0/1/1/1
5	NAG	M	352	5	-	0/6/23/26	0/1/1/1
5	BMA	M	353	5	-	0/2/19/22	0/1/1/1
6	NAG	N	701	6	-	0/6/23/26	0/1/1/1
6	NAG	N	702	6	-	0/6/23/26	0/1/1/1
6	BMA	N	703	6	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	MAN	N	704	6	-	0/2/19/22	0/1/1/1
6	MAN	N	705	6	-	0/2/19/22	0/1/1/1
5	NAG	O	351	5	-	0/6/23/26	0/1/1/1
5	NAG	O	352	5	-	1/6/23/26	0/1/1/1
5	BMA	O	353	5	-	0/2/19/22	0/1/1/1
8	NAG	P	701	8	-	0/6/23/26	0/1/1/1
8	NAG	P	702	8	-	0/6/23/26	0/1/1/1
8	BMA	P	703	8	-	0/2/19/22	0/1/1/1
8	MAN	P	704	8	-	0/2/19/22	0/1/1/1
8	MAN	P	705	8	-	0/2/19/22	0/1/1/1
8	NAG	P	706	8	-	0/6/23/26	0/1/1/1

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	L	701	NAG	O5-C1	-2.72	1.39	1.43
6	N	701	NAG	O5-C1	-2.59	1.39	1.43
8	P	704	MAN	O5-C1	-2.56	1.39	1.43
6	J	701	NAG	O5-C1	-2.49	1.39	1.43
8	P	701	NAG	O5-C1	-2.34	1.39	1.43

The worst 5 of 80 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	N	701	NAG	C2-N2-C7	-6.32	114.92	123.04
8	L	701	NAG	C2-N2-C7	-5.33	116.19	123.04
6	J	702	NAG	C2-N2-C7	-4.53	117.22	123.04
5	M	352	NAG	O4-C4-C3	-4.33	100.59	110.34
6	J	703	BMA	C1-C2-C3	-4.20	104.58	109.54

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	O	352	NAG	O7-C7-N2-C2

There are no ring outliers.

11 monomers are involved in 24 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	I	351	NAG	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	J	701	NAG	1	0
6	J	703	BMA	1	0
6	J	705	MAN	1	0
7	K	351	NAG	2	0
8	L	701	NAG	3	0
6	N	701	NAG	2	0
6	N	702	NAG	1	0
5	O	351	NAG	6	0
5	O	352	NAG	1	0
8	P	701	NAG	4	0

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	222/226 (98%)	1.27	46 (20%) 1 1	73, 105, 211, 230	0
1	C	222/226 (98%)	1.20	42 (18%) 2 2	85, 116, 174, 196	0
1	E	222/226 (98%)	1.12	39 (17%) 2 2	87, 110, 158, 187	0
1	G	222/226 (98%)	1.38	50 (22%) 1 1	90, 116, 186, 207	0
2	B	216/217 (99%)	1.74	71 (32%) 0 1	87, 137, 195, 219	0
2	D	216/217 (99%)	1.44	64 (29%) 1 1	100, 140, 162, 194	0
2	F	216/217 (99%)	1.32	54 (25%) 1 1	104, 123, 156, 176	0
2	H	216/217 (99%)	1.88	75 (34%) 0 1	101, 140, 203, 234	0
3	I	236/334 (70%)	0.86	31 (13%) 5 4	79, 104, 139, 161	0
3	K	232/334 (69%)	1.03	32 (13%) 4 4	85, 118, 157, 184	0
3	M	230/334 (68%)	0.84	23 (10%) 9 9	84, 108, 144, 172	0
3	O	225/334 (67%)	1.23	50 (22%) 1 1	94, 122, 159, 182	0
4	J	98/131 (74%)	0.82	6 (6%) 25 23	70, 98, 155, 186	0
4	L	93/131 (70%)	1.04	13 (13%) 4 4	76, 106, 138, 177	0
4	N	95/131 (72%)	1.02	10 (10%) 8 8	69, 98, 141, 183	0
4	P	93/131 (70%)	1.12	12 (12%) 5 4	80, 105, 135, 163	0
All	All	3054/3632 (84%)	1.23	618 (20%) 1 1	69, 116, 179, 234	0

The worst 5 of 618 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	127	SER	13.4
1	A	126	PRO	12.5
1	A	135	GLY	11.9
1	A	140	LEU	9.4
1	G	126	PRO	9.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	NAG	J	701	14/15	0.82	0.29	1.74	94,94,94,94	0
8	NAG	L	701	14/15	0.77	0.31	1.02	98,98,98,98	0
7	NAG	K	351	14/15	0.51	0.50	0.01	156,156,156,156	0
5	NAG	I	351	14/15	0.79	0.38	-0.09	115,115,115,115	0
5	NAG	M	351	14/15	0.78	0.31	-0.43	117,117,117,117	0
5	NAG	O	351	14/15	0.72	0.35	-0.51	140,140,140,140	0
8	NAG	P	701	14/15	0.80	0.28	-0.60	97,97,97,97	0
6	NAG	N	701	14/15	0.85	0.23	-1.25	96,96,96,96	0
5	NAG	I	352	14/15	0.77	0.30	-	175,175,175,175	0
6	MAN	J	704	11/12	0.78	0.21	-	148,148,148,148	0
5	BMA	I	353	11/12	0.54	0.36	-	172,172,172,172	0
7	NAG	K	352	14/15	0.54	0.50	-	167,167,167,167	0
6	NAG	J	702	14/15	0.88	0.29	-	105,105,105,105	0
5	NAG	O	352	14/15	0.69	0.45	-	177,177,177,177	0
8	NAG	L	702	14/15	0.85	0.24	-	126,126,126,126	0
5	NAG	M	352	14/15	0.62	0.41	-	162,162,162,162	0
8	MAN	L	705	11/12	0.54	0.47	-	162,162,162,162	0
8	MAN	P	704	11/12	0.80	0.21	-	144,144,144,144	0
8	MAN	L	704	11/12	0.76	0.25	-	160,160,160,160	0
6	MAN	N	704	11/12	0.71	0.54	-	132,132,132,132	0
6	MAN	J	705	11/12	0.74	0.31	-	171,171,171,171	0
6	NAG	N	702	14/15	0.83	0.29	-	115,115,115,115	0
8	NAG	P	702	14/15	0.88	0.20	-	116,116,116,116	0
8	MAN	P	705	11/12	0.46	0.45	-	174,174,174,174	0
8	NAG	P	706	14/15	0.76	0.45	-	177,177,177,177	0
8	BMA	L	703	11/12	0.78	0.22	-	118,118,118,118	0
5	BMA	M	353	11/12	0.40	0.49	-	168,168,168,168	0
8	NAG	L	706	14/15	0.62	0.44	-	173,173,173,173	0
6	MAN	N	705	11/12	0.79	0.23	-	148,148,148,148	0
6	BMA	J	703	11/12	0.66	0.32	-	135,135,135,135	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	BMA	O	353	11/12	0.11	0.61	-	190,190,190,190	0
8	BMA	P	703	11/12	0.75	0.21	-	154,154,154,154	0
6	BMA	N	703	11/12	0.79	0.32	-	140,140,140,140	0

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.