



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:36 PM GMT

PDB ID : 4CSY
Title : E-selectin lectin, EGF-like and two SCR domains complexed with Sialyl Lewis X
Authors : Preston, R.C.; Jakob, R.P.; Binder, F.P.C.; Sager, C.P.; Ernst, B.; Maier, T.
Deposited on : 2014-03-11
Resolution : 2.41 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3386 (2.44-2.40)
Clashscore	102246	3897 (2.44-2.40)
Ramachandran outliers	100387	3837 (2.44-2.40)
Sidechain outliers	100360	3838 (2.44-2.40)
RSRZ outliers	91569	3396 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	280	<div> <div>5%</div> <div>91%</div> <div>9%</div> </div>
1	B	280	<div> <div>4%</div> <div>90%</div> <div>10%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	B	285	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4821 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E-SELECTIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	280	Total	C	N	O	S	0	3	0
			2199	1361	363	447	28			
1	B	280	Total	C	N	O	S	0	1	0
			2183	1352	361	442	28			

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is a polymer of unknown type called SUGAR (4-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	4	Total	C	N	O	0	0
			57	32	2	23		
3	B	4	Total	C	N	O	0	0
			57	32	2	23		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ca	0	0
			1	1		
4	A	1	Total	Ca	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	61	Total	O	0	0
			61	61		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	66	Total	O	0	0
			66	66		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: E-SELECTIN



• Molecule 1: E-SELECTIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	52.29Å 58.67Å 58.87Å 76.03° 86.28° 86.31°	Depositor
Resolution (Å)	28.32 – 2.41 24.50 – 2.41	Depositor EDS
% Data completeness (in resolution range)	96.5 (28.32-2.41) 88.4 (24.50-2.41)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.13 (at 2.41Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.215 , 0.253 0.230 , 0.270	Depositor DCC
R_{free} test set	1296 reflections (5.42%)	DCC
Wilson B-factor (Å ²)	55.2	Xtriage
Anisotropy	0.580	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 44.8	EDS
Estimated twinning fraction	0.440 for -h,-l,-k	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 25216 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4821	wwPDB-VP
Average B, all atoms (Å ²)	79.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, CA, SIA, GAL, FUC, MAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/2257	0.74	0/3071
1	B	0.51	0/2241	0.71	0/3049
All	All	0.52	0/4498	0.72	0/6120

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2199	0	2009	12	0
1	B	2183	0	1998	12	0
2	A	98	0	91	1	0
2	B	98	0	91	0	0
3	A	57	0	48	0	0
3	B	57	0	48	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	61	0	0	1	0
5	B	66	0	0	0	0
All	All	4821	0	4285	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:30:GLN:HG2	1:A:62:TRP:HH2	1.46	0.81
1:A:39:ASN:HD21	1:A:96:LYS:H	1.29	0.80
1:B:39:ASN:HD21	1:B:96:LYS:H	1.29	0.79
1:A:31:ASN:OD1	1:A:33:GLU:HG2	1.87	0.73
1:B:255:PHE:HD1	1:B:279:LYS:O	1.73	0.71
1:A:263:LEU:HD22	1:A:273:ASN:HB2	1.75	0.67
1:B:263:LEU:HD22	1:B:273:ASN:HB2	1.75	0.67
1:A:154:GLU:HG3	1:B:160:THR:CG2	2.26	0.65
1:A:24:THR:HG22	1:A:119:THR:HB	1.81	0.62
1:A:30:GLN:HG2	1:A:62:TRP:CH2	2.34	0.59
1:B:24:THR:HG22	1:B:119:THR:HB	1.85	0.58
1:B:30:GLN:HG2	1:B:104:TRP:HZ2	1.68	0.57
1:B:255:PHE:CD1	1:B:279:LYS:O	2.56	0.55
1:A:154:GLU:HG3	1:B:160:THR:HG21	1.92	0.51
1:A:154:GLU:HG2	1:A:155:GLN:HG3	1.93	0.49
1:B:223:ALA:HA	1:B:241:PHE:CE2	2.50	0.47
1:B:28:ALA:O	1:B:29:ILE:O	2.33	0.47
1:B:174:HIS:HB3	1:B:177:GLY:O	2.17	0.44
1:A:174:HIS:HB3	1:A:177:GLY:O	2.17	0.43
1:A:236:GLN:HB2	5:A:2048:HOH:O	2.19	0.42
1:B:30:GLN:HE21	1:B:35:ILE:HD11	1.83	0.42
1:A:155:GLN:HB3	2:A:283:NAG:H83	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	281/280 (100%)	269 (96%)	10 (4%)	2 (1%)	26	37
1	B	279/280 (100%)	268 (96%)	9 (3%)	2 (1%)	26	37
All	All	560/560 (100%)	537 (96%)	19 (3%)	4 (1%)	26	37

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	29	ILE
1	A	139	ASN
1	B	139	ASN
1	A	29	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	250/247 (101%)	239 (96%)	11 (4%)	35	52
1	B	248/247 (100%)	239 (96%)	9 (4%)	42	62
All	All	498/494 (101%)	478 (96%)	20 (4%)	38	57

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	ILE
1	A	33	GLU
1	A	59	VAL
1	A	61	VAL
1	A	66	GLN
1	A	85	GLN
1	A	111	LYS
1	A	166	GLU
1	A	229	ASN
1	A	250	ASP
1	A	262	SER
1	B	59	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	87	ASP
1	B	111	LYS
1	B	118	TYR
1	B	166	GLU
1	B	235	PHE
1	B	243	TRP
1	B	253	GLU
1	B	262	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	30	GLN
1	A	39	ASN
1	A	236	GLN
1	A	237	ASN
1	A	270	ASN
1	B	30	GLN
1	B	39	ASN
1	B	270	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SIA	A	301	3	17,21,21	3.93	7 (41%)	19,31,31	1.29	4 (21%)
3	GAL	A	302	3	9,10,12	1.52	2 (22%)	10,13,17	0.98	1 (10%)
3	MAG	A	303	3	16,16,16	1.82	5 (31%)	21,22,22	1.36	3 (14%)
3	FUC	A	304	3,4	10,10,11	1.16	1 (10%)	14,14,16	0.98	1 (7%)
3	SIA	B	301	3	17,21,21	3.89	7 (41%)	19,31,31	1.34	3 (15%)
3	GAL	B	302	3	9,10,12	1.57	2 (22%)	10,13,17	1.03	1 (10%)
3	MAG	B	303	3	16,16,16	1.77	4 (25%)	21,22,22	1.39	4 (19%)
3	FUC	B	304	3,4	10,10,11	1.20	1 (10%)	14,14,16	1.03	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SIA	A	301	3	-	0/14/38/38	0/1/1/1
3	GAL	A	302	3	-	0/2/15/22	0/1/1/1
3	MAG	A	303	3	-	0/8/28/28	0/1/1/1
3	FUC	A	304	3,4	-	0/0/17/20	0/1/1/1
3	SIA	B	301	3	-	0/14/38/38	0/1/1/1
3	GAL	B	302	3	-	0/2/15/22	0/1/1/1
3	MAG	B	303	3	-	0/8/28/28	0/1/1/1
3	FUC	B	304	3,4	-	0/0/17/20	0/1/1/1

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	303	MAG	C4-C5	2.00	1.57	1.53
3	B	303	MAG	C3-C2	2.03	1.57	1.53
3	B	301	SIA	C3-C4	2.04	1.56	1.53
3	B	304	FUC	O5-C1	2.07	1.47	1.43
3	A	304	FUC	O5-C1	2.16	1.47	1.43
3	A	303	MAG	C3-C2	2.20	1.57	1.53
3	A	302	GAL	C4-C5	2.34	1.56	1.52
3	B	303	MAG	O5-C1	2.44	1.48	1.41
3	A	303	MAG	O5-C1	2.46	1.48	1.41
3	B	302	GAL	C4-C5	2.54	1.56	1.52
3	A	301	SIA	C11-C10	2.82	1.56	1.50
3	B	302	GAL	C3-C2	2.87	1.57	1.51
3	A	302	GAL	C3-C2	2.89	1.57	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	301	SIA	C7-C6	3.00	1.56	1.52
3	A	301	SIA	C6-C5	3.06	1.58	1.53
3	A	301	SIA	C7-C6	3.11	1.56	1.52
3	B	303	MAG	C1-C2	3.50	1.59	1.53
3	A	303	MAG	C1-C2	3.63	1.59	1.53
3	B	301	SIA	C6-C5	3.69	1.59	1.53
3	B	303	MAG	O1-C1	3.83	1.46	1.40
3	B	301	SIA	O6-C6	3.83	1.50	1.44
3	A	303	MAG	O1-C1	3.86	1.46	1.40
3	A	301	SIA	O6-C6	4.41	1.51	1.44
3	B	301	SIA	C3-C2	4.89	1.58	1.51
3	A	301	SIA	C3-C2	5.13	1.58	1.51
3	A	301	SIA	O6-C2	8.24	1.50	1.42
3	B	301	SIA	O6-C2	8.70	1.50	1.42
3	B	301	SIA	O2-C2	10.41	1.51	1.40
3	A	301	SIA	O2-C2	10.67	1.51	1.40

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	301	SIA	C11-C10-N5	-2.31	111.68	116.11
3	B	303	MAG	C3-C2-N2	-2.17	106.18	110.66
3	A	301	SIA	O10-C10-C11	-2.01	118.38	122.06
3	A	303	MAG	O5-C1-C2	2.04	115.22	110.78
3	A	302	GAL	O5-C1-C2	2.05	113.16	110.85
3	B	303	MAG	CM-O1-C1	2.11	116.74	113.29
3	B	304	FUC	O5-C5-C6	2.14	109.67	106.13
3	B	301	SIA	O6-C6-C7	2.18	110.55	107.26
3	B	303	MAG	O5-C1-C2	2.18	115.52	110.78
3	A	301	SIA	O6-C6-C7	2.19	110.56	107.26
3	A	303	MAG	CM-O1-C1	2.20	116.90	113.29
3	A	304	FUC	O5-C5-C6	2.21	109.79	106.13
3	A	301	SIA	C5-N5-C10	2.26	128.91	123.10
3	B	302	GAL	O5-C1-C2	2.43	113.59	110.85
3	A	303	MAG	C4-C3-C2	2.61	114.05	110.43
3	A	301	SIA	C11-C10-N5	2.66	121.19	116.11
3	B	303	MAG	C4-C3-C2	2.66	114.12	110.43
3	B	301	SIA	O10-C10-N5	2.87	127.71	121.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.6 Ligand geometry

Of 16 ligands modelled in this entry, 2 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	281	1	14,14,15	0.24	0	15,19,21	0.89	1 (6%)
2	NAG	A	282	1	14,14,15	0.30	0	15,19,21	0.50	0
2	NAG	A	283	1	14,14,15	0.37	0	15,19,21	0.66	0
2	NAG	A	284	1	14,14,15	0.33	0	15,19,21	0.87	1 (6%)
2	NAG	A	285	1	14,14,15	0.33	0	15,19,21	0.66	1 (6%)
2	NAG	A	286	1	14,14,15	0.30	0	15,19,21	0.37	0
2	NAG	A	287	1	14,14,15	0.25	0	15,19,21	0.43	0
2	NAG	B	281	1	14,14,15	0.32	0	15,19,21	0.86	1 (6%)
2	NAG	B	282	1	14,14,15	0.33	0	15,19,21	0.47	0
2	NAG	B	283	1	14,14,15	0.39	0	15,19,21	0.56	0
2	NAG	B	284	1	14,14,15	0.36	0	15,19,21	1.11	1 (6%)
2	NAG	B	285	1	14,14,15	0.34	0	15,19,21	0.64	1 (6%)
2	NAG	B	286	1	14,14,15	0.27	0	15,19,21	0.89	1 (6%)
2	NAG	B	287	1	14,14,15	0.27	0	15,19,21	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	281	1	-	0/6/23/26	0/1/1/1
2	NAG	A	282	1	-	0/6/23/26	0/1/1/1
2	NAG	A	283	1	-	0/6/23/26	0/1/1/1
2	NAG	A	284	1	-	0/6/23/26	0/1/1/1
2	NAG	A	285	1	-	0/6/23/26	0/1/1/1
2	NAG	A	286	1	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	287	1	-	0/6/23/26	0/1/1/1
2	NAG	B	281	1	-	0/6/23/26	0/1/1/1
2	NAG	B	282	1	-	0/6/23/26	0/1/1/1
2	NAG	B	283	1	-	0/6/23/26	0/1/1/1
2	NAG	B	284	1	-	0/6/23/26	0/1/1/1
2	NAG	B	285	1	-	0/6/23/26	0/1/1/1
2	NAG	B	286	1	-	0/6/23/26	0/1/1/1
2	NAG	B	287	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	285	NAG	C2-N2-C7	2.06	125.69	123.04
2	B	285	NAG	C2-N2-C7	2.13	125.77	123.04
2	A	284	NAG	C1-O5-C5	2.33	115.20	112.25
2	B	286	NAG	C2-N2-C7	2.58	126.36	123.04
2	B	281	NAG	C1-O5-C5	2.96	116.00	112.25
2	A	281	NAG	C1-O5-C5	3.20	116.31	112.25
2	B	284	NAG	C1-O5-C5	3.52	116.71	112.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	283	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	280/280 (100%)	0.34	14 (5%) 32 32	47, 73, 115, 137	0
1	B	280/280 (100%)	0.27	11 (3%) 43 43	48, 73, 119, 143	1 (0%)
All	All	560/560 (100%)	0.31	25 (4%) 37 37	47, 73, 117, 143	1 (0%)

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	23	TYR	5.6
1	B	250	ASP	5.0
1	B	231	PHE	4.9
1	B	258	MET	4.8
1	B	28	ALA	4.6
1	A	255	PHE	4.5
1	B	235	PHE	4.0
1	B	249	PHE	3.8
1	B	101	VAL	3.5
1	A	252	GLU	3.1
1	A	33	GLU	3.0
1	A	259	GLY	3.0
1	A	28	ALA	2.7
1	A	71	GLU	2.6
1	A	280	ALA	2.4
1	B	260	ALA	2.4
1	A	258	MET	2.4
1	B	29	ILE	2.4
1	A	29	ILE	2.3
1	B	94	TYR	2.3
1	A	77	ALA	2.2
1	A	273	ASN	2.2
1	B	280	ALA	2.1
1	A	25	HIS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	256	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	GAL	A	302	10/12	0.90	0.20	0.83	104,107,110,111	0
3	FUC	A	304	10/11	0.94	0.15	-0.26	93,102,105,106	0
3	GAL	B	302	10/12	0.83	0.17	-0.41	103,106,109,111	0
3	FUC	B	304	10/11	0.96	0.14	-0.53	99,104,106,107	0
3	SIA	A	301	21/21	0.86	0.17	-	105,123,134,136	0
3	MAG	B	303	16/16	0.91	0.18	-	105,107,121,121	0
3	SIA	B	301	21/21	0.80	0.21	-	112,121,130,132	0
3	MAG	A	303	16/16	0.90	0.14	-	109,114,127,129	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NAG	B	285	14/15	0.84	0.25	3.48	124,128,130,130	0
2	NAG	B	284	14/15	0.86	0.20	0.90	68,78,89,90	0
2	NAG	A	285	14/15	0.77	0.23	0.86	105,110,113,115	0
2	NAG	B	283	14/15	0.91	0.13	-0.22	36,49,54,63	0
2	NAG	A	283	14/15	0.94	0.12	-0.29	36,53,57,64	0
2	NAG	A	284	14/15	0.91	0.15	-0.34	69,81,100,103	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	CA	B	306	1/1	0.96	0.08	-1.34	82,82,82,82	0
4	CA	A	306	1/1	0.94	0.09	-1.43	82,82,82,82	0
2	NAG	B	286	14/15	0.55	0.43	-	118,127,133,134	0
2	NAG	A	286	14/15	0.69	0.17	-	107,116,121,122	0
2	NAG	B	282	14/15	0.83	0.30	-	106,109,115,116	0
2	NAG	A	282	14/15	0.88	0.18	-	101,107,112,112	0
2	NAG	B	287	14/15	0.89	0.16	-	89,95,109,112	0
2	NAG	B	281	14/15	0.88	0.14	-	66,76,84,85	0
2	NAG	A	281	14/15	0.96	0.17	-	71,82,90,91	0
2	NAG	A	287	14/15	0.88	0.33	-	89,94,106,112	0

6.5 Other polymers [i](#)

There are no such residues in this entry.