



Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 01:58 PM BST

PDB ID : 1CTD
Title : DETERMINATION OF THE SOLUTION STRUCTURE OF A SYNTHETIC
TWO-SITE CALCIUM-BINDING HOMODIMERIC PROTEIN DOMAIN
BY NMR SPECTROSCOPY
Authors : Shaw, G.S.; Sykes, B.D.
Deposited on : 1992-11-12

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	36	
1	B	36	

2 Ensemble composition and analysis ⓘ

This entry contains 7 models. Model 4 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:7-A:30, B:6-B:30 (49)	0.44	4

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 1 clusters and 2 single-model clusters were found.

Cluster number	Models
1	2, 3, 4, 5, 7
Single-model clusters	1; 6

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 666 atoms, of which 118 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called TROPONIN C SITE III - SITE III HOMODIMER.

Mol	Chain	Residues	Atoms					Trace
1	A	36	Total	C	H	N	O	1
			332	170	59	45	58	
1	B	36	Total	C	H	N	O	1
			332	170	59	45	58	

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	9	ALA	CYS	CONFLICT	UNP P02588
A	20	TYR	PHE	CONFLICT	UNP P02588
B	9	ALA	CYS	CONFLICT	UNP P02588
B	20	TYR	PHE	CONFLICT	UNP P02588

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	
2	B	1	Total	Ca
			1	1
2	A	1	Total	Ca
			1	1

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: TROPONIN C SITE III - SITE III HOMODIMER



- Molecule 1: TROPONIN C SITE III - SITE III HOMODIMER




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: TROPONIN C SITE III - SITE III HOMODIMER



- Molecule 1: TROPONIN C SITE III - SITE III HOMODIMER



4.2.2 Score per residue for model 2

- Molecule 1: TROPONIN C SITE III - SITE III HOMODIMER




- Molecule 1: TROPONIN C SITE III - SITE III HOMODIMER




4.2.3 Score per residue for model 3

- Molecule 1: TROPONIN C SITE III - SITE III HOMODIMER




- Molecule 1: TROPONIN C SITE III - SITE III HOMODIMER



4.2.4 Score per residue for model 4 (medoid)

- Molecule 1: TROPONIN C SITE III - SITE III HOMODIMER




- Molecule 1: TROPONIN C SITE III - SITE III HOMODIMER



4.2.5 Score per residue for model 5

- Molecule 1: TROPONIN C SITE III - SITE III HOMODIMER




- Molecule 1: TROPONIN C SITE III - SITE III HOMODIMER




4.2.6 Score per residue for model 6

- Molecule 1: TROPONIN C SITE III - SITE III HOMODIMER



- Molecule 1: TROPONIN C SITE III - SITE III HOMODIMER



4.2.7 Score per residue for model 7

- Molecule 1: TROPONIN C SITE III - SITE III HOMODIMER



- Molecule 1: TROPONIN C SITE III - SITE III HOMODIMER



5 Refinement protocol and experimental data overview ⓘ

Of the ? calculated structures, 7 were deposited, based on the following criterion: ?.

The authors did not provide any information on software used for structure solution, optimization or refinement.

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ACE, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.90±0.07	0±0/195 (0.1±0.2%)	1.38±0.13	1±1/263 (0.5±0.3%)
1	B	0.84±0.01	0±0/203 (0.0±0.0%)	1.34±0.05	1±1/274 (0.4±0.4%)
All	All	0.87	1/2786 (0.0%)	1.36	18/3759 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	2.4±1.6
1	B	0.0±0.0	0.7±0.5
All	All	0	22

All unique bond outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	16	ASN	CG-OD1	8.85	1.43	1.24	4	1

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	18	ASP	CA-C-N	-10.69	94.82	116.20	1	1
1	A	13	PHE	CB-CG-CD2	-7.17	115.78	120.80	6	2
1	B	20	TYR	CB-CG-CD1	-6.83	116.91	121.00	2	1
1	B	13	PHE	CB-CG-CD2	-6.49	116.26	120.80	2	3
1	A	7	ALA	CB-CA-C	6.49	119.83	110.10	6	1
1	B	11	ARG	NE-CZ-NH1	6.34	123.47	120.30	6	2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	B	20	TYR	CB-CG-CD2	-5.92	117.45	121.00	5	1
1	A	18	ASP	CB-CG-OD2	-5.79	113.09	118.30	1	1
1	A	20	TYR	CB-CG-CD2	-5.74	117.56	121.00	3	1
1	B	10	PHE	CB-CG-CD2	-5.64	116.85	120.80	2	1
1	A	11	ARG	NE-CZ-NH2	-5.44	117.58	120.30	5	1
1	A	20	TYR	CB-CG-CD1	-5.30	117.82	121.00	4	2
1	A	11	ARG	NE-CZ-NH1	5.17	122.88	120.30	2	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	9	ALA	Mainchain	3
1	A	20	TYR	Sidechain	3
1	A	10	PHE	Mainchain	3
1	A	13	PHE	Sidechain	3
1	B	20	TYR	Sidechain	2
1	B	17	ALA	Mainchain	2
1	A	7	ALA	Mainchain	1
1	A	11	ARG	Sidechain	1
1	B	7	ALA	Peptide	1
1	A	17	ALA	Mainchain	1
1	A	18	ASP	Mainchain,Peptide	1

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	192	37	182	1±2
1	B	200	38	193	1±1
All	All	2758	525	2625	11

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:18:ASP:HA	1:A:20:TYR:H	0.80	1.32	1	1
1:A:18:ASP:HA	1:A:20:TYR:N	0.68	2.02	1	1
1:A:20:TYR:HB3	1:B:20:TYR:HB3	0.60	1.74	3	1
1:B:11:ARG:HH12	1:B:19:GLY:HA2	0.50	1.65	3	1
1:A:26:LEU:HA	1:A:29:ILE:HD12	0.49	1.83	1	1
1:A:10:PHE:O	1:A:14:ASP:HB2	0.48	2.07	1	1
1:A:14:ASP:OD1	1:A:21:ILE:HG13	0.48	2.08	1	1
1:A:7:ALA:O	1:A:10:PHE:HB3	0.46	2.10	6	1
1:B:8:ASN:ND2	1:B:8:ASN:H	0.44	2.09	5	1
1:B:11:ARG:NH1	1:B:19:GLY:HA2	0.42	2.28	3	1
1:A:8:ASN:O	1:A:12:ILE:HG12	0.41	2.15	4	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	24/36 (67%)	22±0 (92±1%)	2±0 (7±2%)	0±0 (1±2%)	21	68
1	B	25/36 (69%)	23±1 (91±4%)	2±1 (7±4%)	0±0 (1±2%)	23	69
All	All	343/504 (68%)	315 (92%)	24 (7%)	4 (1%)	21	68

All 4 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	B	29	ILE	1
1	A	29	ILE	1
1	B	14	ASP	1
1	A	7	ALA	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation

was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	19/27 (70%)	17±1 (87±5%)	2±1 (13±5%)	10	51
1	B	20/27 (74%)	17±1 (86±3%)	3±1 (14±3%)	8	48
All	All	273/378 (72%)	236 (86%)	37 (14%)	9	49

All 14 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	22	ASP	7
1	B	22	ASP	7
1	B	24	GLU	4
1	A	30	LEU	3
1	B	11	ARG	3
1	A	24	GLU	3
1	B	26	LEU	2
1	B	6	LEU	2
1	A	14	ASP	1
1	A	15	LYS	1
1	B	30	LEU	1
1	A	11	ARG	1
1	A	21	ILE	1
1	B	21	ILE	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided