



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 06:53 PM GMT

PDB ID : 1CX2
Title : CYCLOOXYGENASE-2 (PROSTAGLANDIN SYNTHASE-2) COM-
PLEXED WITH A SELECTIVE INHIBITOR, SC-558
Authors : Kurumbail, R.; Stallings, W.
Deposited on : 1996-12-17
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	587	
1	B	587	
1	C	587	
1	D	587	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	661	-	-	-	X
2	NAG	B	661	-	-	-	X
2	NAG	B	681	-	-	-	X
2	NAG	C	661	-	-	-	X
2	NAG	D	661	-	-	-	X
2	NAG	D	681	-	-	-	X
4	S58	A	701	-	-	X	-
4	S58	C	701	-	-	X	-
4	S58	D	701	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 22376 atoms, of which 4040 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYCLOOXYGENASE-2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	552	Total	C	H	N	O	S	0	0	0
			5439	2886	966	748	814	25			
1	B	552	Total	C	H	N	O	S	0	0	0
			5439	2886	966	748	814	25			
1	C	552	Total	C	H	N	O	S	0	0	0
			5439	2886	966	748	814	25			
1	D	552	Total	C	H	N	O	S	0	0	0
			5439	2886	966	748	814	25			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	310	GLN	ASN	CONFLICT	UNP Q05769
A	333	LYS	ARG	CONFLICT	UNP Q05769
B	310	GLN	ASN	CONFLICT	UNP Q05769
B	333	LYS	ARG	CONFLICT	UNP Q05769
C	310	GLN	ASN	CONFLICT	UNP Q05769
C	333	LYS	ARG	CONFLICT	UNP Q05769
D	310	GLN	ASN	CONFLICT	UNP Q05769
D	333	LYS	ARG	CONFLICT	UNP Q05769

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	A	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	B	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	B	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	B	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	C	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	C	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	C	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	D	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	D	1	Total	C	H	N	O	0	0
			28	8	14	1	5		
2	D	1	Total	C	H	N	O	0	0
			28	8	14	1	5		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 4 is 1-PHENYLSULFONAMIDE-3-TRIFLUOROMETHYL-5-PARABROMOPHENYLPYRAZOLE (three-letter code: S58) (formula: $C_{16}H_{11}BrF_3N_3O_2S$).




Mol	Chain	Residues	Atoms								ZeroOcc	AltConf
4	A	1	Total	Br	C	F	H	N	O	S	0	0
			28	1	16	3	2	3	2	1		
4	B	1	Total	Br	C	F	H	N	O	S	0	0
			28	1	16	3	2	3	2	1		
4	C	1	Total	Br	C	F	H	N	O	S	0	0
			28	1	16	3	2	3	2	1		
4	D	1	Total	Br	C	F	H	N	O	S	0	0
			28	1	16	3	2	3	2	1		


3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CYCLOOXYGENASE-2




• Molecule 1: CYCLOOXYGENASE-2





Chain C:



LYS
THR
ALA
THR
ILE
ASN
ALA
SER
ALA
SER
ILE
HIS
SER
ARG
LEU
ASP
S49
ILE
ASN
PRO
THR
VAL
LEU
ILE
LYS
ARG
ARG
SER
THR
GLU
LEU

• Molecule 1: CYCLOOXYGENASE-2

Chain D: 31% 52% 11% 6%

L494	G432	L365	M299	D239	K166	M101	A33
Y495	R433	L366	Y300	R240	L171	N104	N34
S496	V434	F367	X301	Q241	P172	M105	G36
D497	A435		A302	E242	D173	I105A	C37
L498		Q374	T303	K243	S174	P106	S38
D499	R438	R375	I304	L244	E175	F107	N39
V500		R376	X305	R245	K176	L108	P40
M501	P441	L377	L306	L246	E176	R109	Q41
E502	L442	S579	R307	F247	V177	S110	C42
S503	A443	S379	E308	K248	L178	N111	N43
L504	V444	E380	H309	D249	E179	L112	R44
P505	Q445	F381	Q310	G250	K180	M112	
A506	A446	L382	R311	R251	V181	K113	C47
ASP	V447	T383	V312	L252	L182	K114	M48
PRO	A448	L384	C313	K253	L183	Y115	S49
GLN	K449	Y385	D314	Q254	R184	V116	T50
PRO	A450	H386	L315	Q255		L117	
THR	S451	N387	L316	V256	F187		G51
	L452	H388	K317	I257	I188	R120	F52
THR	D453	P389	Q318	G258	P189	S121	D53
ALA	Q454	L390	E319	G259	D190	Y122	Q54
THR	S455	L391	H320	E260	P191	L123	Y55
ILE	F518		P321	V261	Q192	I124	K56
ASN	G519		E322	Y262	G193	D125	C57
ALA	E520		K323	P263	S194	S126	
SER	T521		G324	T264	M195	P127	F64
M522	K459	N396	D325	T265	M196	P128	Y65
SER	Q461	L397	E326	V266	M197		G66
V523	S462	E398	Q327	K267	F198	M131	E67
E524	L463	D399	L328	D268		V132	N68
SER	M464	Q400	F329	T269	Q203	H133	C69
ARG	E465	E401		T270	H204	Y134	T70
ASP	Y466	Y402	K333	V271	F205	G135	P72
ASP	R467		L334	E272	T206	Y136	T71
ILE	K468		I335	M273	H207	K137	E73
ASN	R469	Q406	L336	L274	Q208	S138	F74
PRO	F470		I337	I275	F209		L75
THR	S471	L408	G338	P276	F210	A141	T76
VAL	L472	Y409		P277	K211	F142	R77
LEU	K473	N410	K342	E278	T212	S143	I78
P542	P474	N411	I343	L279	D213	M144	K79
Q543	Y475	S412	V344	P280	H214	L145	L80
LYS	T476	L413	I345	E281	K215	S146	L81
ARG	F477	L414	E346	N282	R216		L82
ARG	F478		D347	L283		T149	K83
THR	E479	H417	Y348	Q284	G219	R150	P84
SER	L480	G418	V349	F285	F220	A151	T85
GLU	E481	L419	Q350	A286		L152	P86
LEU	T482	T420	H351	V287	R226	P153	N87
	G483	Q421	L352	G288	G227	P154	T88
V554	E484	F422	S353	Q289	V228	V155	
G555	K485	V423	E354	E290	D229	A156	I92
F556	E486	E424	Y355	V291	L230	D157	L93
K557	N487	S425	H356	F292	N231	D158	T94
L558	A488	F426	F357	G293	I232	C159	H95
L559	A489	T427	K358	L294	I233	P160	F96
	E490	L427	L359	V295	Y234	T161	K97
L564	L491	Q429	K360	P296	G235	P162	G98
	K492	L430		G297		Y69	
V568	A493		F264	T298		Y165	H100
C569							

4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	180.34Å 133.92Å 121.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 3.00 20.00 – 3.00	Depositor EDS
% Data completeness (in resolution range)	54.0 (8.00-3.00) 61.9 (20.00-3.00)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.66 (at 2.98Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.216 , 0.218 0.226 , 0.317	Depositor DCC
R_{free} test set	3411 reflections (11.13%)	DCC
Wilson B-factor (Å ²)	28.9	Xtriage
Anisotropy	0.960	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 62.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.38$, $\langle L^2 \rangle = 0.20$	Xtriage
Outliers	0 of 36581 reflections	Xtriage
F_o, F_c correlation	0.84	EDS
Total number of atoms	22376	wwPDB-VP
Average B, all atoms (Å ²)	9.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 39.14 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 3.3164e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, NAG, S58

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	0/4600	0.88	4/6237 (0.1%)
1	B	0.71	0/4600	0.88	4/6237 (0.1%)
1	C	0.69	0/4600	0.88	3/6237 (0.0%)
1	D	0.72	0/4600	0.89	3/6237 (0.0%)
All	All	0.71	0/18400	0.88	14/24948 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	2
1	C	0	2
1	D	0	4
All	All	0	11

There are no bond length outliers.

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	575	CYS	N-CA-C	-6.66	93.02	111.00
1	D	575	CYS	N-CA-C	-6.47	93.54	111.00
1	C	575	CYS	N-CA-C	-6.38	93.78	111.00
1	B	575	CYS	N-CA-C	-6.35	93.85	111.00
1	D	287	VAL	N-CA-C	5.94	127.03	111.00

There are no chirality outliers.

5 of 11 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	134	TYR	Sidechain
1	A	262	TYR	Sidechain
1	A	348	TYR	Sidechain
1	B	262	TYR	Sidechain
1	B	348	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4473	966	4375	399	0
1	B	4473	966	4375	409	0
1	C	4473	966	4375	389	0
1	D	4473	966	4375	392	0
2	A	42	42	39	1	0
2	B	42	42	39	6	0
2	C	42	42	39	1	0
2	D	42	42	39	7	0
3	A	43	0	30	3	0
3	B	43	0	30	6	0
3	C	43	0	30	7	0
3	D	43	0	30	5	0
4	A	26	2	11	9	0
4	B	26	2	11	5	0
4	C	26	2	11	11	0
4	D	26	2	11	7	0
All	All	18336	4040	17820	1552	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 43.

The worst 5 of 1552 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:52:PHE:N	1:D:322:GLU:HG2	1.80	0.96
1:C:322:GLU:HG2	1:D:52:PHE:N	1.83	0.93
1:A:275:TYR:CE2	1:A:284:GLN:HA	2.04	0.93

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:184:ARG:HA	1:C:438:ARG:O	1.71	0.89
1:C:275:TYR:CE2	1:C:284:GLN:HA	2.07	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	550/587 (94%)	439 (80%)	86 (16%)	25 (4%)	3	18
1	B	550/587 (94%)	438 (80%)	85 (16%)	27 (5%)	3	16
1	C	550/587 (94%)	439 (80%)	84 (15%)	27 (5%)	3	16
1	D	550/587 (94%)	444 (81%)	78 (14%)	28 (5%)	2	15
All	All	2200/2348 (94%)	1760 (80%)	333 (15%)	107 (5%)	3	16

5 of 107 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	67	GLU
1	A	138	SER
1	A	226	HIS
1	A	270	GLN
1	A	284	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	493/525 (94%)	422 (86%)	71 (14%)	4	18
1	B	493/525 (94%)	411 (83%)	82 (17%)	3	13
1	C	493/525 (94%)	423 (86%)	70 (14%)	4	19
1	D	493/525 (94%)	418 (85%)	75 (15%)	3	16
All	All	1972/2100 (94%)	1674 (85%)	298 (15%)	3	17

5 of 298 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	428	ARG
1	C	150	ARG
1	D	412	SER
1	B	469	ARG
1	B	577	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 71 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	351	HIS
1	C	87	ASN
1	D	318	GLN
1	B	382	ASN
1	B	464	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

20 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	661	1	14,14,15	0.73	0	15,19,21	0.92	1 (6%)
2	NAG	A	671	1	14,14,15	0.54	0	15,19,21	1.22	2 (13%)
2	NAG	A	681	1	14,14,15	0.80	0	15,19,21	0.93	1 (6%)
3	HEM	A	682	1	30,50,50	2.83	9 (30%)	24,82,82	2.02	6 (25%)
4	S58	A	701	-	27,28,28	2.49	6 (22%)	40,43,43	1.69	7 (17%)
2	NAG	B	661	1	14,14,15	0.61	0	15,19,21	0.95	1 (6%)
2	NAG	B	671	1	14,14,15	0.59	0	15,19,21	0.95	1 (6%)
2	NAG	B	681	1	14,14,15	0.67	0	15,19,21	0.67	0
3	HEM	B	682	1	30,50,50	3.19	10 (33%)	24,82,82	2.08	7 (29%)
4	S58	B	701	-	27,28,28	2.60	8 (29%)	40,43,43	1.70	9 (22%)
2	NAG	C	661	1	14,14,15	0.90	0	15,19,21	0.82	1 (6%)
2	NAG	C	671	1	14,14,15	0.87	1 (7%)	15,19,21	1.20	2 (13%)
2	NAG	C	681	1	14,14,15	0.85	0	15,19,21	0.72	0
3	HEM	C	682	1	30,50,50	2.85	7 (23%)	24,82,82	2.02	6 (25%)
4	S58	C	701	-	27,28,28	2.61	8 (29%)	40,43,43	1.53	7 (17%)
2	NAG	D	661	1	14,14,15	0.61	0	15,19,21	0.84	1 (6%)
2	NAG	D	671	1	14,14,15	0.60	0	15,19,21	1.26	2 (13%)
2	NAG	D	681	1	14,14,15	0.54	0	15,19,21	0.85	0
3	HEM	D	682	1	30,50,50	2.90	9 (30%)	24,82,82	2.01	6 (25%)
4	S58	D	701	-	27,28,28	2.70	9 (33%)	40,43,43	1.69	9 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	661	1	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	671	1	-	0/6/23/26	0/1/1/1
2	NAG	A	681	1	-	0/6/23/26	0/1/1/1
3	HEM	A	682	1	-	0/10/54/54	0/0/8/8
4	S58	A	701	-	-	0/20/20/20	0/3/3/3
2	NAG	B	661	1	-	0/6/23/26	0/1/1/1
2	NAG	B	671	1	-	0/6/23/26	0/1/1/1
2	NAG	B	681	1	-	0/6/23/26	0/1/1/1
3	HEM	B	682	1	-	0/10/54/54	0/0/8/8
4	S58	B	701	-	-	0/20/20/20	0/3/3/3
2	NAG	C	661	1	-	0/6/23/26	0/1/1/1
2	NAG	C	671	1	-	0/6/23/26	0/1/1/1
2	NAG	C	681	1	-	0/6/23/26	0/1/1/1
3	HEM	C	682	1	-	0/10/54/54	0/0/8/8
4	S58	C	701	-	-	0/20/20/20	0/3/3/3
2	NAG	D	661	1	-	0/6/23/26	0/1/1/1
2	NAG	D	671	1	-	0/6/23/26	0/1/1/1
2	NAG	D	681	1	-	0/6/23/26	0/1/1/1
3	HEM	D	682	1	-	0/10/54/54	0/0/8/8
4	S58	D	701	-	-	0/20/20/20	0/3/3/3

The worst 5 of 67 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	682	HEM	C3B-C4B	-9.69	1.43	1.51
3	C	682	HEM	C3B-C4B	-7.55	1.45	1.51
3	B	682	HEM	C3B-CAB	-7.43	1.37	1.51
3	D	682	HEM	C3B-C4B	-6.94	1.45	1.51
3	D	682	HEM	C3B-CAB	-6.83	1.38	1.51

The worst 5 of 69 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	701	S58	O2-S1-O1	-6.47	109.71	118.80
4	B	701	S58	O2-S1-O1	-6.25	110.02	118.80
4	D	701	S58	O2-S1-O1	-5.99	110.38	118.80
4	C	701	S58	O2-S1-O1	-5.16	111.55	118.80
2	D	671	NAG	C2-N2-C7	-3.46	118.59	123.04

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

14 monomers are involved in 68 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	671	NAG	1	0
3	A	682	HEM	3	0
4	A	701	S58	9	0
2	B	661	NAG	2	0
2	B	681	NAG	4	0
3	B	682	HEM	6	0
4	B	701	S58	5	0
2	C	661	NAG	1	0
3	C	682	HEM	7	0
4	C	701	S58	11	0
2	D	661	NAG	5	0
2	D	681	NAG	2	0
3	D	682	HEM	5	0
4	D	701	S58	7	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	552/587 (94%)	-0.53	0 100 100	2, 6, 16, 30	0
1	B	552/587 (94%)	-0.56	1 (0%) 95 87	2, 6, 16, 27	0
1	C	552/587 (94%)	-0.53	0 100 100	2, 6, 16, 26	0
1	D	552/587 (94%)	-0.50	1 (0%) 95 87	2, 7, 17, 30	0
All	All	2208/2348 (94%)	-0.53	2 (0%) 95 90	2, 6, 16, 30	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	105	ASN	2.6
1	B	583	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NAG	C	661	14/15	0.80	0.31	6.62	12,17,21,22	0
2	NAG	A	661	14/15	0.80	0.30	5.08	12,17,20,21	0
2	NAG	D	681	14/15	0.84	0.43	4.83	12,17,20,24	0
2	NAG	D	661	14/15	0.88	0.31	4.55	15,17,19,21	0
2	NAG	B	681	14/15	0.80	0.30	2.63	11,17,22,24	0
2	NAG	B	661	14/15	0.89	0.30	2.61	15,17,19,20	0
2	NAG	C	681	14/15	0.85	0.29	1.93	11,17,21,22	0
3	HEM	B	682	43/43	0.94	0.18	0.51	2,3,7,9	0
2	NAG	A	681	14/15	0.89	0.23	0.32	7,17,18,21	0
3	HEM	D	682	43/43	0.94	0.17	0.07	2,3,7,9	0
3	HEM	A	682	43/43	0.94	0.17	-0.06	2,3,8,10	0
2	NAG	A	671	14/15	0.93	0.16	-0.31	4,12,17,17	0
2	NAG	D	671	14/15	0.92	0.15	-0.33	5,11,17,17	0
2	NAG	C	671	14/15	0.92	0.15	-0.47	7,13,17,17	0
2	NAG	B	671	14/15	0.93	0.17	-0.50	5,12,17,17	0
4	S58	C	701	26/26	0.97	0.13	-0.71	2,8,15,17	0
4	S58	A	701	26/26	0.98	0.13	-0.78	2,9,15,17	0
4	S58	D	701	26/26	0.97	0.12	-0.90	2,8,13,17	0
3	HEM	C	682	43/43	0.96	0.14	-0.91	2,3,7,8	0
4	S58	B	701	26/26	0.98	0.12	-1.45	2,8,14,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.